



Taylor & Francis  
Taylor & Francis Group



---

Sums of Reciprocals of Integers Missing a Given Digit

Author(s): Robert Baillie

Source: *The American Mathematical Monthly*, Vol. 86, No. 5 (May, 1979), pp. 372-374

Published by: Taylor & Francis, Ltd. on behalf of the Mathematical Association of America

Stable URL: <https://www.jstor.org/stable/2321096>

Accessed: 23-09-2024 18:26 UTC

---

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

Your use of the JSTOR archive indicates your acceptance of the Terms & Conditions of Use, available at <https://about.jstor.org/terms>



JSTOR

*Taylor & Francis, Ltd., Mathematical Association of America* are collaborating with JSTOR to digitize, preserve and extend access to *The American Mathematical Monthly*

2. T. J. P. A. Bromwich, *An Introduction to the Theory of Infinite Series*, London, 1931.
3. Harold T. Davis, *The Theory of Linear Operators from the Standpoint of Differential Equations of Infinite Order*, Principia Press, Bloomington, Ind., 1936.
4. A. Erdélyi, *Asymptotic Expansions*, Dover, New York, 1965.
5. G. H. Hardy, *Divergent Series*, Oxford, 1949.
6. Morris Kline, *Mathematical Thought from Ancient to Modern Times*, Oxford, New York, 1972.
7. L. M. Milne-Thomson, *The Calculus of Finite Differences*, London, 1933.
8. N. E. Nörlund, *Leçons sur les Séries d'Interpolation*, Paris, 1926.
9. F. W. J. Olver, *Asymptotics and Special Functions*, Academic, New York, 1974.
10. H. S. Wall, *Analytic Theory of Continued Fractions*, Van Nostrand, 1948.
11. D. V. Widder, *The Laplace Transform*, Princeton, 1941.
12. L. Euler, *De seriebus divergentibus*, *Novi. Comm. Acad. Sci. Petrop.*, 5 (1754/55) 19–23, 205–237 = *Opera Omnia* (1) 14, 585–617. An English translation and paraphrase by E. J. Barbeau and P. J. Leah appears in *Historia Mathematica*, 3 (1976) 141–160.
13. L. Euler, *Opera Postuma, Mathematica et Physica*, 1 (ed. P. H. Fuss and N. Fuss) St. Petersburg, 1862, Six letters to N. Bernoulli, 519–549.
14. C. I. Gerhardt, ed., G. W. Leibniz *Mathematische Schriften* 4, Olms, Hildesheim, New York, 1971.
15. Judith V. Grabiner, Is mathematical truth time-dependent? this MONTHLY, 81 (1974) 354–365.
16. G. H. Hardy, Note on a divergent series, *Proc. Cambridge Philos. Soc.*, 37 (1941) 1–8.
17. Howard K. Hughes, On the analytical extension of functions defined by factorial series, *Amer. J. Math.*, 53 (1931) 757–780.
18. E. Laguerre, Sur l'intégrale  $\int_0^\infty e^{-t}/t dt$ , *Bull. Soc. Math. France*, 7 (1879) 72–81.
19. E. Laguerre, Sur la réduction en fractions continues d'une fonction qui satisfait une équation différentielle linéaire du premier ordre dont les coefficients sont rationnels, *Jour. de Math.* (4), 1 (1885) 135–165.
20. C. N. Moore, Summability of series, this MONTHLY, 39 (1932) 62–71 = Selected papers on calculus (MAA, 1968) 333–341.
21. T. J. Stieltjes, Recherches sur quelques séries semi-convergentes. *Ann. Sci. École Norm. Sup.* (3), 3 (1886) 201–258 = *Oeuvres* 2, 2–58.
22. T. J. Stieltjes, Recherche sur les fractions continues, *Ann. Fac. Sci. Univ. Toulouse*, 8 (1894) J 1–122; 9 (1895) A 1–47 = *Oeuvres* 2, 402–566.
23. John Tucciarone, The development of the theory of summable divergent series from 1880 to 1925, *Arch. History Exact Sci.*, 10 (1973) 1–40.
24. A. P. Youschkevitch, The concept of function up to the middle of the 19th century. *Arch. History Exact Sci.*, 16 (1976) 37–85.
25. W. T. Reid, A note on the Hamburger and Stieltjes moment problems, *Proc. Amer. Math. Soc.*, 5 (1954) 521–525.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TORONTO, TORONTO, ONTARIO, CANADA M5S 1A1.

## MATHEMATICAL NOTES

EDITED BY DEBORAH TEPPER HAIMO AND FRANKLIN TEPPER HAIMO

*Material for this department should be sent to Deborah Tepper Haimo, Department of Mathematical Sciences, University of Missouri, St. Louis, MO 63121.*

### SUMS OF RECIPROCAL OF INTEGERS MISSING A GIVEN DIGIT

ROBERT BAILLIE

The harmonic series  $\sum_{n=1}^{\infty} 1/n$  diverges. If we omit those terms for which  $n$  has, say, at least one "9" in its base 10 representation, then the remaining series converges [6]. In fact, this result holds for any base  $b \geq 2$  and any digit  $m$ ,  $0 \leq m \leq b-1$ . (See [4, Theorem 144, p. 120].) Various

estimates for these sums are obtained in [1]–[7], although nowhere is a method discussed that easily produces accurate values of the sums.

This note gives a simple technique for accurately summing these series. We also give, to twenty decimal places, the values of the ten sums where the “missing digit” ranges from 0 through 9, base 10. (See Table 1.) This method easily generalizes to other bases.

Choose the missing digit  $m$ ,  $0 \leq m \leq 9$ . Let  $S$  be the set of positive integers that have no digit equal to  $m$ , and let  $S_i$  be the set of integers in  $S$  which have exactly  $i$  digits. Define

$$s(i, j) = \sum_{x \in S_i} 1/x^j \quad (j \geq 1). \quad (1)$$

The sum we are interested in is

$$t_m = \sum_{i=1}^{\infty} s(i, 1), \quad (2)$$

the sum of the reciprocals of all positive integers missing the digit  $m$ .

Now  $t_m$  cannot be computed from (2) because the convergence is much too slow: the remainder after summing  $10^{27}$  terms still exceeds one! We can, however, derive a recursion formula which gives  $s(i+1, j)$  from  $s(i, j+n)$ ,  $n \geq 0$ , and for small  $i$ ,  $s(i, j+n)$  can be computed directly.

The recursion formula is based on the observation that

$$S_{i+1} = \bigcup_{x \in S_i} [\{10x, 10x+1, \dots, 10x+9\} \setminus \{10x+m\}].$$

Therefore,

$$s(i+1, j) = \sum_x \left[ \sum'_k (10x+k)^{-j} \right] \quad (3)$$

where the outer sum extends over all  $x$  in  $S_i$ , and  $\Sigma'$  is over  $k=0, 1, \dots, 9, k \neq m$ .

Let us make these definitions:

$$\begin{aligned} c(j, n) &= (-1)^n \binom{j+n-1}{n} \quad (j > 0, n \geq 0) \\ b_n &= 0^n + 1^n + \dots + 9^n - m^n \quad (n > 0), b_0 = 9 \\ a(j, n) &= b_n c(j, n) / 10^{j+n}. \end{aligned}$$

Next, rearrange (3) using the binomial theorem to expand  $(10x+k)^{-j}$ :

$$\begin{aligned} s(i+1, j) &= \sum_x \left\{ (10x)^{-j} \sum'_k \left[ \sum_{n=0}^{\infty} c(j, n) (k/10x)^n \right] \right\} \\ &= \sum_x \left\{ (10x)^{-j} \left[ \sum_{n=0}^{\infty} b_n c(j, n) (10x)^{-n} \right] \right\} \\ &= \sum_x \left[ \sum_{n=0}^{\infty} a(j, n) x^{-(j+n)} \right]. \end{aligned}$$

According to (1), we can write this last sum as

$$s(i+1, j) = \sum_{n=0}^{\infty} a(j, n) s(i, j+n). \quad (4)$$

The needed  $s(i, j)$  for  $i \leq 4$  can be computed explicitly. (4) is then used with at most 10 terms to get  $s(i, 1)$  for  $5 \leq i \leq 30$ . For  $i \geq 31$ , we use the estimate

$$\sum_{i=31}^{\infty} s(i, 1) \sim 9 \cdot s(30, 1).$$

This comes from using only the first term of (4). It can be shown that the round-off and truncation errors will not affect the twentieth decimal place of  $t_m$ .

It seems plausible that the  $t_m$  are irrational.

TABLE 1.

| $m$ |          | $t_m$ |       |       |
|-----|----------|-------|-------|-------|
| 0   | 23.10344 | 79094 | 20541 | 61603 |
| 1   | 16.17696 | 95281 | 23444 | 26657 |
| 2   | 19.25735 | 65328 | 08072 | 22453 |
| 3   | 20.56987 | 79509 | 61230 | 37107 |
| 4   | 21.32746 | 57995 | 90036 | 68663 |
| 5   | 21.83460 | 08122 | 96918 | 16340 |
| 6   | 22.20559 | 81595 | 56091 | 88416 |
| 7   | 22.49347 | 53117 | 05945 | 39817 |
| 8   | 22.72636 | 54026 | 79370 | 60283 |
| 9   | 22.92067 | 66192 | 64150 | 34816 |

References

1. R. Alexander, Remarks about the digits of integers, *J. Austral. Math. Soc.*, 12 (1971) 239–241.  
2. B. D. Craven, On digital distribution in some integer sequences, *J. Austral. Math. Soc.*, 5 (1965) 325–330.  
3. N. J. Fine, Solution to problem E 2204, this MONTHLY, 77 (1970) 1009–1010.  
4. G. H. Hardy and E. M. Wright, *An Introduction to the Theory of Numbers*, Oxford University Press, 1965.  
5. F. Irwin, A curious convergent series, this MONTHLY, 23 (1916) 149–152.  
6. A. J. Kempner, A curious convergent series, this MONTHLY, 21 (1914) 48–50.  
7. A. D. Wadhwa, An interesting subseries of the harmonic series, this MONTHLY, 82 (1975) 931–933.

COMPUTER-BASED EDUCATION RESEARCH LAB, UNIVERSITY OF ILLINOIS AT URBANA-CHAMPAIGN, URBANA, IL 61801.

A GENERALIZATION OF ZOLOTAREV'S THEOREM

PATRICK MORTON

In 1872 Zolotarev [4] published the following theorem:

**THEOREM 1.** *Let  $p$  be an odd prime and let  $a$  be an integer not divisible by  $p$ . Let  $\pi_a$  be the permutation on the group of reduced residues mod  $p$  given by*

$$\pi_a(k) \equiv ak \pmod{p}, \quad k = 1, \dots, p-1.$$

*Then  $\pi_a$  is an even or an odd permutation according as  $a$  is a quadratic residue or nonresidue mod  $p$ ; in symbols*

$$\operatorname{sgn} \pi_a = \left( \frac{a}{p} \right),$$

*where  $\operatorname{sgn}$  is the signum of the permutation and  $\left( \frac{a}{p} \right)$  is the Legendre symbol.*

Several proofs of this theorem have been given [1], [2], [3]. A group-theoretic proof can be constructed by considering the map  $a \rightarrow \operatorname{sgn} \pi_a$  as a homomorphism from the group of reduced residues onto the group  $\{\pm 1\}$  and showing that the kernel is exactly the subgroup of squares mod  $p$ . This argument leads to the following generalization.

**THEOREM 2.** *Let  $G$  be a finite group, and let  $G^+$  be the subgroup of  $G$  generated by the squares in  $G$ . For  $a$  in  $G$  let  $\pi_a$  be the permutation on  $G$  given by*

$$\pi_a(k) = ka, \quad k \text{ in } G.$$

*Then if the order of  $G$  is even and its 2-Sylow subgroups are cyclic,*

The author is an NSF Predoctoral Fellow.