Peaks or thresholds?

Non-Gaussian primordial black hole abundances for broad spectra and the pulsar-timing-array tension

[Author name]*

Abstract

The abundance of primordial black holes (PBHs) that accompanies an enhanced curvature spectrum is computed either with threshold statistics or with peaks theory. With local non-Gaussianity (NG), peaks theory had been applied only to monochromatic spectra or with the peak condition imposed on a Gaussian field. We construct peaks theory on the compaction function for broad spectra and local NG as the exact counterpart of threshold statistics, with the same threshold, window, transfer function and cut-off. The maxima of the non-Gaussian linear compaction $C_\ell = F'(\zeta_r)C_\mathrm{g}$ are the BBKS maxima of the Gaussian field $C_\mathrm{g} + \lambda\zeta_r$ with $\lambda = C_\ell F''/F'^2$, times a rank-one Hessian factor that we compute exactly to first order. An independent Monte Carlo and counts of maxima on 256^3 Gaussian random fields confirm the result; the Gaussian-peak approximation used so far miscounts peaks by up to 18%. The required amplitude ratio $R = A_*^\mathrm{peaks}/A_*^\mathrm{threshold}$ is 0.74–0.82 for $|f_\mathrm{NL}| \leq 2$ and lognormal widths $0.1 \leq \Delta \leq 2$. The NG dependence cancels almost completely in R/R_Gauss, to within $\pm 4\%$ (16% for a strong ultra-slow-roll tail). Propagated to the NANOGrav 15-year data, peaks theory strengthens the exclusion of a scalar-induced gravitational-wave explanation by 0.2–0.8σ (0.4σ in the BBKS normalisation), for every NG template and spectral shape we consider. This is smaller than the effect of NG itself for $|f_\mathrm{NL}| \gtrsim 1$. It is comparable to three analysis choices that are rarely stated:

- the transfer function in the collapse threshold;
- the normalisation of f_PBH (the expression in common use overestimates f_PBH by a factor of 3–13);
- the frequency range of the PTA fit.

We quantify all three.

1 Introduction

Pulsar timing arrays (PTAs) have found evidence for a nanohertz gravitational-wave background [1]. One cosmological interpretation is the gravitational waves induced at second order by a peak in the primordial curvature spectrum [3, 4, 5, 7, 2]. A peak large enough to produce the signal also forms PBHs when the perturbations re-enter the horizon. Requiring that PBHs are not overproduced disfavours the scenario at the 2–3σ level [8, 2, 9]. How strongly depends on the NG of the curvature perturbation and on how the PBH abundance is computed.

Two prescriptions are in use for the abundance:

- **Threshold statistics** counts the volume fraction in which the compaction function exceeds a threshold. It includes the non-linear relation between curvature and density and any local NG non-perturbatively [12, 13].
- **Peaks theory** counts maxima [14]. It has been developed for Gaussian curvature with finite width [15, 16, 17] and for local NG with monochromatic spectra [18, 19, 20].

*[Affiliation, e-mail]. Draft of 24 October 2026; the code `ngpbh` and every number below are in the accompanying repository.

Several recent papers state that peaks theory with local NG for broad spectra is an open problem [13, 9, 10, 15, 17]. 2306.17149 [8] add, in a footnote, that peaks theory "provides slightly smaller amplitudes", so their threshold-statistics bounds are conservative.

This note fills that gap and quantifies the footnote.

- **Construction** (Sec. 2). We build peaks theory on the compaction function for broad spectra with local NG (quadratic, logarithmic or ultra-slow-roll, and curvaton templates). It is the exact counterpart of threshold statistics, so that the two differ only by the peak condition.
- **Main technical step.** We count maxima of the non-Gaussian compaction itself, a local function of two correlated Gaussian fields, rather than of its Gaussian component. This differs from the closest existing work [21, 22, 23].
- **The gap** (Sec. 4). We give the amplitude ratio $R(\Delta, \mathrm{NG})$ between the statistics.
- **PTA consequences** (Sec. 5). We propagate the ratio to the significance of the PBH-overproduction tension for NANOGrav 15, with every analysis choice stated and varied.
- **Two pitfalls.** The second appears widely in the literature.
  - 2306.17149 compute the collapse threshold without the radiation transfer function, although their formula includes it.
  - The $f_{\rm PBH}$ normalisation in common use is too large by a factor of 3–13.

2 Two statistics for the non-Gaussian compaction

2.1 Local NG and the compaction function

The curvature perturbation is $\zeta = F(\zeta_{\rm g})$ with $\zeta_{\rm g}$ Gaussian and power spectrum $\mathcal{P}_\zeta$. We use three templates:

- quadratic, $F = \zeta_{\rm g} + \frac{3}{5} f_{\rm NL} \zeta_{\rm g}^2$;
- ultra-slow-roll (USR), $F = -\mu_* \ln(1 - \zeta_{\rm g}/\mu_*)$, which is $F = -\frac{2}{\beta}\ln(1 - \frac{\beta}{2}\zeta_{\rm g})$ of 2306.17149 with $\mu_* = 2/\beta$ and has $f_{\rm NL} = 5/(6\mu_*)$ at second order;
- the sudden-decay curvaton, $\zeta = \ln X(r_{\rm dec}, \zeta_{\rm g})$ [12].

At smoothing scale r (the scale that re-enters the horizon at formation, $k = 1/r$), define the Gaussian fields

$$C_{\rm g}(\boldsymbol{x}; r) = \int \frac{\mathrm{d}^3 k}{(2\pi)^3} e^{i\boldsymbol{k}\cdot\boldsymbol{x}} \tfrac{2\Phi}{3} (kr)^2 W(kr) T(kr)\, \zeta_{\rm g}(\boldsymbol{k}), \tag{1}$$

$$\zeta_r(\boldsymbol{x}; r) = \int \frac{\mathrm{d}^3 k}{(2\pi)^3} e^{i\boldsymbol{k}\cdot\boldsymbol{x}} W_s(kr) T(kr)\, \zeta_{\rm g}(\boldsymbol{k}), \tag{2}$$

with the top-hat $W(y) = 3(\sin y - y\cos y)/y^3$, the shell $W_s(y) = \sin y/y$, the radiation transfer function $T(y) = W(y/\sqrt{3})$ and $\Phi = 2/3$. All moments use a hard cut $kr < 4.49$, which the peak moments need for broad spectra; it changes threshold-statistics amplitudes by $\leq 2\%$. For a spherical configuration $C_{\rm g} = -2\Phi r \zeta_{\rm g}'$, $\zeta_r = \zeta_{\rm g}(r)$, and the linear and full compaction of $\zeta = F(\zeta_{\rm g})$ are [12, 18]

$$C_\ell = F'(\zeta_r)\, C_{\rm g}, \qquad C = C_\ell - \frac{C_\ell^2}{4\Phi}. \tag{3}$$

A PBH of mass $M = \mathcal{K}(C - C_{\rm th})^\gamma M_H$ forms where $C > C_{\rm th}$ and $C_\ell < 2\Phi$ (type I). We use $\mathcal{K} = 3.3$ and $\gamma = 0.36$. The threshold $C_{\rm th} = \delta_c(q)$ follows the shape-dependent prescription of Musco et al. [24, 25], with the transfer function in the moments.

2.2 Threshold statistics

With $P_{\rm G}(c,z)$ the bivariate Gaussian of $(C_{\rm g},\zeta_r)$, the mass fraction at one scale is [12]

$$\beta_{\rm th} = \int {\rm d}z \int_{C_{\ell,\rm th}}^{2\Phi} {\rm d}C_\ell \, \frac{P_{\rm G}(C_\ell/F'(z),z)}{|F'(z)|} \, \mathcal{K}(C-C_{\rm th})^\gamma, \tag{4}$$

where $C_{\ell,\rm th} = 2\Phi[1-(1-C_{\rm th}/\Phi)^{1/2}]$. Equation (4) includes the non-linear $C(C_\ell)$ and the full F.

2.3 Peaks of the Gaussian component (Gaussian-peak approximation)

The natural generalisation of earlier work is to count BBKS maxima of the Gaussian field $C_{\rm g}$, with ζ_r as a correlated scalar, and map them through F'. Where $F'<0$, maxima of C_ℓ are minima of $C_{\rm g}$. This replaces the volume weight in (4) by

$$w = \frac{V\,G(m,s)}{(2\pi)^{3/2}R_*^3}, \qquad G(m,s) = \int_0^\infty {\rm d}x\, f(x)\,\mathcal{N}(x;m,s^2), \qquad V = \tfrac{4}{3}\pi r^3, \tag{5}$$

where:

- f is the BBKS function [14];
- m and s^2 are the conditional mean and variance of $x = -\nabla^2 C_{\rm g}/\sigma_2$ given $(C_{\rm g},\zeta_r)$;
- $R_* = \sqrt{3}\sigma_1/\sigma_2$.

This is exact for $F'=$ const, but not in general: $\nabla C_\ell = F'\nabla C_{\rm g} + F''C_{\rm g}\nabla\zeta_r$, so the maxima of C_ℓ are displaced from those of $C_{\rm g}$.

2.4 Peaks of the non-Gaussian compaction

The Kac–Rice density of maxima of C_ℓ at $(C_{\rm g},\zeta_r)=(c,z)$ follows from three steps:

1. The gradient factorises:

$$\nabla C_\ell = F'(z)\,\nabla Y_\lambda, \qquad Y_\lambda \equiv C_{\rm g} + \lambda\zeta_r, \qquad \lambda = \frac{cF''(z)}{F'(z)}, \tag{6}$$

   with λ frozen at the point, so that Y_λ is a Gaussian field.

2. At a stationary point, $\nabla\nabla C_\ell = F'[\nabla\nabla Y_\lambda + \kappa\nabla\zeta_r\nabla\zeta_r^T]$, with $\kappa = -cF'(1/F')''$. The rank-one term vanishes identically for the USR template, since $1/F'$ is linear there.

3. Conditional on $\nabla Y_\lambda = 0$, the gradient $\nabla\zeta_r$ is Gaussian with variance $v = (\sigma^2_{1,cc}\sigma^2_{1,rr} - \sigma^4_{1,cr})/(3\sigma^2_{1,\lambda})$. Here $\sigma^2_{n,AB} = \int {\rm d}\ln k\, k^{2n} K_A K_B \mathcal{P}_\zeta$ and $\sigma^2_{n,\lambda} = \sigma^2_{n,cc} + 2\lambda\sigma^2_{n,cr} + \lambda^2\sigma^2_{n,rr}$.

The determinant lemma then gives the peak density exactly to first order in $\tau = s\kappa v/\sigma_{2,\lambda}$ (Appendix A):

$$\boxed{n(c,z) = P_{\rm G}(c,z)\,\frac{G(s\,m_\lambda, s_\lambda)}{(2\pi)^{3/2}R_{*,\lambda}^3}\,\big[1-\tau R_2(s\,m_\lambda,s_\lambda)\big], \qquad R_{*,\lambda} = \frac{\sqrt{3}\,\sigma_{1,\lambda}}{\sigma_{2,\lambda}},} \tag{7}$$

with $s = \operatorname{sign} F'$, m_λ and s_λ the conditional moments of $-\nabla^2 Y_\lambda/\sigma_{2,\lambda}$, and $R_2 = 27\int f_2\mathcal{N}/\int f\mathcal{N} \to 9/m$. The peak mass fraction is (4) with the integrand multiplied by $Vn/P_{\rm G}$. Equation (7) has three limits:

- it reduces to (5) for $\lambda = 0$ (Gaussian ζ or $F''=0$);




Figure 1: Maxima of $C_\ell = F'(\zeta_r)C_g$ counted on 256^3 Gaussian random fields, divided by the prediction of eq. (7) (left) and of the Gaussian-peak approximation (5) (right), as a function of the threshold in units of σ_c. Gray: Gaussian ζ, which calibrates the grid.

- it reduces to (5) for a monochromatic spectrum, where $Y_\lambda \propto \zeta_r$;
- it is finite as $F' \to 0$.

The rank-one factor is formally one power of the r.m.s. amplitude smaller, but $\kappa = -2\lambda^2/c$ for the quadratic template. It changes the local density by up to 36% at nodes with $f_{\rm NL} = -2$, $\Delta = 1$.

Checks.

- An independent Monte Carlo of the 20 Gaussian variables with the exact Hessian reproduces (7) to 3×10^{-4} at 15 of 16 test nodes, and to 0.85% at the most extreme one.
- We counted local maxima of C_ℓ on 256^3 Gaussian random fields (lognormal $\Delta = 1$, 24 realisations), with predictions from the lattices' own mode sums (Fig. 1). Eq. (7) follows the counts to about 1% beyond the grid trend seen in the Gaussian case. The Gaussian-peak approximation (5) is off by up to 18% for $f_{\rm NL} = -2$.

3 Abundance and its conventions

The abundance integrates the mass fraction over the horizon mass M_H at which $k = 1/r$ re-enters:

$$f_{\rm PBH} = \int \mathrm{d}\ln M_H \; w(M_H)\,\beta(M_H). \tag{8}$$

Three normalisations of w are in use (Appendix B):

- the $M_{\rm eq}$ form $w = (M_{\rm eq}/M_H)^{1/2}/\Omega_{\rm DM}$ [12, 17];
- eq. (14) of 2306.17149, $w = (M_H/M_\odot)^{-1/2}(g_*/106.75)^{3/4}(g_{*s}/106.75)^{-1}/(7.9\times10^{-10}\,\Omega_{\rm DM})$;
- the first-principles result from entropy conservation, which is the same expression with 1.0×10^{-8} in place of 7.9×10^{-10}.

The first two overestimate $f_{\rm PBH}$ by factors of 3–5 and 12.7 respectively. Since β depends exponentially on $1/A$, this lowers the amplitude for $f_{\rm PBH} = 1$ by only $\ln(\text{factor})/|\ln\beta| \simeq 8\%$ and 12%. These factors cancel in R but not in the PTA tension.

The threshold prescription is a second, larger choice. The Musco et al. shape parameter q [24] depends on whether the transfer function enters its determination:

template	$\Delta = 0.1$	0.3	0.5	1	2
Gaussian	0.770	0.775	0.778	0.783	0.789
$f_{\rm NL} = -2$	0.738	0.744	0.769	0.794	0.810
$f_{\rm NL} = -1$	0.748	0.751	0.760	0.784	0.803
$f_{\rm NL} = +1$	0.788	0.792	0.796	0.803	0.811
$f_{\rm NL} = +2$	0.796	0.800	0.803	0.809	0.817
USR $\mu_* = 5/6$	0.808	0.816	0.825	0.851	0.914
USR $\mu_* = 5/2$	0.779	0.785	0.789	0.797	0.809

Table 1: $R = A_*^{\rm peaks}/A_*^{\rm threshold}$ with peaks of the non-Gaussian compaction (7), lognormal spectra of width Δ, $f_{\rm PBH} = 1$, $k_* = 10^7\,{\rm Mpc}^{-1}$.

- for a lognormal of width $\Delta = 1$, $\delta_c = 0.537$ with it and 0.458 without;
- for the broken power law of 2306.17149 $(\alpha, \beta, \gamma = 4, 3, 1)$, $\delta_c = 0.570$ and 0.544.

The thresholds in Fig. S3 of 2306.17149 are the values without the transfer function. The difference moves A_* by 18% (BPL) and 62% (lognormal, $\Delta = 1$), for every statistic, because the linear threshold $C_{\ell,\rm th}$ grows faster than $C_{\rm th}$.

Our implementation `ngpbh` was validated as follows:

- it reproduces the public code `pbhstat` [17] in its conventions to 0.017% in A_* for $\Delta = 0.1$–2, for both statistics;
- it reproduces the broad-spectrum benchmark of Ferrante et al. (amplitude 9.09×10^{-3} vs 9.2×10^{-3});
- it reproduces their curvaton amplitudes to 1% [12].

The integration uses Gauss–Legendre nodes in C_ℓ that absorb the $(C - C_{\rm th})^\gamma$ cusp and an adaptive window in ζ_r, all in $\log\beta$. It is stable to 10^{-5} when every grid is doubled.

4 The gap between the statistics, with NG

Table 1 and Fig. 2 give $R = A_*^{\rm peaks}/A_*^{\rm threshold}$ for $f_{\rm PBH} = 1$, with the same threshold, window, transfer function, cut, $\mathcal{K}$ and γ for both statistics.

- **Gaussian ζ:** $R = 0.770$–0.789 in the BBKS normalisation with $V = \frac{4}{3}\pi r^3$.
- **Conventions:** R spans 0.75–0.89 across the normalisation conventions in use (Press–Schechter factor, peak volume, Jacobian). It is about 0.6–0.67 with the high-peak formula of Ref. [9] as used in Ref. [11]. These conventions cancel to better than 1% in the double ratio $R/R_{\rm Gauss}$.
- **The double ratio:** it stays within 0.958–1.036 for $|f_{\rm NL}| \le 2$. The strongly non-perturbative USR tail $\mu_* = 5/6$ reaches 1.05–1.16, the largest value at $\Delta = 2$.
- **True NG peaks vs the Gaussian-peak approximation:** the amplitudes differ by $A_*^{\rm 3b}/A_*^{\rm 3a} = 0.983$–$1.000$. The 18% miscount of peaks moves A_* by at most 1.7%, because $\delta \ln A_* \simeq \delta \ln n/|\ln\beta|$.

For comparison, NG moves the threshold-statistics amplitude itself by factors of 0.2–2 over the same grid.




Figure 2: Left and middle: R for the quadratic and logarithmic templates; solid lines are peaks of C_ℓ (7), dashed lines are the Gaussian-peak approximation (5). Right: A_*^{3b}/A_*^{3a}.

5 Consequences for the PTA tension

5.1 Gravitational waves and the NANOGrav 15 likelihood

Induced GWs. For Gaussian ζ in radiation domination [5, 6, 8],

$$\Omega_{\rm GW,RD}(k) = \frac{1}{6}\int_1^\infty \mathrm{d}t \int_0^1 \mathrm{d}s \left[\frac{(t^2-1)(1-s^2)}{t^2-s^2}\right]^2 I_{t,s}^2\, \mathcal{P}_\zeta\Big(k\frac{t+s}{2}\Big)\mathcal{P}_\zeta\Big(k\frac{t-s}{2}\Big), \tag{9}$$

$$h^2\Omega_{\rm GW}(f) = h^2\Omega_r\,\frac{g_*}{g_*^0}\Big(\frac{g_{*s}}{g_{*s}^0}\Big)^{-4/3}\Omega_{\rm GW,RD}(k), \tag{10}$$

with $f = k/2\pi$, $h^2\Omega_r = 4.2\times10^{-5}$ and the Standard-Model $g_*(T)$ of [26]. The kernel is [5, 8]

$$I_{t,s}^2 = \frac{288(s^2+t^2-6)^2}{(t^2-s^2)^6}\left[\frac{\pi^2}{4}(s^2+t^2-6)^2\Theta(t-\sqrt{3}) + \Big(t^2-s^2-\tfrac{1}{2}(s^2+t^2-6)\ln\Big|\frac{t^2-3}{3-s^2}\Big|\Big)^2\right]. \tag{11}$$

We rewrite the logarithmic bracket as $(3-s^2)\,g(e)$ with $e=(t^2-s^2)/(3-s^2)$ and $g=-e^3/12-e^4/12-\ldots$, so that the cancellation at $t,s\to1$ is done analytically. We also split t at the resonance $t=\sqrt{3}$. Checks:

- a flat spectrum gives $0.8222436\,A^2$ [5];
- the Dirac closed form is recovered;
- direct two-dimensional integration agrees to 10^{-8}–10^{-12};
- the infrared k^3 tail matches its asymptotic form.

NG corrections to $\Omega_{\rm GW}$, of relative size $\sim A(3f_{\rm NL}/5)^2$, are neglected, as in [8]. They change the amplitude at the overproduction line by $\lesssim 3\%$.

Likelihood. We use the NANOGrav 15-year free spectrum with Hellings–Downs correlations (the lowest 14 bins), represented by kernel density estimates of the per-bin posteriors in $\log_{10}\rho$ [27, 28]:

$$\ln L = \sum_j \ln p_j(\log_{10}\rho_j), \qquad \rho_j^2 = \frac{H_{100}^2\,h^2\Omega_{\rm GW}(f_j)}{8\pi^4 f_j^5 T}, \qquad T = 16.03\,\mathrm{yr}. \tag{12}$$

The KDEs use a compact kernel with bandwidths down to 0.004 dex. Outside the sampled range their density is floored, and their sparse tails are combs of single-sample spikes. We therefore regularise them by taking the maximum of each density and its 0.05-dex Gaussian blur (Appendix C). This leaves the bulk unchanged and moves every significance below by $\leq 0.14\sigma$.

Validation of the likelihood:

- for $h_c \propto f^{-2/3}$ it gives $A = 2.52^{+0.59}_{-0.61} \times 10^{-15}$ (90%), against $2.4^{+0.7}_{-0.6} \times 10^{-15}$ from NANOGrav [1];
- with a free slope it gives $\gamma = 3.07^{+0.63}_{-0.56}$, against 3.2 ± 0.6;
- for the lognormal SIGW fit with Δ marginalised it gives $\log_{10} A = -0.57^{+0.54}_{-0.84}$, against $-0.56^{+0.53}_{-0.74}$ in 2306.17149;
- for the broken power law with the shape marginalised it gives $\log_{10} A = -0.84^{+0.79}_{-0.74}$, against $-0.86^{+0.79}_{-0.65}$, and $\log_{10} k_* = 7.95$ against 7.92.

Significance. The posterior is taken in $(\log_{10} k_*, \log_{10} A)$, uniform on $[5, 10] \times [-3, 0]$, for the same spectral shape as the PBH line. The "allowed region" is $A < A_*(k_*)$, where $f_{\rm PBH}(A_*) = 1$. We quote two measures:

- σ_P, the one-sided Gaussian equivalent of the posterior mass in the allowed region;
- $\sigma_{\rm HPD}$, the two-sided equivalent of the highest-posterior-density level at which the region is first reached. This is the definition of 2306.17149: their $1\sigma/2\sigma/3\sigma$ contours lie at our 69/94/99% levels.

$\sigma_{\rm HPD}$ saturates on local features of a lumpy posterior ridge, so we lead with σ_P. Values above 4σ lie where the KDEs carry no information and are quoted only as "> 4". The lines $A_*(k_*)$ for all k_* come from one set of β grids, because the moments depend on kr only.

5.2 Reproducing 2306.17149

Figure 3 compares our threshold-statistics bands with Fig. 3 of 2306.17149, read exactly from the vector paths of the arXiv source. Their setup reproduces their bands only with two of their unstated conventions: the threshold without the transfer function and their $f_{\rm PBH}$ normalisation. With both, the agreement for $k_* \geq 10^{7.5}\,{\rm Mpc}^{-1}$ is:

- Gaussian (non-linearities only): 1–6% in A;
- $f_{\rm NL} = -2$: within 6%;
- USR with $\beta = 3$: within 2.3%;
- curvaton $r_{\rm dec} = 0.9$: 8–11%.

Their bands dip by up to 0.05 dex near $k_* \simeq 10^{6.5}$, where they include the QCD softening of the formation parameters and we do not. Putting their own lines on our posterior changes the significance by $\leq 0.07\sigma$, so the QCD omission does not matter, as they state.

5.3 Exclusion per statistic and NG template

Table 2 gives σ_P for threshold statistics and for peaks of the non-Gaussian compaction, in the conventions of 2306.17149. Figure 5 shows all templates.

- **The statistics shift.** Over the 81 cases with $\sigma_P \leq 4$, peaks theory adds $\Delta\sigma_P = +0.24$ to $+0.78$ (median $+0.42$). The amplitude shift itself is nearly uniform: -0.07 to -0.11 dex, and -0.05 for the strongest USR tail at $\Delta = 2$. It translates into different $\Delta\sigma$ depending on where the line meets the posterior.
- **Their setup.** With the shape-marginalised posterior and the BPL(4,3,1) lines, the Gaussian case goes from 2.3 to $2.8\sigma_P$ (1.4 to $1.9\sigma_{\rm HPD}$). $f_{\rm NL} = -2$ goes from 1.6 to $1.9\sigma_P$, and the curvaton from 1.7 to $2.0\sigma_P$. The USR $\beta = 3$ case, which they place beyond 3σ, gives $3.9\sigma_P$ with threshold statistics and > 4 with peaks.




Figure 3: Left: threshold-statistics bands $f_{\rm PBH} = 1$ (upper) and 10^{-3} (lower) for BPL(4,3,1); wide lines are 2306.17149 Fig. 3 (with QCD), dashed lines are `ngpbh` (no QCD). Right: the NANOGrav 15 posterior with the BPL shape marginalised (fill: 68.3/95.4/99.7% HPD) and the contours of 2306.17149.

template	BPL(4,3,1)	LN $\Delta = 0.5$	LN $\Delta = 1$	LN $\Delta = 2$
Gaussian	$2.98 \to 3.62$	$3.02 \to 3.52$	$3.54 \to > 4$	$3.84 \to > 4$
$f_{\rm NL} = -2$	$1.87 \to 2.18$	$1.88 \to 2.31$	$3.43 \to 3.84$	> 4
$f_{\rm NL} = -1$	$1.98 \to 2.40$	$1.90 \to 2.36$	$3.05 \to 3.46$	$3.89 \to > 4$
$f_{\rm NL} = -0.5$	$2.25 \to 2.93$	$2.22 \to 2.89$	$3.09 \to 3.51$	$3.60 \to > 4$
$f_{\rm NL} = +0.5$	$3.67 \to > 4$	$3.60 \to > 4$	> 4	> 4
curvaton $r_{\rm dec} = 0.9$	$2.01 \to 2.44$	$2.00 \to 2.43$	$3.09 \to 3.49$	$3.91 \to > 4$
USR $\mu_* = 5/2$	$3.53 \to > 4$	$3.49 \to 3.91$	> 4	> 4
$f_{\rm NL} \geq +1$, USR $\mu_* \leq 5/6$	> 4	> 4	> 4	> 4

Table 2: Exclusion of the PBH-allowed region by NANOGrav 15, σ_P, threshold statistics $\to$ peaks of C_ℓ (7), $f_{\rm PBH} = 1$. Conventions of 2306.17149 (threshold without the transfer function, their $f_{\rm PBH}$ normalisation); fixed spectral shape in both the line and the posterior.

- **True NG peaks vs the Gaussian-peak approximation.** They agree to $|\Delta\sigma_P| \leq 0.04$.
- **The NG shift.** Relative to the Gaussian case it runs from -1.2σ (negative $f_{\rm NL}$ and the curvaton, for narrow spectra) to well beyond $+2\sigma$ (positive $f_{\rm NL}$ and USR tails). For $\Delta = 2$ the sign of the $f_{\rm NL} < 0$ shift flips, because large negative $\zeta_{\rm g}$ also produce large compaction.

5.4 Robustness

Table 3 varies one analysis choice at a time.

- **The statistics shift.** It survives every change: $+0.2$ to $+0.8\sigma$ across the peak-normalisation conventions in use, and $+0.8$ to $+1.3\sigma$ with the high-peak formula.
- **The absolute significance.** It depends mainly on the frequency range: 5, 14 or 30 bins move it by -1.5 to $+1\sigma$. The part of the posterior nearest the allowed region, $k_* \simeq 10^7\,\mathrm{Mpc}^{-1}$, places the GW peak at 10–20 nHz, and the bins above it bound it.
- **The other conventions:**
  - the transfer function in the threshold moves it by -0.2 to -1.0σ;




Figure 4: NANOGrav 15 posteriors (68.3/95.4/99.7% HPD) for BPL(4,3,1) and a lognormal of width 1, with the $f_{\rm PBH} = 1$ lines from threshold statistics (solid) and peaks of C_ℓ (dashed) for three NG templates. The legend gives σ_P.




Figure 5: σ_P per NG template for threshold statistics (circles), the Gaussian-peak approximation (open triangles, hidden behind the squares) and peaks of C_ℓ (squares). Shaded: beyond 4σ.

  - the entropy-conserving $f_{\rm PBH}$ normalisation by -0.15 to -0.4σ;
  - $f_{\rm PBH} = 10^{-3}$ instead of 1 by $+0.4$ to $+0.8\sigma$.

- **Fixed vs marginalised shape.** Fixing the shape raises the significance relative to marginalising over it (Gaussian BPL: 3.0 vs $2.3\sigma_P$). The fixed BPL(4,3,1) has a steep causality tail, which disfavours the low-k_* corner closest to the allowed region.

6 Discussion

What changes. The question we set out to answer is whether the gap between threshold statistics and peaks theory, once local NG is included for broad spectra, changes how strongly PBH overproduction disfavours a scalar-induced explanation of the PTA signal. It does, modestly and uniformly.

- Peaks theory lowers the overproduction line by about 0.1 dex for every NG template. This strengthens the exclusion by $\simeq 0.4\sigma$, or 0.2–0.8σ depending on the peak normalisation.

change from the reference	$\Delta\sigma_P$, threshold → peaks	max. change of σ_P
none (14 bins, KDE blur 0.05 dex)	+0.31 to +0.68	—
prior $\log_{10} k_* \in [6, 9]$, or $\log_{10} A \geq -4$	unchanged	0.00
posterior grid twice as fine	unchanged	0.00
KDE blur 0, 0.03 or 0.10 dex	+0.30 to +0.72	0.14
5 frequency bins (2–10 nHz)	+0.27 to +0.72	1.45
30 frequency bins (2–59 nHz)	+0.23 to +0.55	1.06
peak volume $(2\pi)^{3/2} r^3$	+0.40 to +0.81	0.14
Press–Schechter factor 2	+0.27 to +0.59	0.09
`pbhstat` v0.0.6 normalisation	+0.16 to +0.39	0.33
high-peak formula [9] (indicative)	+0.82 to +1.29	0.65

Table 3: Robustness of the PTA significances for BPL(4,3,1) and lognormals with $\Delta = 0.5$ and 1, over seven NG templates (cases with $\sigma_P \leq 4$). Reference prior: $\log_{10} k_* \in [5, 10]$, $\log_{10} A \in [-3, 0]$.

- The NG dependence of the gap cancels almost completely: the double ratio stays within ±4%. NG and the choice of statistics therefore add rather than interact.
- Negative $f_{\rm NL}$ and the curvaton remain the ways to relieve the tension, but peaks theory takes back 0.3–0.5σ of the relief.
- None of the conclusions of 2306.17149 changes qualitatively. Their footnote is correct and now quantified.

What dominates. For an absolute statement about the PTA tension, the larger uncertainties lie elsewhere. The frequency range of the fit, the treatment of the collapse threshold and the $f_{\rm PBH}$ normalisation each move the significance by about as much as the choice of statistics, or more. All three are rarely stated. We recommend quoting them with any PBH bound on the PTA signal, together with the peak normalisation when peaks theory is used. On the theory side, our result says that the Gaussian-peak approximation is adequate for the PTA question (to 0.04σ), although it miscounts peaks by up to 18%.

Limitations.

- As in the literature we compare with, we assume spherical symmetry for the compaction of a peak [18], use the mean profile (profile dispersion can matter for negative NG [22]), neglect QCD effects on the formation parameters and on the GW spectrum, use the free-spectrum refit rather than the full PTA likelihood, and ignore an astrophysical foreground. A supermassive-black-hole-binary component would relax every tension quoted here.
- The comparison between statistics is like-for-like, since all of these limitations affect both statistics in the same way.
- The absolute numbers carry the systematics of Table 3.

7 Conclusions

- **Construction.** We have constructed peaks theory with local NG for broad spectra on the compaction function, as the exact counterpart of threshold statistics. Maxima of the non-Gaussian compaction are maxima of a shifted Gaussian field, corrected by a rank-one term (eq. 7). Monte Carlo and lattice counts confirm this.
- **The gap.** Peaks require 18–26% smaller amplitudes than threshold statistics. This ratio is almost independent of NG for $|f_{\rm NL}| \leq 2$.

- **PTA consequence.** For NANOGrav 15 this strengthens the exclusion of a scalar-induced explanation by 0.2–0.8σ for every NG template. The effect is smaller than that of NG itself for $|f_{\rm NL}| \gtrsim 1$ and comparable to three analysis conventions, which we have quantified.
- **Code.** `ngpbh` implements every piece and reproduces published results: `pbhstat`, Ferrante et al., and 2306.17149.

A Peaks of the non-Gaussian compaction: the rank-one factor

At a stationary point of C_ℓ, write $H_\lambda = \nabla\nabla Y_\lambda$ and $\boldsymbol{g} = \nabla\zeta_r$, with $\boldsymbol{g} \sim \mathcal{N}(0, v\mathbb{I})$ independent of H_λ. The exact density is

$$n(c,z) = P_{\rm G}(c,z)\Big(\frac{3}{2\pi\sigma_{1,\lambda}^2}\Big)^{3/2}\mathbb{E}\big[|\det(H_\lambda + \kappa\boldsymbol{g}\boldsymbol{g}^T)|\,\Theta_s\big], \tag{13}$$

where Θ_s requires negative- (positive-) definiteness for $s = +1$ (-1). For $H \prec 0$ the matrix determinant lemma gives $\det(-H - \kappa\boldsymbol{g}\boldsymbol{g}^T) = \det(-H)\,[1 - \kappa\,\boldsymbol{g}^T(-H)^{-1}\boldsymbol{g}]$, which is linear in $\kappa|\boldsymbol{g}|^2$. Averaging over the direction and length of $\boldsymbol{g}$ gives $v\,\det(-H)\mathrm{tr}(-H)^{-1} = v\,e_2(-H)$. In the BBKS shape variables $e_2 = x^2/3 - (3y^2+z^2)/3$, so $f_2(x) = \mathbb{E}[e_2\Theta|x]$ follows from the same two-dimensional integral as f; that integral reproduces BBKS's f to 10^{-10}. The interior part of the first-order factor is then exactly $1 - \tau R_2$.

- For $\tau > 0$, maxima can only be lost, so $1 - \tau R_2$ is a lower bound.
- For $\tau < 0$, configurations that become maxima add a positive $\mathcal{O}(\tau^2)$ term.

The Monte Carlo bounds the remainder at 0.85% at the most extreme node tested, a shift of 0.02% in A_*.

B The PBH abundance from entropy conservation

After formation at temperature T, the PBH number per entropy is conserved, so $\rho_{\rm PBH,0} = \frac{3}{4}\beta\,(g_*/g_{*s})\,T\,s_0$. Here $s_0 = \frac{2\pi^2}{45}g_{*s}^0 T_0^3$ and β is the mass fraction at formation (including $\mathcal{K}(C - C_{\rm th})^\gamma$). With $M_H = 4\pi M_P^2/H$ and $H = \pi(g_*/90)^{1/2}T^2/M_P$ this gives

$$f_{\rm PBH} = \frac{1}{\Omega_{\rm DM}}\int \mathrm{d}\ln M_H \Big(\frac{M_H}{M_\odot}\Big)^{-1/2}\Big(\frac{g_*}{106.75}\Big)^{3/4}\Big(\frac{g_{*s}}{106.75}\Big)^{-1}\frac{\beta(M_H)}{1.0\times10^{-8}}, \tag{14}$$

in agreement with Carr et al. [29] to 2%. The constant 7.9×10^{-10} in eq. (14) of 2306.17149 makes $f_{\rm PBH}$ 12.7 times larger. The form $(M_{\rm eq}/M_H)^{1/2}\beta/\Omega_{\rm DM}$ drops the factor Ω_m and the heating of the radiation by g_{*s} changes after formation, and is 3–5 times larger. Both lower the amplitude for $f_{\rm PBH} = 1$ by $\ln(\text{factor})/|\ln\beta|$, i.e. 8–12% at $|\ln\beta| \simeq 21$.

C Free-spectrum density estimates

The per-bin densities distributed with PTArcade [28, 27] are Epanechnikov KDEs on a grid of 10^4 points in $\log_{10}\rho$. The kernel has compact support, so the density is exactly zero (stored as $\ln p = -36$) outside the range of the samples. Within the sparse tails it alternates between single-sample spikes and zero. The two most constraining bins (3.9 and 5.9 nHz) have almost no low-ρ tail. Tail probabilities beyond $\sim 2.5\sigma$ are therefore lumpy and cliff-like. We replace each density by the maximum of itself and its convolution with a 0.05-dex Gaussian: the bulk is unchanged and the holes are filled. Table 3 shows that the result changes by at most 0.14σ between no blur and 0.10 dex.

Code and data

`ngpbh` (MIT licence) is in the accompanying repository, with:

- the scripts that produce every table and figure here (`README.md`);
- 74 unit tests;
- the digitised Fig. 3 of 2306.17149.

The NANOGrav 15 free-spectrum densities are those of PTArcade (Zenodo record 10495907) and are downloaded with an MD5 check.

References

[1] G. Agazie *et al.* (NANOGrav), Astrophys. J. Lett. 951, L8 (2023), arXiv:2306.16213.

[2] A. Afzal *et al.* (NANOGrav), Astrophys. J. Lett. 951, L11 (2023), arXiv:2306.16219.

[3] K. N. Ananda, C. Clarkson, D. Wands, Phys. Rev. D 75, 123518 (2007), arXiv:gr-qc/0612013.

[4] D. Baumann, P. J. Steinhardt, K. Takahashi, K. Ichiki, Phys. Rev. D 76, 084019 (2007), arXiv:hep-th/0703290.

[5] K. Kohri, T. Terada, Phys. Rev. D 97, 123532 (2018), arXiv:1804.08577.

[6] J. R. Espinosa, D. Racco, A. Riotto, JCAP 09 (2018) 012, arXiv:1804.07732.

[7] G. Domènech, Universe 7, 398 (2021), arXiv:2109.01398.

[8] G. Franciolini, A. J. Iovino, V. Vaskonen, H. Veermäe, Phys. Rev. Lett. 131, 201401 (2023), arXiv:2306.17149.

[9] A. J. Iovino, G. Perna, A. Riotto, H. Veermäe, arXiv:2406.20089.

[10] A. J. Iovino, G. Perna, H. Veermäe, arXiv:2512.13648.

[11] D. Blas, J. W. Foster, Y. Gouttenoire, A. J. Iovino, arXiv:2602.12252.

[12] G. Ferrante, G. Franciolini, A. J. Iovino, A. Urbano, Phys. Rev. D 107, 043520 (2023), arXiv:2211.01728.

[13] A. Ianniccari, A. J. Iovino, A. Kehagias, D. Perrone, A. Riotto, Phys. Rev. D 109, 123549 (2024), arXiv:2402.11033.

[14] J. M. Bardeen, J. R. Bond, N. Kaiser, A. S. Szalay, Astrophys. J. 304, 15 (1986).

[15] S. Pi, M. Sasaki, V. Takhistov, J. Wang, JCAP 09 (2025) 045, arXiv:2501.00295.

[16] A. Kushwaha, T. Suyama, arXiv:2603.23025.

[17] P. S. Cole, J. Fumagalli, arXiv:2509.14326 (`pbhstat`).

[18] S. Young, arXiv:2201.13345.

[19] N. Kitajima, Y. Tada, S. Yokoyama, C.-M. Yoo, arXiv:2109.00791.

[20] R. Inui *et al.*, arXiv:2411.07647.

[21] F. Riccardi, M. Taoso, A. Urbano, arXiv:2102.04084.

[22] A. Escrivà, arXiv:2607.08738.

[23] C. Germani, M. A. Gorji, M. Uwabo-Niibo, M. Yamaguchi, arXiv:2503.05434.

[24] I. Musco, V. De Luca, G. Franciolini, A. Riotto, Phys. Rev. D 103, 063538 (2021), arXiv:2011.03014.

[25] A. Escrivà, C. Germani, R. K. Sheth, Phys. Rev. D 101, 044022 (2020), arXiv:1907.13311.

[26] K. Saikawa, S. Shirai, JCAP 08 (2020) 011, arXiv:2005.03544.

[27] W. G. Lamb, S. R. Taylor, R. van Haasteren, Phys. Rev. D 108, 103019 (2023), arXiv:2303.15442.

[28] A. Mitridate *et al.*, arXiv:2306.16377 (PTArcade).

[29] B. J. Carr, K. Kohri, Y. Sendouda, J. Yokoyama, Phys. Rev. D 81, 104019 (2010), arXiv:0912.5297.