EXP.NO:24

Pair with Difference-O(n^2)Time Complexity, O(1) Space Complexity

PROBLEM STATEMENT:

Given an array A of sorted integers and another non negative integer k, find if there exists 2 indices i and j such that A[j] - A[i] = k, i != j.

Input Format:

First Line n - Number of elements in an array

Next n Lines - N elements in the array

k - Non - Negative Integer

Output Format:

1 - If pair exists

0 - If no pair exists

Explanation for the given Sample Testcase:

YES as 5 - 1 = 4

So Return 1.

For example:

Input	Result
3 1 3 5 4	1

CODE

```c
#include <stdio.h>
int main(){
    int n;
    scanf("%d",&n);
    int a[n];
    for(int i=0;i<n;i++){
        scanf("%d",&a[i]);
    }
    int key;
    scanf("%d",&key);
    for(int i=0;i<n;i++){
        for(int j=0;j<n;j++){
            if(a[i]-a[j]==key && i!=j){
                printf("%d",1);
                return 0;
            }
        }
    }
    printf("%d",0);
}
```

EXP.NO:24

OUTPUT:



	Input	Expected	Got	
✔	3 1 3 5 4	1	1	✔
✔	10 1 4 6 8 12 14 15 20 21 25 1	1	1	✔
✔	10 1 2 3 5 11 14 16 24 28 29 0	0	0	✔
✔	10 0 2 3 7 13 14 15 20 24 25 10	1	1	✔

Passed all tests! ✔

Correct

Marks for this submission: 1.00/1.00.

RESULT:

The code is executed successfully.