STAT410: Intro to Probability Theory

Yuval Amit

Contents

1 Probability: The Mathematical Study of Uncertainty **2**

2 Continuity of Probability Measures **3**

3 Inclusion-Exclusion Principle **4**

4 Intro to Random Variables **5**

5 Conditional Probability **5**

6 Independence **7**

7 Random Variables **7**

8 Expected Values **10**

9 Special Distributions **12**

9.1 Bernoulli Distribution 12

9.2 Binomial Distribution 13

9.3 Hypergeometric Distributions 13

9.4 Poisson Distribution 14

1. Probability: The Mathematical Study of Uncertainty

Definition 1.1

A **probability space** $(\Omega, \mathcal{F}, P)$ is composed of

(i) The **sample space** Ω. It contains all possible outcomes of the "experiment".

(ii) The **event space** $\mathcal{F}$ consisting of **events**. An event is a collection of outcomes.

(iii) The **probability measure** $P : \mathcal{F} \to \mathbb{R}$.

Example 1.2. *Toss a fair coin. This has the sample space* $\Omega := \{H, T\}$*, the event space* $\mathcal{F} = \mathbb{P}(\Omega)$*, and the probability measure* $P : \mathcal{F} \to \mathbb{R}$ *given by*

$$P(\{H\}) = 1/2, \quad P(\{T\}) = 1/2, \quad P(\emptyset) = 0, \quad P(\{H, T\}) = 1$$

Example 1.3. *Roll a fair dice. This has the sample space* $\Omega = \{1, \ldots, 6\}$*, the event space* $\mathcal{F} = \mathbb{P}(\Omega)$*. Note that* $\mathcal{F}$ *is too big to succinctly write what the probability measure* P *does. More on this later.*

Example 1.4. *Pick a number uniformly at random from* $[0, 1]$*. Here,* $\Omega = [0, 1]$*. We pretend that* $\mathcal{F} = \mathbb{P}([0, 1])$*, but in reality* $\mathcal{F} \subsetneq \mathbb{P}(\Omega)$*. Again, we can't explicitly write what* P *does.*

Definition 1.5

We define the following axioms for the event space $\mathcal{F}$:

1. $\emptyset \in \mathcal{F}$ and $\Omega \in \mathcal{F}$.
2. If $A \in \mathcal{F}$, then $A^c \in \mathcal{F}$.
3. If $\{A_i\}$ is a sequence of events in $\mathcal{F}$, then $\bigcup A_i \in \mathcal{F}$.

This is the definition of a **σ-algebra**.

Remark 1.6. *If* $A_1, \ldots, A_n \in \mathcal{F}$*, then* $\bigcup_{i=1}^n A_i \in \mathcal{F}$.

Proof. Let $A_i = \emptyset$ for all $i > n$. Then, we know $\bigcup_{i=1}^n A_i = \bigcup_{i=1}^\infty A_i \in \mathcal{F}$ □

Proposition 1.7

If $\{A_i\}$ is a sequence of events in $\mathcal{F}$, then $\bigcap A_i \in \mathcal{F}$.

Proof.

$$\bigcap_{i=1}^\infty A_i = \left(\bigcup_{i=1}^\infty A_i^c \right)^c \in \mathcal{F}$$

by axiom 2, 3, and De Morgan's law. □

Definition 1.8

We define the following axioms for the probability measure P:

1. $P(\emptyset) = 0$ and $P(\Omega) = 1$.
2. For any $A \in \mathcal{F}$, $P(A) \geq 0$.

3. (Countable additivity) If $\{A_i\}$ is a sequence of pairwise-disjoint events, then
$$P\left(\bigcup_{i=1}^{\infty} A_i\right) = \sum_{i=1}^{\infty} P(A_i)$$

Example 1.9. *Let $\Omega = \{1, \dots, 6\}$ be the sample space for rolling a fair dice, and $\mathcal{F} = \mathbb{P}(\Omega)$. Recall that $P(\Omega) = 1$, and moreover*
$$\Omega = \{1\} \sqcup \{2\} \sqcup \cdots \sqcup \{6\}.$$
Thus,
$$1 = P(\Omega) = \sum_{i=1}^{6} P(\{i\}).$$
By the assumption that the dice is fair, we know $P(\{i\}) = P(\{j\})$, hence $P(\{i\}) = 1/6$ for all $i \in \{1, \dots, 6\}$.

Example 1.10. *Suppose $\Omega = [0, 1]$, and for any subinterval $I \subset [0, 1]$, $P(I) =$ length of I. What is the probability you see a rational number? An irrational number?*

Note that for any $x \in [0, 1]$, $P(\{x\}) = 0$ since $\{x\}$ can be considered a closed interval with length 0. Thus, since $\mathbb{Q}$ is countable, it follows that
$$P(\mathbb{Q} \cap [0, 1]) = 0, \quad P((\mathbb{R} \setminus \mathbb{Q}) \cap [0, 1]) = 1$$

Proposition 1.11

$$P(B^c) = 1 - P(B)$$

2. Continuity of Probability Measures

Let $B_1 \supset B_2 \supset \cdots$ be a decreasing sequence of sets. Then, we write
$$\lim_{n\to\infty} B_n = B = \bigcap_{i=1}^{\infty} B_i,$$
which is the greatest lower bound, or the infimum.

Theorem 2.1

The probability measure P commutes with limits. That is, given an ascending sequence $C_1 \subset C_2 \subset \cdots$,
$$P\left(\bigcup_{n=1}^{\infty} C_n\right) = P\left(\lim_{n\to\infty} C_n\right) = \lim_{n\to\infty} P(C_n)$$

Proof. Let $D_1 = C_1$, $D_2 = C_2 - C_1$, and generally $D_{i+1} := C_{i+1} - C_i$. Then, observe
$$\bigcup_{i=1}^{\infty} C_i = \bigcup_{i=1}^{\infty} D_i,$$
hence
$$P\left(\bigcup C_i\right) = P\left(\bigcup D_i\right).$$

Moreover, since the D_i's are pairwise disjoint, one shows that

$$P(D_{i+1}) = P(C_{i+1}) - P(C_i),$$

hence

$$\sum_{i=1}^{n} P(D_i) = P(C_n).$$

Thus,

$$\begin{aligned} P\left(\bigcup_{i=1}^{\infty} C_i\right) = P\left(\bigcup_{i=1}^{\infty} D_i\right) &= \sum_{i=1}^{\infty} P(D_i) \\ &= \lim_{n\to\infty} \sum_{i=1}^{n} P(D_i) = \lim_{n\to\infty} P(C_n) \end{aligned}$$

as desired. □

3. Inclusion-Exclusion Principle

Inclusion-Exclusion Principle **3.1**

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

Proof. Observe

$$P(A \cup B) = P(A \cup (B - A)) = P(A) + P(B - A).$$

Moreover,

$$P(B) = P(A \cap B) + P(B - A) \implies P(B - A) = P(B) - P(A \cap B),$$

thus

$$P(A \cup B) = P(A) + P(B) - P(A \cap B).$$

□

Corollary **3.2**

$$P(A \cup B \cup C) = P(A) + P(B) + P(C) - P(A \cap B) - P(B \cap C) - P(A \cap C) + P(A \cap B \cap C)$$

Proposition **3.3**

$$P\left(\bigcup_{i=1}^{n} A_i\right) \leq \sum_{i=1}^{n} P(A_i)$$

$$P\left(\bigcup_{i=1}^{\infty} A_i\right) \leq \sum_{i=1}^{\infty} P(A_i)$$

Proposition **3.4**

If $A \subset B$, then $P(A) \leq P(B)$.

4. Intro to Random Variables

Definition 4.1

A **random variable** is a (measurable) function $X : \Omega \to \mathbb{R}$.

Definition 4.2

The **cumulative distribution function**, abbreviated **cdf**, is

$$F_X(a) := P(X \leq a) = P(X \in [-\infty, a]).$$

Now, suppose X takes countably many values, i.e. $\{X(\omega) \mid \omega \in \Omega\}$ is countable. A **probability mass function**, abbreviated **pmt**, is a function p_X such that there's sequence of real numbers $\{x_n\}$ with $p_X(x_n) = P(X = x_n) > 0$ for all n, and $p_X(x) = P(X = x) = 0$ for all other x.

Example 4.3. *Any experiment with finitely many outcomes X is called a discrete random variable. Then, $P_X(x)$ is the probability mass function of X. The set where $P_X(x) > 0$ is called the **support**.*

For any set B, we have

$$P(x \in B) = \sum_{x \in \text{supp}(X) \cap B} p_X(x).$$

Definition 4.4

A random variable X is said to be **(absolutely) continuous** if there exists a function $f_X(x)$ such that $f_X(x) \geq 0$, and for any interval $I = [a, b]$,

$$P(x \in I) = \int_a^b f_X(x)dx$$

5. Conditional Probability

Question 5.1. *Does knowing A has taken place affect our chances of seeing B?*

Definition 5.2

Let A, B be events. Then, the probability of B given A is denoted $P(B|A)$.

Theorem 5.3

$$P(B|A) = \frac{P(A \cap B)}{P(A)}$$

Proof. We can give a frequentist proof with

$$P(B|A) = \lim_{n \to \infty} \frac{\frac{\#A,B \text{ both take place}}{n}}{\frac{\#A \text{ takes place}}{n}} = \frac{P(A \cap B)}{P(A)}.$$

□

Example 5.4. *An experiment consists of tossing a fair coin, followed by rolling a fair D6 if you see heads, and rolling a fair D4 if you see tails. What is the probability that you see heads and two?*

If $A =$ coin is heads and $B =$ dice is two, then our probability is

$$P(A \cap B) = P(A) \cdot P(B|A) = \frac{1}{2} \cdot \frac{1}{6} = \frac{1}{12}.$$

Theorem 5.5

$P'(X) := P(X|A)$ is a probability measure. However, $P''(X) := P(B|X)$ is not a probability measure.

***Law of Total Probability* 5.6**

Let $A_1, \ldots, A_n$ partition Ω. and $P(A_i) > 0$ for all i. Then,

$$P(B) = \sum_{i \in I} P(B|A_i)P(A_i)$$

This also holds for countable partitions.

***Bayes' Theorem* 5.7**

Suppose $A_1, \ldots, A_n$ partition Ω, and $P(A_i) > 0$. Suppose further that B is an event with $P(B) > 0$. Then, for any $j \in \{1, \ldots, n\}$,

$$P(A_j|B) = \frac{P(B|A_j)P(A_j)}{\sum_{i=1}^{n} P(B|A_i)P(A_i)},$$

as desired.

Proof. By Theorem 5.3,

$$P(A_j|B) = \frac{P(A_j \cap B)}{P(B)},$$

and by the law of total probability

$$\frac{P(A_j \cap B)}{P(B)} = \frac{P(A_j \cap B)}{\sum_{i=1}^{n} P(B|A_i)P(A_i)} = \frac{P(B|A_j) \cdot P(A_j)}{\sum_{i=1}^{n} P(B|A_i)P(A_i)}$$

□

Example 5.8. *The prevalence of a particular disease is 1%. A rapid test for this disease has a 2% false positive rate, and a 3% false negative rate. If the test comes up positive, how likely is it that you have the disease?*

Solution. Let D be the event that you have the disease, and T be the event that you tested positive. Then,

$$P(D) = 0.01,$$
$$P(T|D^c) = 0.02,$$
$$P(T^c|D) = 0.03.$$

By Bayes' Theorem,

$$P(D|T) = \frac{P(T|D)P(D)}{P(T|D)P(D) + P(T|D^c)P(D^c)} = \frac{(1-0.03)\cdot 0.01}{(1-0.03)\cdot 0.01 + 0.02\cdot(1-0.01)} = 0.3288$$

□

6. Independence

Definition 6.1

Events A and B are said to be **independent** of $P(A \cap B) = P(A)P(B)$.

Theorem 6.2

Let A, B be events with $P(A), P(B) > 0$. If they are independent, they cannot be disjoint.

Proof. $P(A \cap B) = P(A)P(B) > 0$. □

Theorem 6.3

For any events $A_1, \ldots, A_n$ with $P(A_i) > 0$,

$$P\left(\bigcap_{i=1}^{n} A_i\right) = P\left(A_n | \bigcap_{i=1}^{n} A_i\right) P\left(A_{n-1} | \bigcap_{i=1}^{n-2} A_i\right) \cdots P(A_1).$$

Example 6.4. *An experiment consists of tossing a fair coin until you see heads for the first time, at which point the experiment terminates. Let A_i be the event you see tails on the i'th toss. Are the A_i's independent? What is the probability we see heads for the first time on the n'th toss?*

No, the A_i's are not independent. Next, let B_n be the event that heads is seen on the n'th toss. Then,

$$B_n = A_1 \cap A_2 \cap \cdots \cap A_{n-1} \cap A_n^c.$$

Therefore,

$$P(B_n) = P\left(A_n^c | \bigcap_{i=1}^{n-1} A_i\right) \cdot \prod_{k=1}^{n-1} P\left(A_k | \bigcap_{i=1}^{k-1} A_i\right) = (1/2)^n.$$

Exercise 1. *If A, B are independent, show that A, B^c are independent.*

7. Random Variables

Definition 7.1

For a random variable X, and a (measurable) $A \subset \mathbb{R}$, we define the distribution D_X to be

$$D_X(A) = P(X \in A)$$

Now, given a random variable $X : \Omega \to \mathbb{R}$, we have

$$D_X(A) = P(X \in A) = P\left(\{\omega \mid X(\omega) \in \mathbb{R}\}\right),$$

and

$$F_X(a) = P(X \leq a) = P(X \in (-\infty, a]).$$

Clearly, D_X determins F_X, but in fact we'll see that F_X determines D_X.

Example 7.2. *Let X be the outcome of rolling a fair die. Then,*

$$F_X(a) = \begin{cases} 0 & a < 1 \\ 1/6 & 1 \le a < 2 \\ 2/6 & 2 \le a < 3 \\ \vdots & \\ 5/6 & 5 \le a < 6 \\ 1 & 6 \le a. \end{cases}$$

In other words,

$$F_X(a) = \begin{cases} 0 & a < 1 \\ 1 & a \ge 6 \\ \frac{\lfloor a \rfloor}{6} & 1 \le a < 6. \end{cases}$$

Proposition 7.3

The function $F_X : \mathbb{R} \to \mathbb{R}$ satisfies the following properties:

1. $\lim_{X \to \infty} F_X(a) = 1$ and $\lim_{X \to -\infty} F_X(a) = 0$
2. F_X is monotonically increasing
3. F_X is right continuous, i.e.

$$\lim_{x \to a^+} F_X(x) = F_X(a)$$

Definition (*Discrete Random Variables*) 7.4

X is called **discrete** if there is a countable $A \subset \mathbb{R}$ such that $P(X \in A) = 1$.

Then, the function $a \mapsto P(X = a)$ is called the **probability mass function** (pmf), and the probabilites of interest are computed by taking a suitable sum of the pmf.

Definition (*Continuous Random Variables*) 7.5

X is called **(absolutely) continuous** if there is some function $f_X(x)$ such that $f_X(x) \ge 0$, and for any interval I with endpoints $-\infty \le a \le b \le \infty$,

$$P(X \in I) = \int_a^b f_X(x)dx$$

Note that we then have

$$F_X(a) = \int_{-\infty}^a f_X(x)dx$$

Definition (*Identical Distributions*) 7.6

Random variables X, Y are said to be **identically distributed** if $D_X = D_Y$, which holds if and only if $F_X = F_Y$.

A good way to think about identical distributions is with an example: how do you generate a random integer in $\{0,1\}$?

One way to do it is flipping a coin, taking 0 if the coin is tails, and 1 if the coin is heads. Another way to do it is generating a random real number in $[0,1]$, taking 0 if it's $< 1/2$, and 1 if it's $\geq 1/2$.

Let X be the coin method and Y be the real number method. Note that X and Y have completely different sample spaces:

$$X : \{H,T\} \to \mathbb{R} \quad X(\{H\}) = 1, X(\{T\}) = 0$$
$$Y : [0,1] \to \mathbb{R} \qquad Y(\omega) = \begin{cases} 0 & \omega < 0.5 \\ 1 & \omega \geq 0.5. \end{cases}$$

However, observe

$$F_X(a) = F_Y(a) = \begin{cases} 0 & a < 0 \\ 1/2 & 0 \leq a < 1 \\ 1 & a \geq 1. \end{cases}$$

Thus, X and Y are identically distributed, even though they're completely different (they have different sample spaces!).

Moreover, let $W = 1 - X$, i.e. $W = 1$ on tails and $W = 0$ on heads. Then, X and W are still identically distributed!

Remark 7.7. 1. *If F_X is piecewise constant, then X is a discrete random variable. In this case, $P(X = a)$ can be computed by $F(a) - \lim_{X \to a^-} F_X(a)$*

2. *If X is (absolutely) continuous, F_X is continuous*

3. *If F_X is differentiable with only finitely many exceptions, then $F_X'(x) = f_X(x)$ for all points where F_X is differentiable. At other points, we may define f_X arbitrarily.*

Example 7.8. *Suppose X is a continuous random variable with pdf $f_X(x) = 1/2$ for $x \in [-1,1]$.*

(i) Compute $P(X > 0)$

We know

$$P(X > 0) = \int_0^\infty f_X(x)dx = \int_0^1 \frac{1}{2}dx = 1/2$$

(ii) Compute $P(X > 1/2 | X > 0)$.

Using conditional probability, we see that

$$P(X > 1/2 | X > 0) = \frac{P(X > 1/2 \cap X > 0)}{P(X > 0)} = \frac{1/4}{1/2} = \frac{1}{2}.$$

(iii) Compute the pdf of X^2.

Observe that

$$P(X^2 \leq a) = P(-\sqrt{a} \leq X \leq \sqrt{a}) = \int_{-\sqrt{a}}^{\sqrt{a}} f_X(x)dx.$$

When $0 \leq a < 1$, this evaluates to $\sqrt{a}$. Thus, we see

$$F_{X^2}(a) = \begin{cases} 0 & a < 0 \\ \sqrt{a} & 0 \leq a < 1 \\ 1 & a \geq 1. \end{cases}$$

Note that this is the cdf of X^2. Differentiating, we get

$$f_{X^2}(a) = \begin{cases} 0 & a < 0 \\ \frac{1}{2\sqrt{a}} & 0 \leq a < 1 \\ 0 & a \geq 1. \end{cases}$$

Definition **7.9**

We say that a random variable X has a **uniform distribution** on $[a, b]$, denoted $X \sim u[a, b]$, if

$$f_X(x) = \frac{1}{b-a}$$

for $x \in a, b$.

Example 7.10. *Suppose $X \sim u[0, 1]$. Find the pdf of $Y := e^X$.*

Solution. If $a < 1$, then e^X doesn't attain a on $[0, 1]$, so $F_Y(a) = 0$ for $a < 1$. Similarly, when $a > e$, we know with certainty that $e^X < a$, so $F_Y(a) = 1$. Finally, if $a \in [1, e]$, then

$$F_Y(a) = P(Y \leq a) = P(e^X \leq a) = P(X \leq \log a) = P(0 \leq X \leq \log a) = \log a.$$

Thus,

$$F_Y(a) = \begin{cases} 0 & a < 1 \\ \log(a) & 1 \leq a \leq e \\ 1 & a > e, \end{cases}$$

and differentiating yields

$$f_Y(a) = \begin{cases} 0 & a \leq 1 \\ \frac{1}{a} & 1 < a < e \\ 0 & a \geq e. \end{cases}$$

□

8. Expected Values

Definition (*Expected Value (Discrete)*) **8.1**

Let X be a discrete random variable with support A. Then, the **expected value** of X, denoted $E[X]$, is

$$E[X] = \sum_{x \in A} x p_X(x) = \sum_{x \in A} x P(X = x)$$

Definition (*Expected Value (Continuous)*) **8.2**

Let X be a continuous random variable. Then, the **expected value** of X, denoted $E[X]$, is

$$E[X] = \int_{-\infty}^{\infty} x f_X(x) dx$$

Definition **8.3**

We say that the expected value $E[X]$ **exists** if $E[|X|] < \infty$.

Example 8.4. *Suppose $X \sim u[0,1]$. Compute $E[X]$ and $E[e^X]$.*

Solution. By definition,

$$E[X] = \int_{-\infty}^{\infty} x f_X(x) dx = \int_0^1 x dx = \frac{1}{2}.$$

For $E[e^X]$, one method involves using the pdf we computed earlier, which gives

$$E[e^X] = \int_{-\infty}^{\infty} y f_Y(y) dy = \int_1^e \frac{y}{y} dy = e - 1.$$

However, we may use the following general result:

$$E[g(x)] = \int_{-\infty}^{\infty} g(x) f_X(x) dx,$$

which gives

$$E[e^X] = \int_0^1 e^x dx = e - 1.$$

□

LOTUS **8.5**

Let X be a random variable with density function f_X. Then, for $g : \mathbb{R} \to \mathbb{R}$,

$$E[g(X)] = \int_{-\infty}^{\infty} g(x) f_X(x) dx$$

Linearity of Expectation **8.6**

If X, Y are random variables, and a is a real number, then $E[aX + Y] = aE[X] + E[Y]$.

Markov's Inequality **8.7**

Suppose $X \geq 0$ and $c \in \mathbb{R}^+$. Then

$$P(X \geq c) \leq E[X]/c,$$

Chebyshev's Inequality **8.8**

Let X be a random variable with finite mean μ and finite variance σ^2. For any $c \in \mathbb{R}^+$,

$$P\left(|X - \mu| \geq c\right) \leq \frac{\sigma^2}{c^2}$$

Proposition 8.9

If $E[X^k]$ is finite, then $E[X^n]$ is finite for all $n \in \{1, \dots, k\}$. In particular, for Chebyshev's inequality, it's sufficient to know that σ^2 is finite.

Proof. Observe

$$E\left[|X^m|\right] = \int_{-\infty}^{\infty} |x^m| f_X(x) dx = \int_{|x^m|>1} |x^m| f_X(x) dx + \int_{|x^m|\leq 1} |x^m| f_X(x) dx$$

by LOTUS (Theorem 8.5), for any m. Let A be the first term above when $m = k$, and B the second term when $m = k$. When $x^n > 1$, then $|x^n| f_X(x) \leq |x^k| f_X(x)$. When $|x^m| \leq 1$, then $|x^m| f_X(x) \leq f(x)$. Thus,

$$E[|X^n|] \leq \int_{|x^n|>1} |x^k| f_X(x) dx + \int_{-\infty}^{\infty} f_X(x) \leq A + 1 < \infty,$$

so $E[X^n]$ is finite. □

9. Special Distributions

In this section, we study some important distributions to know.

Discrete Distributions

9.1. Bernoulli Distribution

Definition 9.1

A **Bernoulli trial** is a random variable which has two outcomes: success (1) and failure (0).

Proposition 9.2

A bernoulli random variable X has a probability mass function

$$p_X(x) = \begin{cases} 1-p & x = 0 \\ p & x = 1 \end{cases}$$

where p is the probability of success. Moreover,

$$E[X] = p$$
$$V[X] = p(1-p)$$

Proof. We compute

$$E[X] = \sum x p_X(x) = 0 \cdot (1-p) + 1 \cdot p = p$$
$$V[X] = E[X^2] - E[X]^2 = p - p^2 = p(1-p)$$

□

Example 9.3. *Repeat a Bernoulli trial in an independent manner. Let X be a random variable that counts*

the number of trials until the first success. Then, we know X has the probability mass function

$$p_X(x) = p(1-p)^{x-1} \qquad x = 1, 2, \dots$$

Moreover,

$$E[X] = \sum_{n=1}^{\infty} n \cdot p(1-p)^{n-1} = \frac{1}{p}$$

To compute this sum, use the derivative trick. Remember the sums and derivatives commute when we're within the radius of convergence.

Question 9.4. *The probability of an independent event happening is p. What's the expected number of attempts does it take for the event to happen?*

Solution. Let X be the number of attempts to make the event happen, and we saw in the previous example that $X \sim \text{Geom}(1/10^7)$. Then, $E[X] = 10^7$. □

9.2. Binomial Distribution

We repeat a Bernoulli trial, with success chance p, n times, where n is fixed. Let X be the random variable counting the number of successes.

Let's compute $P(X = k)$. This involves k successes, which has probability p^k, and $n - k$ failures, with probability $(1-p)^k$. Now, we must count the number of ways in which we can achieve k successes. This is $\binom{n}{k}$, and so

$$P(X = k) = \binom{n}{k} p^k (1-p)^{n-k}$$

Moreover,

$$E[X] = \sum_{k=0}^{n} k \binom{n}{k} p^k (1-p)^{n-k}.$$

This sum can be computed with some neat tricks. However, we will eventually see that X can be written $X = X_1 + \cdots + X_n$ as the sum of n Bernoulli random variables, which are independent and identically distributed (**IID**). Note that $E[X_i] = p$, so $E[X] = np$.

9.3. Hypergeometric Distributions

Say we want to sample a portion of a population for some property. Do we sample with or without replacement? If we sample with replacement, we get a binomial distribution. However, it can be better to sample with replacement, in which case we get a hypergeometric distribution.

Definition 9.5

Let N, N_A, n be integers with $0 \le n \le N_A \le N$. A random variable X is said to follow a **hypergeometric distribution** with respect to the parameters (N, N_A, n) if X takes values in $\{1, \dots, n\}$, and

$$P(X = k) = \frac{\binom{N_A}{k}\binom{N-N_A}{n-k}}{\binom{N}{n}}.$$

This is interpreted as follows: there are $\binom{N}{n}$ ways to select n people out of your population of N. Then, if N_A of those people have the condition, there are $\binom{N_A}{k}$ ways to find k of them, and $\binom{N-N_A}{n-k}$ for the rest of them to not have it.

Example 9.6. *Suppose that a collection of 20 rare books contains 5 first editions, each costing* \$5000*, and 15 second editions, each costing* \$1000*. Four books are picked at random to be sold. What is the expected income?*

Solution. Let X be the number of first editions, so $4 - X$ is the number of second editions. Then, the revenue is

$$r(X) := 5000X + 1000(4 - X) = 4000X + 4000.$$

By linearity of expectation,

$$E[r(X)] = E[4000X + 4000] = 4000E[X] + 4000.$$

We're now left to compute $E[X]$. Note that X is sampling randomly from a population without replacement, so it's a hypergeometric distribution. The population size is $N = 20$, the number of first editions is $N_A = 5$, and the sample size is $n = 4$. Computing $E[X]$ is left as an exercise. □

9.4. Poisson Distribution

Definition 9.7

Fix $\lambda > 0$. A random variable X is said to have a **Poisson distribution** with parameter λ if $X \geq 0$ is integer valued, and has a probability mass function

$$p_X(k) = e^{-\lambda}\frac{\lambda^k}{k!}$$

for $k \in \{0, 1, 2, \dots\}$.

Remark 9.8. *If* $X \sim \text{Poisson}(\lambda)$*, then* $E[X] = V[X] = \lambda$.

Let $X \sim \text{Bin}(n.p)$ and $Y \sim \text{Poisson}(np)$. Then, for any $A \subset \{0, 1, 2, \dots\}$, we have

$$|P(X \in A) - P(Y \in A)| \leq np^2$$

Example 9.9. *Assume that a random variable X counts the occurrence of rare events that are not "strongly dependent". Then, the distribution of X can be approximated with a* $\text{Poisson}(E[X])$ *distribution.*

Example 9.10. *Suppose that a large factory experiences, on average, three accidents each month. What is the probability that on a particular month there are two accidents?*

Solution. Since the factory is large, we can assume that three accidents per month is very small, so we may assume that the number of accidents is represented by $X \sim \text{Poisson}(3)$. Then,

$$P(X = 2) = e^{-3}\frac{3^2}{2!} \approx 0.224$$

□

Continuous Distributions

There are three types of continuous distributions: uniform, exponential, and normal.

Uniform distributions try to compute the idea that the regions with equal "length" have the same probability of having something "drawn" from them.

Definition 9.11

A random variable X is said to have a **uniform distribution**, denoted $X \sim \text{Unif}[a, b]$ for $a < b$ if

$$f_X(x) = \frac{1}{b-a}, \quad x \in [a, b]$$