

Assignment 5

BY YUVAL BERNARD

Date:

Page 168

Question 23

$$t^2 y'' - 4t y' + 6y = 0, \quad t > 0, \quad y_1 = t^2$$

The general method is as follows: for a second order equation of the form:

$$y_2'' + p(x) y_2' + q(x) y_2 = 0 \quad (1)$$

where $y_1(t)$ is a solution, write a second solution as $y_2(t) = y_1(t) v(t)$.

$$\begin{aligned} y_2' &= y_1' v + y_1 v' \\ y_2'' &= y_1'' v + 2y_1' v' + y_1 v'' \end{aligned}$$

Get

$$(y_1'' v + 2y_1' v' + y_1 v'') + p(x) (y_1' v + y_1 v') + q(x) (y_1 v) = 0$$

Rewrite as

$$(y_1'' + p(x) y_1' + q(x) y_1) v + (2y_1' + p(x) y_1) v' + y_1 v'' = 0$$

The red term is equal to zero because y_1 is a solution. Let's look at what's left.

$$(2y_1' + p y_1) v' + y_1 v'' = 0 \quad (2)$$

This is a 2nd order ODE in v , or a 1st order linear ODE in v' . Therefore we can find y_2 by “reduction of order”.

Rewrite the equation to match the form of eq. (1). Divide by $t^2 \neq 0$:

$$y'' - \frac{4}{t} y' + \frac{6}{t^2} y = 0, \quad t > 0$$

Write a second solution $y_2 = y_1 v = t^2 v$. According to eq. (1), $p(t) = -\frac{4}{t}$, $q(t) = \frac{6}{t^2}$.

Get the reduced ODE according to eq. (2):

$$\underbrace{\left(2 \cdot 2t - \frac{4}{t} \cdot t^2\right)}_{=0} v' + t^2 v'' = 0$$

We are left with the equation

$$t^2 v'' = 0$$

Therefore, $v'' \equiv 0 \forall t$.

$$v' = \int v'' dt = c_1$$

$$v = \int v' dt = c_1 t + c_2$$

In conclusion,

$$y_2 = y_1 v = t^2 (c_1 t + c_2) = c_1 t^3 + c_2 t^2, \quad t > 0$$

For a specific y_2 , choose $c_1 = 1, c_2 = 0$. $y_2 = t^3, (t > 0)$.

Question 24

$$t^2 y'' + 2t y' - 2y = 0, \quad t > 0, \quad y_1 = t$$

Rewrite the equation to match the form of eq. (1). Divide by $t^2 \neq 0$:

$$y'' + \frac{2}{t} y' - \frac{2}{t^2} y = 0, \quad t > 0$$

Here $p = \frac{2}{t}, q = -\frac{2}{t^2}$. Select a solution $y_2 = y_1 v$. According to eq. (2),

$$\left(2 \cdot 1 + \frac{2}{t} t\right) v' + t v'' = 0$$

$$4v' + t v'' = 0, \quad t > 0$$

Divide by $t \neq 0$:

$$v'' + \frac{4}{t} v' = 0, \quad t > 0$$

This is a second order linear ODE in v . Substitute $f = v'$ to reduce equation order. The equation becomes

$$f' + \frac{4}{t} f = 0, \quad t > 0$$

Solve using integration factor method.

$$f(t) = e^{-\int \frac{4}{t} dt} \left[\int 0 \cdot e^{\int \frac{4}{t} dt} dt + c \right]$$

$$f(t) = \frac{c_1}{t^4}$$

Integrate to find v .

$$v(t) = \int f(t) dt = -\frac{c_1}{3} t^{-3} + c_2$$

Substitute in expression for y_2 :

$$y_2 = y_1 v = -\frac{c_1}{3} t^{-2} + c_2 t, \quad t > 0$$

For a specific solution, choose $c_1 = -3, c_2 = 0$. $y_2 = t^{-2}, \quad t > 0$.

Question 27

$$x y'' - y' + 4x^3 y = 0, \quad x > 0, \quad y_1(x) = \sin x^2$$

Rewrite the equation to match the form of eq. (1). Divide by $x \neq 0$:

$$y'' - \frac{1}{x} y' + 4x^2 y = 0, \quad x > 0$$

Here $p = -\frac{1}{x}, q = 4x^2$. Select a solution of the form $y_2 = y_1 v = \sin x^2 v$. According to eq. (2),

$$\left(2 \cdot \cos x^2 \cdot 2x - \frac{1}{x} \cdot \sin x^2 \right) v' + \sin x^2 v'' = 0$$

$$\left(4x \cos x^2 - \frac{1}{x} \sin x^2 \right) v' + \sin x^2 v'' = 0$$

Divide by $\sin x^2 \neq 0$: ($x \neq \pm \sqrt{\pi k}, k = 1, 2, 3, \dots$)

$$v'' + \left(\frac{4x}{\tan x^2} - \frac{1}{x} \right) v' = 0$$

Substitute $f = v'$,

$$f' + \left(\frac{4x}{\tan x^2} - \frac{1}{x} \right) f = 0$$

This is a linear first order ODE in f . Solve using integration factor method.

$$f(x) = e^{-\int \left(\frac{4x}{\tan x^2} - \frac{1}{x}\right) dx} \left[\int 0 \cdot e^{\int \left(\frac{4x}{\tan x^2} - \frac{1}{x}\right) dx} + c \right]$$

$$f = e^{\int \left(\frac{1}{x} - \frac{4x}{\tan x^2}\right) dx} \cdot c_1$$

$$\int \left(\frac{1}{x} - \frac{4x}{\tan x^2}\right) dx = \ln |x| - 2 \int \frac{2x}{\tan x^2} dx$$

$$\int \frac{2x}{\tan x^2} dx = [u = x^2, du = 2x dx] \int \frac{1}{\tan u} du = \int \frac{\cos u}{\sin u} du$$

Substitute $\ell = \sin u$. $d\ell = \cos u du$.

$$\int \frac{2x}{\tan x^2} dx = \dots = \int \frac{1}{\ell} d\ell = \ln |\ell| = \ln |\sin u| = \ln |\sin x^2|$$

In summary,

$$f(x) = c_1 e^{(\ln |x| - 2 \ln |\sin x^2|)} = c_1 \frac{x}{\sin^2 x^2}, \quad \sin x^2 \neq 0$$

Integrate to get v :

$$v = \int f dx = c_1 \int \frac{x}{\sin^2 x^2} dx = [u = x^2] = \frac{c_1}{2} \int \frac{1}{\sin^2 u} du$$

$$\int \frac{1}{\sin^2 u} du = \int \frac{\frac{1}{\cos^2 u}}{\frac{\sin^2 u}{\cos^2 u}} du = \int \frac{1}{\tan^2 u} du$$

Substitute $\ell = \tan u$, $d\ell = \frac{1}{\cos^2 u} du$.

$$\int \frac{1}{\sin^2 u} du = \int \frac{1}{\ell^2} d\ell = -\frac{1}{\ell} = -\frac{1}{\tan u} = -\frac{\cos x^2}{\sin x^2}$$

Therefore,

$$v = -\frac{c_1 \cos x^2}{2 \sin x^2} + c_2$$

$$y_2 = y_1 v = -\frac{c_1}{2} \cos x^2 + c_2 \sin x^2$$

For a specific solution, choose $c_1 = -2$, $c_2 = 0$. $y_2 = \cos x^2$, $x > 0$.

Page 220

Question 11

$$y''' - y'' - y' + y = 0$$

As we've seen in class, given: $a_n y^{(n)} + a_{n-1} y^{(n-1)} + \dots + a_0 y = 0$, $a_n \neq 0$, we look for solutions of the form $e^{\lambda x}$ and get:

$$y^{(k)}(x) = \lambda^k e^{\lambda x} \quad \forall x$$

And when we substitute in the ODE:

$$e^{\lambda x} (a_n \lambda^n + a_{n-1} \lambda^{n-1} + \dots + a_1 \lambda + a_0) = 0$$

Then $e^{\lambda x}$ is a solution if and only if $a_n \lambda^n + a_{n-1} \lambda^{n-1} + \dots + a_1 \lambda + a_0 = 0$. (This equation is also called the *characteristic polynomial of the ODE*.)

If the polynomial has n distinct solutions $\lambda_1, \dots, \lambda_n$ then $e^{\lambda_1 x}, \dots, e^{\lambda_n x}$ are n linearly independent solutions and form a basis for space of solutions.

Additionally, if λ_0 is a root of the characteristic polynomial of multiplicity r , then $e^{\lambda_0 x}, x e^{\lambda_0 x}, x^2 e^{\lambda_0 x}, \dots, x^{r-1} e^{\lambda_0 x}$ will all solve the ODE and are linearly independent, and will be linearly independent of solutions we obtain from other roots of the characteristic polynomial.

The equation for the characteristic polynomial of the equation is:

$$\lambda^3 - \lambda^2 - \lambda + 1 = 0$$

Note that $\lambda = -1$ solves the equation. The equation may be simplified (via long polynomial division):

$$(\lambda + 1)(\lambda^2 - 2\lambda + 1) = (\lambda + 1)(\lambda - 1)^2 \Rightarrow \lambda_1 = -1, \lambda_2 = 1$$

- From $\lambda = -1$ we get the solution: e^{-x} .
- From $\lambda = 1$ we get the solutions: $e^x, x e^x$.

The general solution is a linear combination of all solutions we've found:

$$y = c_1 e^{-x} + c_2 e^x + c_3 x e^x$$

Question 20

$$y^{(4)} - 8y' = 0$$

The characteristic polynomial equation is

$$\lambda^4 - 8\lambda = 0$$

$$\lambda(\lambda^3 - 8) = 0$$

$$\lambda(\lambda - 2)(\lambda^2 + 2\lambda + 4) = 0$$

$$\lambda^2 + 2\lambda + 4 = 0 \Rightarrow \lambda = \frac{-2 \pm \sqrt{2^2 - 4 \cdot 4}}{2} = -1 \pm \sqrt{3}i$$

- For $\lambda = 0$ the solution is $e^0 = 1$
- For $\lambda = 2$ the solution is e^{2x}
- For $\lambda = -1 \pm \sqrt{3}i$ the solutions are $e^{-x} \cos(\sqrt{3}x)$ and $e^{-x} \sin(\sqrt{3}x)$

The general solution is

$$c_1 + c_2 e^{2x} + c_3 e^{-x} \cos(\sqrt{3}x) + c_4 e^{-x} \sin(\sqrt{3}x)$$

Question 29

$$y''' + y' = 0, \quad \begin{cases} y(0) = 0 \\ y'(0) = 1 \\ y''(0) = 2 \end{cases}$$

Find roots of characteristic polynomial:

$$\lambda^3 + \lambda = 0$$

$$\lambda(\lambda^2 + 1) = 0$$

$$\lambda(\lambda - i)(\lambda + i)$$

$$y = c_1 + c_2 \cos t + c_3 \sin t$$

$$y' = -c_2 \sin t + c_3 \cos t$$

$$y'' = -c_2 \cos t - c_3 \sin t$$

Inputting $t = 0$ gives:

$$y(0) = 0 = c_1 + c_2 \rightarrow c_1 = -c_2$$

$$y'(0) = 1 = c_3$$

$$y''(0) = 2 = -c_2$$

The specific solution is $y = 2 - 2\cos t + \sin t$.

The limit of $y(t)$ as $t \rightarrow \infty$ is indeterminate. The result is a sinusoid.

Question 34

$$4y''' + y' + 5y = 0, \quad \begin{cases} y(0) = 2 \\ y'(0) = 1 \\ y''(0) = -1 \end{cases}$$

Find roots of characteristic polynomial:

$$4\lambda^3 + \lambda + 5 = 0$$

Note that $\lambda = -1$ solves the equation. Using polynomial division the polynomial can be rewritten as:

$$(\lambda + 1)(4\lambda^2 - 4\lambda + 5) = 0$$

$$(4\lambda^2 - 4\lambda + 5) = 0 \Rightarrow \lambda = \frac{4 \pm \sqrt{4^2 - 4 \cdot 4 \cdot 5}}{8} = \frac{1}{2} \pm i$$

The general solution is

$$y = c_1 e^{-t} + c_2 e^{t/2} \cos t + c_3 e^{t/2} \sin t$$

$$y' = -c_1 e^{-t} + c_2 \left(-e^{t/2} \sin t + \frac{1}{2} e^{t/2} \cos t \right) + c_3 \left(e^{t/2} \cos t + \frac{1}{2} e^{t/2} \sin t \right)$$

$$y' = -c_1 e^{-t} + e^{t/2} \sin t \left(\frac{c_3}{2} - c_2 \right) + e^{t/2} \cos t \left(\frac{c_2}{2} + c_3 \right)$$

$$y'' = c_1 e^{-t} + e^{t/2} \sin t \left(-c_2 - \frac{3}{4} c_3 \right) + e^{t/2} \cos t \left(c_3 - \frac{3}{4} c_2 \right)$$

Input ICs:

$$y(0) = 2 = c_1 + c_2 \tag{3}$$

$$y'(0) = 1 = -c_1 + \frac{c_2}{2} + c_3 \tag{4}$$

$$y''(0) = -1 = c_1 + c_3 - \frac{3}{4} c_2 \tag{5}$$

$$(4) + (5) : 0 = -\frac{c_2}{4} + 2c_3 \rightarrow c_2 = 8c_3$$

$$(3) + (4) : 3 = \frac{3}{2} c_2 + c_3 \rightarrow c_3 = 3 - \frac{3}{2} c_2$$

Therefore,

$$c_3 = 3 - \frac{3}{2} \cdot 8c_3 = 3 - 12c_3 \rightarrow c_3 = \frac{3}{13} \rightarrow c_2 = \frac{24}{13} \rightarrow c_1 = \frac{2}{13}$$

The specific solution is

$$y = \frac{2}{13} e^{-t} + \frac{24}{13} e^{t/2} \cos t + \frac{3}{13} e^{t/2} \sin t$$

As $t \rightarrow \infty$ the term $\frac{2}{13} e^{-t}$ approaches zero, leaving us with a sinusoid with an indeterminate limit.

Question 40

$$c_1 e^{r_1 t} + \dots + c_n e^{r_n t} = 0, \quad \forall t \quad (6)$$

(a) Multiply eq. (6) by $e^{-r_1 t}$.

$$c_1 + c_2 e^{(r_2 - r_1)t} + c_3 e^{(r_3 - r_1)t} + \dots + c_n e^{(r_n - r_1)t} = 0$$

Differentiate wrt t .

$$c_2(r_2 - r_1)e^{(r_2 - r_1)t} + c_3(r_3 - r_1)e^{(r_3 - r_1)t} + \dots + c_n(r_n - r_1)e^{(r_n - r_1)t} = 0 \quad (7)$$

(b) Multiply eq. (7) by $e^{-(r_2 - r_1)t}$.

$$c_2(r_2 - r_1) + c_3(r_3 - r_1)e^{(r_3 - r_2)t} + \dots + c_n(r_n - r_1)e^{(r_n - r_2)t} = 0$$

Differentiate wrt t .

$$c_3(r_3 - r_1)(r_3 - r_2)e^{(r_3 - r_2)t} + c_4(r_4 - r_1)(r_4 - r_2)e^{(r_4 - r_2)t} + \dots + c_n(r_n - r_1)(r_n - r_2)e^{(r_n - r_2)t} = 0 \quad (8)$$

(c) As we can see, each time we multiply by the negative version of the first exponent and differentiate, we reduce the number of terms in the equation and multiply all the remaining terms by the power of the exponent we multiplied with.

Therefore, repeating the process (the third time) for eq. (8) we get

$$c_4(r_4 - r_1)(r_4 - r_2)(r_4 - r_3)e^{(r_4 - r_3)t} + \dots + c_n(r_n - r_1)(r_n - r_2)(r_n - r_3)e^{(r_n - r_3)t} = 0$$

and for the forth time:

$$c_4(r_5 - r_1)(r_5 - r_2)(r_5 - r_3)(r_5 - r_4)e^{(r_5 - r_4)t} + \dots + c_n(r_n - r_1)(r_n - r_2)(r_n - r_3)(r_n - r_4)e^{(r_n - r_4)t} = 0$$

After $n - 1$ times, we are left with only one term:

$$c_n(r_n - r_{n-1})(r_n - r_{n-2}) \dots (r_n - r_1)e^{(r_n - r_{n-1})t} = 0$$

Because $e^{(r_n - r_{n-1})t} \neq 0$ for all t and because all r_n are different, **c_n must be zero.**

The original equation can then be reduced to

$$c_1 e^{r_1 t} + \cdots + c_{n-1} e^{r_{n-1} t} = 0$$

If we repeat the process again $n - 2$ times, we get $c_{n-1} = 0$. Then, after we rewrite the original equation again and repeat the multiplication and differentiation process $n - 3$ times, we get $c_{n-2} = 0$. Eventually, we see that all $c_k, k = 1, 2, 3 \dots$ are zero, thus proving that all $e^{r_1 t}, e^{r_2 t}, \dots, e^{r_n t}$ are linearly independent.