Notes on *Manifolds, sheaves and cohomology*

Author: 秦宇轩（Qin Yuxuan）
Last compiled at 2026-03-17

Contents

1. Notations ... 1
2. Presheaf of (E-valued) funcions ... 2
3. Sheaf valued at the empty set ... 2
4. Morphisms between stalks are alyways more convenient ... 2
5. Constant sheaves ... 2
6. Restriction of presheaves ... 2
7. Locally constant sheaves ... 2
8. Constant sheaves and trivial etale spaces ... 3
9. Ringed spaces ... 3
10. Locally ringed spaces ... 3
11. Properties of morphisms between $\mathbb{K}^n$ subspaces with C^α sheaves (Eg. 4.5) ... 3
12. Restriction is the left Kan extension (Def. 4.7) ... 3
13. Open subspaces of locally ringed space are also locally ringed (Def. 4.7) ... 3
14. C^α-premanifolds ... 4
15. Complex premanifolds ... 4
16. Manifolds ... 4
17. $(\mathbb{K}, C^\alpha_{\mathbb{K}})$ is a manifold ... 4
18. A Yoneda-style lemma of structure sheaf (Prop. 4.17) ... 4
19. Torsor ... 4
20. Cech cohomology ... 5
    20.1. Motivation ... 5
    20.2. Definition ... 6
21. Ch 5: Linearization of Manifolds ... 6
22. Ch 6: Lie Groups ... 7
    22.1. Derivative of a Lie algebra ... 7
23. Ch 7: Torsors and the First Čech Cohomology ... 8
    23.1. Torsors ... 8
        23.1.1. Torsors and principal bundles ... 9
    23.2. $H(\mathcal{U}, \mathcal{A}) \cong \check{H}(\mathcal{U}, \mathcal{A})$ ... 10
    23.3. Extend the short exact sequence ... 10
24. Ch 8: Bundles ... 11
    24.1. Vector bundles ... 11

---

1. Notations

- X, Y, Z: topological spaces.
  - τ_X: the topology of space X, i.e., the set of open sets.
- F, G: general (pre)sheaves.
- $\mathcal{O}_X$: structure sheaf associated with ringed space X.
  - For open set $U \subset X$, we use $\mathcal{O}_U$ to denote $\mathcal{O}_X|_U$.
- $\mathbb{K}$: used for denoting $\mathbb{R}$ or $\mathbb{C}$.
- α: a generalized natural number, i.e., $\alpha \in \mathbb{N}$ or $\alpha = \infty$.

- C_X^α: Sheaf of C^α functions, on a suitable space X.

Warning: All "(pre)manifolds" are either C^α real-(pre)manifolds for some α, or C^ω complex-(pre)manifolds.

To be more imformative we may use:
- (M, C_M^α): for real-manifolds;
- (M, hol_M): for complex-manifolds;
- $(M, \mathcal{O}_M)$: for general manifolds;

---

2. Presheaf of (E-valued) funcions

Given a bare set E, those presheaves P such that $P(U)$ is some kinds of subset of $\mathrm{Set}(U, E)$ for all opens U, where the restriction map is the obvious one.

3. Sheaf valued at the empty set

For sheaf F defined on a topological space X, we claim that $F(\emptyset) =$ singleton. The reason is that the empty-indexed product in any category is its (if exists) initial object, in the category $\mathbf{Set}$ this means that $\prod_\emptyset =$ singleton. Also note that $\coprod_\emptyset = \emptyset$.

So $\coprod_\emptyset = \emptyset$ is a cover of $\emptyset$, and since F is a sheaf,

$$F(\emptyset) \to \prod_\emptyset = \prod_\emptyset$$

is a equalizer, so $F(\emptyset) =$ singleton.

4. Morphisms between stalks are alyways more convenient

- The possisble reason of this phenomena is that, in general you can't go back from a set of local data to global data in presheaves, and *sheaves* are about local data, so it is difficult to define a morphism from $\mathrm{Sh}(F)$ to F for presheaves F. But the stalk F_x *is* about local data and permit you to construct an element by pure local data.
- For example, in the proof of **Example 3.30**, one can easily define a morphism $\alpha_x : \tilde{F}_x := \mathrm{Sh}(F)_x \to F_x$ by specify that, for an element $(g, \{U_i\})$ in $\tilde{F}U$, the equvilence class defined by $\left(g|_{U_x}, U_x\right)$ where $x \in U_x$. And this is indeed a well morphism, which induces a map $\tilde{F}_x \to F_x$.

5. Constant sheaves

Unfortunately, the name is a little bit confusing (at least in some kind), since for such a sheaf F and an open U, the elements of FU is *not* constant functions but *locally constant* function.
- But it can be fixed by viewing that such sheaves are the sheafification of *constant presheaves.* The proof is straightforward by **Example 3.30**.

6. Restriction of presheaves

For a presheaf $F : \tau_X^{\mathrm{op}} \to \mathrm{Set}$, we can restrict it to a open $U \subset X$ by pre-composing the inclusion functor inckusion : $\tau_U^{\mathrm{op}} \to \tau_X^{\mathrm{op}}$. This is a presheaf by definition.

7. Locally constant sheaves

Those are sheaves F such that for any open U, there exist an open cover $\{U_i\}$ on where F is a constant sheaf, i.e., FU_i is a constant sheaf for all i. This name should be interpreted as "Locally (constant sheaf)".

8. Constant sheaves and trivial etale spaces

We all agree something to be "constant" is some kinds of triviality, and under the equivalence of categories, this triviality should be preserved. And it is the fact that **constant sheaves correspond to trivial etale spaces**! To prove so, one need to prove that $\mathrm{Et}(F)$ for some constant sheaf F is a trivial etale space, *or* $\mathrm{Sh}(S, \pi)$ is a constant sheaf. And after trials one find that the latter is much easier.

- Remark of "trivial etale spaces": The most trivial etale space of a topological space X is ... X it self with the identity map. The second class of trivial etale spaces is the class of $X \times E$ where E is a set equipped with discrete topology, with the projection map.
- Add the prefix "locally" we immediately obtain the finer correspondence between *locally constant sheaves* and *locally trivial etale space*, i.e., *covering spaces.*

9. Ringed spaces

- Objects: A ringed space is a topological space X with a R-algebra sheaf $\mathcal{O}_X$, where R is a ring.
- Morphisms: A morphism between two ringed spaces $(X, \mathcal{O}_X)$ and $(Y, \mathcal{O}_Y)$ is a pair $(f, f^\flat)$:
  - $f : X \to Y$ is a continuos map;
  - $f^\flat : \mathcal{O}_Y \to f_* \mathcal{O}_X$ is a morphism between sheaves. (pronounced as "f flat")
- *Remark:* By the adjunction of $f^{-1} \dashv f_*$, to specify the $f^\flat$ is equivalent to specify $f^\natural : f^{-1}\mathcal{O}_Y \to \mathcal{O}_X$.

10. Locally ringed spaces

- Objects: Those ringed spaces such that $\mathcal{O}_{X,x}$ the stalk at x is a *local ring*.
- Morphisms: Those $(f, f^\flat, f^\natural)$ such that the morphism $f_x^\natural : \mathcal{O}_{Y,f(x)} \to \mathcal{O}_{X,x}$ send the maximal ideal into maximal ideal.

11. Properties of morphisms between $\mathbb{K}^n$ subspaces with C^α sheaves (Eg. 4.5)

For open susbets $X \subset V$ and $Y \subset W$, where both V and W are **finite dimension** $\mathbb{K}$-vector spaces and $\mathbb{K}$-ringed C^α-map $(f, f^\flat) : (X, C_X^\alpha) \to (Y, C_Y^\alpha)$:

- $(f, f^\flat)$ is automaticallt a **locally** ringed morphism.
- For all open $U \subset Y$, the morphism between structure sheaves $f_U^\flat : C_Y^\alpha U \to C_X^\alpha(f^{-1}U)$ is given by precomposition, i.e., $f^\flat = - \circ f$.

12. Restriction is the left Kan extension (Def. 4.7)

For an open subspace $i : U \to X$ and a sheaf $\mathcal{O}_X$ on X, we claim that the restriction of $\mathcal{O}_X$ on U, i.e., $\mathcal{O}_U := \mathcal{O}_X|_U$, is isomporhic to $i^{-1}\mathcal{O}_X$.

- *proof:* It's straightforward to verify that $\mathcal{O}_U$ is exactly the left Kan extension of $\mathcal{O}_X$ along i.
  - *Remark:* The canonical isomorphism is actually $i^\natural : i^{-1}\mathcal{O}_X \to O_U$, the reason is that $i^\natural$ is the correspondence of $i^\flat$ through the adjunction, but since this adjunction is about left Kan extension and precomposition, $i^\natural$ is induced by the universal property of $i^{-1}\mathcal{O}_X$, since there is a morphism $\mathcal{O}_X \to \mathcal{O}_U = \mathcal{O}_X|_U = \mathcal{O}_X \circ \mathrm{bi}$, where $\mathrm{bi} : \tau_X^{\mathrm{op}} \to \tau_U^{\mathrm{op}}$ induced by i.

13. Open subspaces of locally ringed space are also locally ringed (Def. 4.7)

By the above property, the stalk at $u \in U$ of $\mathcal{O}_U$ is isomorphic to $\mathrm{colim}_u i^{-1}\mathcal{O}_X = \mathcal{O}_{X,u}$, which by assumption is a local ring.

14. C^α-premanifolds

C^α-Premanifolds are **locally $\mathbb{K}$-ringed spaces** that are **locally isomorphic to open subspace of $\mathbb{K}^n$**, which endowed with C^α sheaf, for some n.

- Objects: A triple $(M, \mathcal{O}_M, \{U_i\})$ where
  - $(M, \mathcal{O}_M)$ is a locally $\mathbb{K}$-ringed space;
  - $\{U_i\}$ is an open cover of M equipped with a bunch of **isomorphisms** between locally $\mathbb{K}$-ringed spaces: $w_i : \left(U_i, \mathcal{O}_{U_i}\right) \to (Y, C^\alpha_Y)$, where $Y \subset \mathbb{K}^n$ open.
    - *Remarks:* Y and n may depend on i, while α fixed.
- Morphisms: Plain morphisms between locally ringed space. For such morphisms between C^α premanifolds, we usually call them **C^α-maps**.
  - *Remarks 1:* When $\alpha = \infty$, we call them **smooth** maps. When $\alpha = \omega$, we call them **analytic** maps.
  - *Remarks 2:* Isomorphisms between C^α-premanifolds are called **C^α-diffeomorphisms**.

15. Complex premanifolds

When $\mathbb{K} = \mathbb{C}$, all functions that are C^α for any α are automatically C^ω, so it is dumb to differ them.

Therefore we call them just **complex premanifolds**.

The fancy name of morphisms between complex premanifolds is "**holomorphic maps**", for isomorphisms, we use "**bi-holomorphic maps**".

16. Manifolds

- Objects: Those premanifolds whose underlying spaces are **Hausdorff** and **second countable** (admit a countable bunch of basis).
- Morphisms: Inherits from premanifolds.

17. $(\mathbb{K}, C^\alpha_{\mathbb{K}})$ is a manifold

The only needed chart is $\mathrm{id} : (\mathbb{K}, C^\alpha_{\mathbb{K}}) \to (\mathbb{K}, C^\alpha_{\mathbb{K}})$.

We alyways equip $\mathbb{K}$ with this maniold struture, if without warning.

18. A Yoneda-style lemma of structure sheaf (Prop. 4.17)

Given an arbitrary $\mathbb{K}$-premanifolds $(M, \mathcal{O}_M)$, we claim that $\mathrm{PMfd}(-, \mathbb{K}) \simeq \mathcal{O}_M$. The isomorphism is given by

$$\left[(t, t^\flat) : U \to \mathbb{K}\right] \mapsto \left[t^\flat_{\mathbb{K}}(\mathrm{id}_{\mathbb{K}})\right].$$

Note that $t^\flat : C^\alpha_{\mathbb{K}} \to t_* \mathcal{O}_U$ is a morphism between sheaves. And $\mathcal{O}_U V = \mathcal{O}_M(U \cap V)$, so $t^\flat_{\mathbb{K}}(\mathrm{id}_{\mathbb{K}}) \in \mathcal{O}_U U = \mathcal{O}_M U$, which implies these morphism is well-defined, at least in the layer of codomain.

proof: This is indeed a morphism between sheaves of $\mathbb{K}$-algebra. **TODO**

19. Torsor

A torsor is a group forgotten its identity. More precisely, for a group object G in $\mathbf{Sh}(X)$ (thus a sheaf of groups), a G-torsor is a sheaf T such that:

- There is a simply transitive G-action on T;
- T is non trivial for a open covering of X.

When $X =$ singleton we have $\mathbf{Sh}(X) \simeq \mathbf{Set}$. So at this time a torsor is just a set, and there is a bunch of group structures on T induced by a bijection from T to G. More explicitly for a fixed element $t \in T$ we can define a bijection $\alpha_t : T \to G$ as $x \mapsto g$ where g is the unique one such taht $gt = x$.

For any sheaf of groups G we obtain a category $\mathbf{Tors}(G)$, further, it is a groupoid since in $\mathbf{Set}$ any morphism between G-torsors are isomorphism.

G itself is a torsor as well, those isomorphic to G as torsors are called trivial. If $T(X)$ is non-empty then for all opens U we have $T(U) \simeq G(U)$ naturally and thus $T \simeq G$. That is, **if $\boldsymbol{T(U)}$ is non-empty then $\boldsymbol{T}$ is trivial**.

Example. The primitive function sheaf Prim_f for a holomorphic function $f : M \to \mathbb{C}$ on complex plane is an $\mathbb{C}_M$-torsor. Where $\mathbb{C}_M$ is the constant sheaf target M, that is $\mathbb{C}_M(U) := \{g : U \to M \text{ locally constant}\}$.

Example. For a bunch of discrete points $\{p_i\}$ on complex plane and a bunch of principal part f_i, we construct the Mittag-Leffler sheaf ml on $\mathbb{C}$ as a sheaf of meromorphic funtions with only possisble singular points are $\{p_i\}$, where $\mathrm{ml}(U) := \{g : U \to \mathbb{C}, \text{around } p_i \text{ has principal } f_i\}$. Then ml is a $\mathcal{O}_{\mathbb{C}}$ -torsor.

Notation: By $H^1(X, G)$ we mean the isomorphic calsses of G-torsors.

20. Cech cohomology

20.1. Motivation

See https://math.stackexchange.com/questions/4709130/what-is-the-motivation-behind-the-definition-of-cech-cohomology

Notation stolen from section "Torsor".

The Mittag-Leffler problem has a solution if and only if there is a bunch holomorphic functions $\{h_i\}$ around $\{p_i\}$ respectively such that $f_i + h_i = f_j + h_j$ for all i, j. Indeed we can construct a meromorphic function $g := f_i + h_i$ which is a solution. Conversly, if there is a solution g we define $h_i := g - f_i$. **TODO**

20.2. Definition

Definition 20.2.1 (Cech cohomology of a covering)

For a fixed open covering $\mathcal{U} = (U_i)$ of X and a sheaf of groups G, we define:

- A Cech 1-cocycles of G on $\mathcal{U}$ is a tuple $\theta = (g_{ij})$ with $g_{ij} \in G(U_{ij})$ and satisfies the *cocycle condition*:

$$g_{ij} g_{jk} = g_{ik}.$$

  The set of all 1-cocycles are denoted by $\hat{Z}(\mathcal{U}, G)$. It carries a group struture defined as component-wise multiplication.

- Two cocycles θ and θ' are called cohomologous if and only if there exist a bunch of elements $h_i \in G(U_i)$ such that

$$h_i g_{ij} = g'_{ij} h_j.$$

  (Being cohomologous is an equivalent relation.)

Finally we define the **Cech cohomology on $\mathcal{U}$**:

$$\hat{H}(\mathcal{U}, G) := \frac{\hat{Z}(\mathcal{U}, G)}{\text{cohomologous}}.$$

In general $\hat{H}(\mathcal{U}, G)$ is not a group. ♣

It's straightforward to extend this definition to global, first note that all covering of X form a category $\mathbf{Cov}(X)$ with refinements as morphisms and, $\hat{H}(-, G) : \mathbf{Cov}^{\mathrm{op}}(X) \to \mathbf{Grp}$ is a functor.

Definition 20.2.2 (Cech cohomology on X)

The "global" cohomology is defined as a colimit:

$$\hat{H}(X, G) := \operatorname{colim}_{\mathcal{U}} \hat{H}(\mathcal{U}, G).$$

♣

21. Ch 5: Linearization of Manifolds

Definition 21.1 (Rank of a morphism between manifolds)

For a morphism (i.e. smooth map) $F : M \to N$ between manifolds, define the rank of F at a point $p \in M$ as:

$$\mathrm{rk}_p(F) := \dim \operatorname{im} T_p F,$$

where $T_p F : T_p M \to T_{F(p)} N$ is the differential of F. ♣

Note that we obtain a function $\mathrm{rk}_{(-)} F : M \to \mathbb{N}$.

Proposition 21.2 ($\mathrm{rk}_{(-)} F^{-1}(\mathbb{N}_r^+)$ is open)

For morphisms between manifolds $F : M \to N$ and any given $r \in \mathbb{N}$, the following subset of M is open:

$$\{p \in M : \mathrm{rk}_p F \geq r\}.$$

♠

Proof. By restricting the domain to a smaller open subset $U \subset M$ we can assume $M \subset \mathbb{K}^n$ and $N \subset \mathbb{K}^m$, thus $T_p F$ can be identifies with an element A in $\mathrm{Hom}_{\mathbb{K}}(\mathbb{K}^n, \mathbb{K}^m)$, which can be viewd as a matrix.

So we just need to prove $\{A \in \mathrm{Mat}_{n\times m} : \mathrm{rk}\ A \geq r\}$ is open, but this can be done by checking the complement, for simplicity denoted by C, which consists of matrices whose all r-minors vanish.

Since the determinent is continuous and thus preserve the r-minor vanishing properties, C is closed. So the given set is open. □

Corollary 21.2.1

For morphism between manifolds $F : M \to N$:

1. The set $U := \{p \in M : F \text{ is an immersion at } p\} \subset M$ is open.
2. The set $V := \{p \in M : F \text{ is an submersion at } p\} \subset M$ is open.

♡

Proof. We give the proof of 1.

Since M is a manifold there is an atlas $\{U_i\}_{i\in I}$ where $U_i \cong \mathbb{K}^{n_i}$.

Note that

$$U = \bigcup_{i\in I} \{p \in U_i : F \text{ is an immersion at } p\},$$

and for all $p \in U_i$, we have $T_p M \cong T_p U_i \cong \mathbb{K}^{n_i}$, so the differential map of F is

$$T_p F : T_p M \cong T_p U_i \cong \mathbb{K}^{n_i} \to T_{F(p)} N.$$

And thus

$$\{p \in U_i : F \text{ is an immersion at } p\} = \{p \in U_i : \mathrm{rk}_p F \geq \dim T_p M = n_i\}$$

is an open subset since n_i is a fixed natural number.

Finally U is open because it is an union of open sets. □

22. Ch 6: Lie Groups

22.1. Derivative of a Lie algebra

This is (seemingly) a standard result: given a finite dimensional Lie algebra $\mathfrak{g}$, the automorphism group $\mathrm{Aut}(\mathfrak{g})$ is actually a Lie group. TODO: proof.

Definition 22.1.1 (Derivative of a Lie algebra)

Given a finite dimensional Lie algebra $\mathfrak{g}$, the Lie algebra of the Lie group $\mathrm{Aut}(\mathfrak{g})$ is called the **space of derivatives** of $\mathfrak{g}$.

♣

23. Ch 7: Torsors and the First Čech Cohomology

The exact reason why people defined the Čech Cohomology in such a complex way remains unknown to me, but if take for grant then the introduction of torsors is, in some sense, natural out of the consideration of providing an alternative definition.

23.1. Torsors

In the context of group theory, torsor is just a G-set whose action is freely transitive. And it is natural to generalize this definition to other $\mathbf{Set}$-like categories, i.e. topoi.

Torsors are geometric model of the first non-abelian cohomology of a sheaf, that means, the equivalence class of torsors of a group sheaf G is isomorphic to the first Čech cohomology group, see Section 23.2 for more.

We can form the category of torsors $\mathbf{Tors}(\mathcal{A})$ of a given group sheaf $\mathcal{A}$, and a surprising result is

Theorem 23.1.1 ($\mathbf{Tors}(\mathcal{A})$ is a groupoid)

1. $\mathbf{Tors}(\mathcal{A})$ is a groupoid;
2. An $\mathcal{A}$-torsor T is trivial iff $T(X) \neq \emptyset$

The following theorem is a good example of using the equivalence between $\mathbf{Sh}(X)$ and $\acute{\mathbf{E}}\mathbf{t}(X)$:

Theorem 23.1.2

$H^1(X, \mathcal{A}) = 0$ for locally constant sheaf $\mathcal{A}$ and simply connected, path-connected X.

We can use this theorem to prove the classical result

Proposition 23.1.3

Let $M \subset \mathbb{C}$ be open, then a holomorphic function $f : M \to \mathbb{C}$ admits primitives if M is simply-connected

Proof. We define the primitive sheaf Prim_f:

$$U \mapsto \mathrm{Prim}_f(U) := \{\text{Primitives of } f \text{ on } U\},$$

This is a $\mathbb{C}_M$-torsor with action by addition:

$$\begin{aligned} \mathbb{C}_M(U) \times \mathrm{Prim}_f(U) &\to \mathrm{Prim}_f(U), \\ (h, F) &\mapsto h + F. \end{aligned}$$

Note that the assertion is equivalent to $\mathrm{Prim}_f(M) \neq \emptyset$, which is again equivalent to Prim_f is trivial.

Now by Theorem 23.1.2, $H^1(M, \mathbb{C}_M)$ is trivial, so $\mathrm{Prim}_f \cong \mathbb{C}_M$ thus $\mathrm{Prim}_f(M) \neq \emptyset$. $\square$

Proposition 23.1.4 (Logarithm is a torsor)

Let $M \subset \mathbb{C}$ be open and $f : M \to \mathbb{C}$ be holomorphic. Define the sheaf $\log_f \in \mathbf{Sh}(M)$:

$$\log_f U := \{g \in \text{holomorphic}(U) : \exp \circ g = f\}.$$

Then
1. $\log_f$ is a $\mathbb{Z}_M$-pseudotorsor;
2. $\log_f$ is a torsor iff f does not vanish on M

♠

Proof.
1. Trivial;
2. For every point $p \in M$, there exists an open ball $p \in B_p$ such that $B_p \subset M$ because M is open. So When f does not vanish on M, $\{B_p\}_{p \in M}$ trivialises $\log_f$, which is a classical result from complex analysis[1].

□

So, when M is simply connected and f does not vanish on M, we have $H^1(M, \mathbb{Z}_M)$ is trivial, which means $\log_f(M) \neq \emptyset$.

If f admits a primitive on M, then by methods of complex analysis we can give a explicit logarithm of f, which is defined in Freitag, Theorem II.2.9. But we still need to prove the existence of primitives on a simply connected open set, which can be proved by Proposition 23.1.3.

Torsors are amazing!

23.1.1. Torsors and principal bundles

In this subsection, we generalize torsor to any category with product.

Definition 23.1.5 (Torsors in general categories)

Let $\mathcal{C}$ be a category with products, and $G \in \mathcal{C}$ a group object. A G-torsor in $\mathcal{C}$ is an object $T \in \mathcal{C}$ with an action map $\alpha : G \times T \to T$ such that the shear map $G \times T \to T \times T$ is an isomorphism.

♣

Torsors are preserved under the map from $\mathcal{C}$ to the slice catgeory $\mathcal{C}/B$ for $B \in \mathcal{C}$:

Definition 23.1.6 (Torsors over an object)

Suppose $(T, \alpha : G \times T \to T)$ is a torsor in a category $\mathcal{C}$.

For $B \in \mathcal{C}$, we have a slice map $(- \times B) : \mathcal{C} \twoheadrightarrow \mathcal{C}/B$ as:

$$C \mapsto (C \times B, \mathrm{pr}_B).$$

This slice map preserves products so $G \times B$ is also a group in $\mathcal{C}/B$.

Now we define **a G-torsor over $B \in \mathcal{C}$** is a $(G \times B)$-torsor in $\mathcal{C}/B$.

♣

[1]Non-vanishing functions admit logarithm on open balls. See, for example, Freitag Theorem II.2.9.

Definition 23.1.7 (Princial bundles)

Given a topological group G and a space B, a G-princial bundle P over B is defined as a G-torsor over B, with a local trivialization. ♣

23.2. $H(\mathcal{U}, \mathcal{A}) \cong \check{H}(\mathcal{U}, \mathcal{A})$

This is just like $H^{\text{Sing}} \cong H^{\text{Simp}}$.

Sometimes it is more convenient to compute the cohomology in another way.

A good example of using different definition to obtain fast proof of something related to the first Čech cohomology is the following proposition:

Proposition 23.2.1

$\mathcal{A} \in \mathbf{Sh}_{\mathbf{Ab}}(X) \Rightarrow \check{H}^1(X, \mathcal{A}) \in \mathbf{Ab}$ ♠

Proof. When $\mathcal{A}$ is a sheaf of abelian groups, by the definition of Čech cohomology it is obvious that $\check{H}(\mathcal{U}, \mathcal{A})$ is an abelian group.

Since $\mathbf{Ab}$ admit arbitrary colimit and

$$\check{H}^1(X, \mathcal{A}) = \operatorname{colim}_{\mathcal{U}} \check{H}^1(\mathcal{U}, \mathcal{A}),$$

this property is immediate.

□

Remark

To prove this result by the definition using torsor is unimaginable.

23.3. Extend the short exact sequence

Given a exact sequence of group sheaves on X:

$$1 \longrightarrow \mathcal{A}' \xhookrightarrow{\iota} \mathcal{A} \xrightarrow{\pi} \mathcal{A}'' \longrightarrow 1$$

We can extend it[2].

Theorem 23.3.1 (Extend exact sequence of sheaves)

For group sheaves $\mathcal{A}, \mathcal{A}'$ and $\mathcal{A}''$, the following sequence of **pointed sets** is exact:

$$\begin{tikzcd}
1 \arrow[r] & \mathcal{A}'(X) \arrow[r, hook, "\iota"] & \mathcal{A}(X) \arrow[r, "\pi"] & \mathcal{A}''(X) \arrow[dll, "\delta"'] \\
\check{H}^1(X, \mathcal{A}') \arrow[r] & \check{H}^1(X, \mathcal{A}) \arrow[r] & \check{H}^1(X, \mathcal{A}'') &
\end{tikzcd}$$

♡

[2]It can be extended to infinity, but now we just focus on the first cohomology groups.

> ⚠ **Warning**
>
> $\check{H}^1(X, \mathcal{A})$ is *not* a group in general, but it is a pointed set.

24. Ch 8: Bundles

The following example shows the power of Theorem 23.3.1:

Example. Consider this exact sequence of Lie groups:

$$0 \longrightarrow \mathbb{Z} \hookrightarrow \mathbb{C} \xrightarrow{\exp(2\pi i \cdot -)} \mathbb{C}^\times \longrightarrow 0$$

By the above theorem (TODO), for every complex manifold B, the following sequence of sheaves is also exact:

$$0 \longrightarrow \mathbb{Z}_B \hookrightarrow \mathcal{O}_B \xrightarrow{\exp(2\pi i \cdot -)} \mathcal{O}_B^\times \longrightarrow 0$$

By Theorem 23.3.1, we obtain a longer exact sequence

$$\begin{tikzcd}
0 \arrow[r] & \mathbb{Z}^{\pi_0(B)} \arrow[r, hook] & \mathcal{O}_B(B) \arrow[r, two heads, "\exp(2\pi i \cdot -)"] & \mathcal{O}_B^\times(B) \arrow[dll, "\delta"] \\
\check{H}(B, \mathbb{Z}_B) \arrow[r] & \check{H}(B, \mathcal{O}_B) \arrow[r] & \check{H}(B, \mathcal{O}_B^\times) &
\end{tikzcd}$$

So a nowhere vanishing holomorphic function admits a logarithm iff

$$\operatorname{im}(\mathcal{O}_B(B) \twoheadrightarrow \mathcal{O}_B^\times(B)) = \mathcal{O}_B^\times(B),$$

iff

$$\operatorname{im} \delta = 0,$$

iff $\check{H}(B, \mathbb{Z}_B) \to \check{H}(B, \mathcal{O}_B)$ is injective.

So when B is simply connected, every nowhere vanishing function admits a logarithm.

24.1. Vector bundles

Vector bundles are good which embodied in the following equivalence:

$$\mathbf{VectB}_X \cong \text{Finite Locally Free } \mathcal{O}_X - \mathbf{Mod},$$

and the latter is a abelian group, so is $\mathbf{VectB}_X$.

So, by the general operation, i.e. given an abelian group, we can form a K-theory of it, we thus obtain the topological K-theory.

TODO: The precise process of form a K-theory, and is it true that given any abelian category we can do this?