Notes on Riemann Surface

Author: 秦宇轩 (QIN Yuxuan)
Last compiled at 2026-04-07

Contents

1. Chapter III: Sheaves 1
  1.1. The canonical flabby resolution 2
    1.1.1. Godement resolution 2
    1.1.2. Sheaf cohomology 3
    1.1.3. Exactness of sheaves induces exactness of Godementons 3
    1.1.4. Some homological results 3
  1.2. Sheaves on manifolds 5
    1.2.1. Fine sheaves 6
    1.2.2. $\mathbf{Sh}_{\mathbf{Ab}}(X)$ is just like $\mathbf{Ab}$ 6
2. Chapter V: Basic facts about Riemann surface 7
  2.1. $(\mathcal{O}_{X,p}, \mathrm{ord}_p)$ is a valuation ring 7

Ref. Prof. Dr. Freitag's Notes.

1. Chapter III: Sheaves

To implement a homology theory for sheaves, one must first consider the concept of "exactness", and everything is just fine for presheaves, but what if we take a closer look at sheaves?

Suppose we have a sequence of sheaves and maps between them:

$$0 \longrightarrow F \xrightarrow{\alpha} G \xrightarrow{\beta} H \longrightarrow 0,$$

one may want to define the exactness of this sequence by checking if $\ker \beta = \operatorname{im} \alpha$, which works just perfect well in the case of presheaves, but **the image of a map between sheaves need not to be a sheaf again**!

> **Important**
>
> **Non-example** Consider the map in $\mathbf{Sh}(\mathbb{C})$:
>
> $$\exp : \mathcal{O} \to \mathcal{O}^*$$
>
> where $\mathcal{O}$ is the well-known ($\mathbb{Z}$-module valued, under addition) sheaf of holomophic functions on $\mathbb{C}$ and $\mathcal{O}^*$ is the multiplicative version, i.e. the ($\mathbb{Z}$-module valued, under multiplication) sheaf of holomophic functions **without zeros** on $\mathbb{C}$.
>
> Denote the "image" of exp as $K := \operatorname{im} \exp$, where $KU := \{\exp f : f \text{ is holomophic on } U\}$, we claim that it is not a sheaf.
>
> *proof by contradiction.* If K is a sheaf, then consider these two open sets which cover $\mathbb{C}^*$:
>
> $$H_1 := \left\{z : -\frac{\pi}{2} \leq \arg z \leq \pi\right\}, H_2 := \left\{z : \frac{\pi}{2} \leq \arg z \leq 2\pi\right\}.$$

Now let $f_1 = \mathrm{id} \in K(H_1)$ and $f_2 = \mathrm{id} \in K(H_2)$, we note $f_1 \mid_{H_{12}} = f_2 \mid_{H_{12}} = \mathrm{id}$ so by the definition of a sheaf there must exist a $f = \mathrm{id} \in K(H_1 \cup H_2) = K(\mathbb{C}^*)$, but this means there is a holomorphic logarithm on $\mathbb{C}^*$ which is impossible!

So we can not ccompare the kernel and the image, but need to compare the kernel with **the sheafification of image**[1]!

Definition 1.1 (Exactness of sheaves)

A sequence of sheaves

$$0 \longrightarrow F \xrightarrow{\alpha} G \xrightarrow{\beta} H \longrightarrow 0,$$

is exact iff $\mathbf{Sh}(\operatorname{im}\alpha) = \ker\beta$.

Surjectivity can be verified at the stalk level:

Theorem 1.2

A morphism of sheaves $\phi : F \to G$ is surjective if and only if the stalk map $\phi_p : F_p \to G_p$ is surjective for every p.

This can be proved by the following lemma:

Lemma 1.3

Let G be a sheaf, and F, H be two subsheaves of G, then

$$F = H \iff F_x = H_x, \forall x.$$

Proof. See [Jean, Proposition 10.17]. □

1.1. The canonical flabby resolution

1.1.1. Godement resolution

Given a presheaf $F \in \mathbf{PSh}(X)$, we can construct a associated sheaf $\mathrm{Gode}(F) \in \mathbf{Sh}(X)$ for free:

$$\mathrm{Gode}(F)(U) := \prod_{x \in U} F_x,$$

where $U \subset X$ is open.

This new sheaf is called the **Godement sheaf** of F with a obvious natural[2] injection $F \hookrightarrow \mathrm{Gode}(F)$, and it admits some good properties, such as being flabby, which, as we will see later, indicates the cohomology is trivial.

Mathematicians like iterating, so what about $\mathrm{Gode}(\mathrm{Gode}(F))$? The answer is we obtain nothing new: this sheaf is just $\mathrm{Gode}(F)$, since the germ at $\mathrm{Gode}(F)_x$, the germ at x, is just F_x.

So you **can not** get a naive resolution like

$$F \longrightarrow \mathrm{Gode}(F) \longrightarrow \mathrm{Gode}(\mathrm{Gode}(F)) \longrightarrow \mathrm{Gode}(\mathrm{Gode}(\mathrm{Gode}(F))) \longrightarrow \dots$$

[1] Contrary to the pathological behaviour of the image, the kernel is always a sheaf.

[2] Yes, Gode is a functor.

But as noted before, F is a sub(pre)sheaf of $\mathrm{Gode}(F)$ and thus peple plays with the quotient sheaf[3] $\mathrm{Gode}(F)/F$.

And now we can iterate:

$$F^{(0)} := \mathrm{Gode}(F),$$
$$F^{(n+1)} := \mathrm{Gode}\big(F^{(n)}/F^{(n-1)}\big).$$

These sheaves give the exact sequence:

$$0 \longrightarrow F \longrightarrow F^{(0)} \longrightarrow F^{(1)} \longrightarrow F^{(2)} \longrightarrow \dots$$

which is called **the Godement resolution** or the **canonical flabby resolution** (note that every $F^{(i)}$ is flabby).

1.1.2. Sheaf cohomology

Thanks to the composition-preserving property of functors, we can apply the global section functor $\Gamma : \mathbf{Sh}(X) \to \mathbf{Ab}, F \mapsto \Gamma F := F(X)$ to the Godement resolution, and obtain a **cochain complex**:

$$0 \longrightarrow \Gamma F \longrightarrow \Gamma F^{(0)} \longrightarrow \Gamma F^{(1)} \longrightarrow \Gamma F^{(2)} \longrightarrow \dots$$

This cochain complex consists of abelian groups so we can play with its cohomology groups. And this is (one of) the definition of sheaf cohomology:

$$H^n(X, F) := \frac{\ker\big(\Gamma F^{(n)} \to \Gamma F^{(n+1)}\big)}{\mathrm{im}\big(\Gamma F^{(n-1)} \to \Gamma F^{(n)}\big)}.$$

Note that $H^0(X, F) = \Gamma F = FX$.

1.1.3. Exactness of sheaves induces exactness of Godementons

Yes, for simplicity I call $F^{(n)}$ *Godementons*...

Theorem 1.1.1 (Exactness of sheaves induces exactness of Godementons)

If $F \to G \to H$ is an exact sequence of sheaves, then the induced sequence is also exact for all n:

$$F^{(n)} \to G^{(n)} \to H^{(n)}.$$

Proof.

□

1.1.4. Some homological results

Theorem 1.1.2 (Flabbiness implies exactness of Γ)

Given an exact sequence of sheaves $0 \to F \to G \to H \to 0$, if F is flabby, then the following sequence is also exact:

$$0 \to \Gamma F \to \Gamma G \to \Gamma H \to 0.$$

Proof. Zorn's Lemma...!

My thoughts: We may use the theorem that a sequence of sheaves is exact iff it is exact for for any stalks. □

[3] There are at least two definitions of "quotient sheaf", one is given by the sheafification of the co-kernel (as defined in Jean), and the another is given by the sheafification of the image (as defined in Freitag).

Theorem 1.1.3 (Flabbiness can breed!)

Given an exact sequence of sheaves $0 \to F \to G \to H \to 0$, if both F and G are flabby, then H is also flabby.

Proof. Given an open V, we need to prove the restriction $\Gamma H \to HV$ is surjective.

By restricting to V, we obtain a new exact sequence:

$$0 \to F|_V \to G|_V \to H|_V \to 0$$

and both $F|_V$ and $G|_V$ are still flabby.

Apply the former theorem to this new exact sequence we obtain a diagram:

$$\begin{tikzcd}
0 \arrow[r] \arrow[d] & \Gamma F \arrow[r, hook] \arrow[d, two heads, "\mathrm{res}"'] & \Gamma G \arrow[r, two heads] \arrow[d, two heads, "\mathrm{res}"'] & \Gamma H \arrow[r] \arrow[d, dashed, "\mathrm{res}"'] & 0 \quad \text{(Exact)} \\
0 \arrow[r] & FV \arrow[r, hook] & GV \arrow[r, two heads] & HV \arrow[r] & 0 \quad \text{(Exact)}
\end{tikzcd}$$

where all squares commutes.

Now it's clear $\Gamma H \to HV$ is surjective. □

In Theorem 1.1.2 we see flabbiness give trivial short exact sequence of global sections, and we will see this result is also true for long exact sequences of global sections, by means of splitting long exact sequences.

Lemma 1.1.4 (Splicing short exact sequence)

Let $0 \to A \to B \to C \to 0$ and $0 \to C \to D \to E \to 0$ be two exact sequence, then we can splicing them and obtain a longer one:

$$\begin{tikzcd}
0 \arrow[r] & A \arrow[r] & B \arrow[rr, dashed, "h \circ g"] \arrow[dr, "g"'] & & D \arrow[r] & E \arrow[r] & 0 \\
& & & C \arrow[ur, "h"'] \arrow[dr] & & & \\
& & 0 \arrow[ur] & & 0 & &
\end{tikzcd}$$

Proof. Trivial. □

Lemma 1.1.5 (Flabbiness implies long exact sequence)

Let

$$0 \longrightarrow F \longrightarrow F_0 \longrightarrow F_1 \longrightarrow \dots$$

be an exact sequence of flabby sheaves, then Γ preserves the exactness, i.e. the following sequence is exact:

$$0 \longrightarrow \Gamma F \longrightarrow \Gamma F_0 \longrightarrow \Gamma F_1 \longrightarrow \dots$$

Proof. From the long sequence we obtain a bunch of short exact sequence like

$$0 \longrightarrow F \longrightarrow F_0 \longrightarrow F_0/F \longrightarrow 0$$
$$0 \longrightarrow F_0/F \longrightarrow F_1 \longrightarrow F_1/F_0 \longrightarrow 0$$

And since in the first exact sequence, both F and F_0 are flabby so F_0/F is also flabby, and we apply this result to the second exact sequence, finally this procedure shows that all F_{i+1}/F_i are flabby.

Thanks to Theorem 1.1.2, after applying Γ those sequence are still exact. So by splicing them we obtain the required long exact sequence. □

Theorem 1.1.6 (Flabby sheaves admits trivial cohomology)

If F is flabby, then

$$H^n(X, F) = 0$$

for all $n > 0$.

Proof. Apply the above lemma. □

Warning

Please note that $H^0(X, F) = \Gamma F$ for all kinds of sheaves F, because F itself does not occur in the global sections complex!

Definition 1.1.7 (Acyclic Sheaf)

We call a sheaf **acyclic** if its higher order cohomology is trivial, i.e.

$$H^n(X, F) = 0$$

for all $n > 0$.

So flabby sheaves are acyclic.

This theorem shows the reason behind the name of "canonical resolution":

Theorem 1.1.8 (Canonical resolution is indeed canonical)

Let $0 \to F \to F_0 \to F_1 \to \dots$ be a acyclic resolution of sheaf F, then there is a natural isomorphism between $H^n(X, F)$ and the n-th cohomology of the following complex:

$$0 \to \Gamma F_0 \to \Gamma F_1 \to \dots$$

Proof. See Freitag. □

1.2. Sheaves on manifolds

Results in this section are mainly used for manifolds, since all of them are paracompact.

Since we have a structure sheaf $\mathcal{O}_X$ in hand for every manifold X and it is always a sheaf of rings, we only consider $\mathcal{O}_X$-modules in this section.

Theorem 1.2.1 (X paracompact $\Longrightarrow$ Γ preserves surjectivity)

Let X be paracompact and $M \to N$ a surjective morphism between $\mathcal{O}_X$-modules, then

$$\Gamma M \to \Gamma N$$

is also surjective.

This implies Γ is exact.

Theorem 1.2.2 (Γ is exact for $\mathcal{O}_X$-modules on manifold)

For a exact sequence of $\mathcal{O}_X$-modules on a paracompact space X

$$M \to N \to P,$$

Γ is exact, i.e.

$$\Gamma M \to \Gamma N \to \Gamma P$$

is also exact.

Theorem 1.2.3 ($\mathcal{O}_X$-moduels on manifolds are acyclic)

For $\mathcal{O}_X$-module F on manifold X, the cohomology is trivial:

$$H^n(X, F) = 0$$

for all $n > 0$.

1.2.1. Fine sheaves

Ref. 杜武亮 - *Introduction to Sheaf Cohomology*

Fine sheaves are another kind of acyclic sheaves on manifold.

1.2.2. $\mathbf{Sh}_{\mathbf{Ab}}(X)$ is just like $\mathbf{Ab}$

Ref. Jean Gallier - *Sheaf Cohomology*

Theorem 1.2.4 (Baer's embedding theorem)

$\mathbf{Ab}$ admits enough injectives.

Proof. See [Jean Theorem 11.6]. □

So given a $F \in \mathbf{Sh}_{\mathbf{Ab}}(X)$, we can construct a injective sheaf I with stalk-wise with a embedding $\iota : F \hookrightarrow I$, also stalk-wise.

We introduce a lemma:

Lemma 1.2.5

For an adjoint pair $L \dashv R : \mathcal{C} \to \mathcal{D}$ between abelian categories, if L is exact (i.e. preserves limits since as a left adjoint it automatically preserves colimits), then R preserves injective objects.

Now we prove the most fundmental theorem which ensures the existence of sheaf cohomology.

Theorem 1.2.6 ($\mathrm{Sh}_{\mathcal{A}}(X)$ admits enough injectives if $\mathcal{A}$ is so)

For example: the category of R-moduels $\mathbf{Sh}_R(X)$ admits enough injectives.

Proof. Based on [Jean, Proposition 13.1], see also Alkaid - *Homological Algebra and Commutative Algebra* (知乎) for more on abstract nonsense.

Now let $\mathcal{A}$ be an abelian category which admits enough injectives, $x \in X$ a point and consider this pair of functors:

$$\mathbf{Sh}_{\mathcal{A}}(X) \underset{\mathrm{sky}_x}{\overset{\mathrm{stalk}_x}{\rightleftarrows}} \mathcal{A}$$

where $\mathrm{stalk}_x : F \mapsto F_x$ is the usual stalk functor, and $\mathrm{sky}_x : A \mapsto \underline{A}$ is the skyscraper functor. It is easy to check $\mathrm{stalk}_x \dashv \mathrm{sky}_x$.

Note stalk_x is exact, then by the claim sky_x preserves injective objects.

Let $F \in \mathbf{Sh}_{\mathcal{A}}(X)$ be a sheaf, if $\mathcal{A}$ admits enough injectives, then choose a monic inclusion to an injective object $\iota_x : F_x \hookrightarrow I^x$ for every point $x \in X$. Now apply sky_x to this inclusion, and the result map $\eta_x := \mathrm{sky}_x(\iota_x)$ is also monic because as a right adjoint sky_x automatically preserves kernel.

Define $I \in \mathbf{Sh}_{\mathcal{A}}(X)$ as $I(U) := \prod_{x \in U} I^x$, this implies $I = \prod_{x \in X} \mathrm{sky}_x I^x$ and since sky_x preserves injective objects I is also an injective object.

Define $\eta : F \Rightarrow I$:

$$\eta_U : F(U) \to I(U) := F(U) \hookrightarrow \prod_{x \in U} F_x \hookrightarrow \prod_{x \in U} I^x.$$

Now $\eta_x = \iota_x$ is monic and I is injective. We are done. □

Remark

Actually, every topos admits enough injective (the subobject classifier Ω is injective, and so is the power Ω^X, see [Mac Lane & Moerdijk, IV.10]), such as $\mathbf{Sh}_{\mathbf{Set}}(X)$. However, we can not deduce $\mathbf{Sh}_{\mathcal{A}}(X)$ admit enough injective from this observation since it is **not** a topos. ($\mathbf{Sh}_{\mathcal{A}}(X)$ is an abelian category, but all non-trivial abelian categories are not cartesian-closed and thus it is not a topos.)

2. Chapter V: Basic facts about Riemann surface

2.1. $(\mathcal{O}_{X,p}, \mathrm{ord}_p)$ is a valuation ring

References of this section: Prof. Victoria Hoskins's Riemann surface course (start on `1:26:22`)

Given a connected riemann surface X with its function sheaf $\mathcal{O}_X$, then the germ stalk $\mathcal{O}_{X,p}$ at a point $p \in X$ is a local ring. With some input from complex analysis we can further claim that it is a ring with discrete valuation:

For $f \in \mathcal{O}_X(U)$ and $p \in U$, we can define the order of f at p as the order of the local normal form associated with f at p. This order function is clearly stable under the equivalence relation which identifies two holomorphic functions around p if thet share the same Tylor expansion in a open set around p, so $\mathrm{ord}_p : \mathcal{O}_{X,p} \to \mathbb{N}_\infty$ is a well-defined function.

We checked ord_p is indeed a valuation.