Q2

```
path = "C:\\Users\\LEGION\\Downloads\\btrips2015-7-1-4.rds"
bike_data <- readRDS(path)
library(ggplot2)
library(lme4)
library(dplyr)</pre>
```

(a)

```
bike_data$log_duration <- log(bike_data$duration)
anova_model <- aov(log_duration ~ station, data = bike_data)
anova_summary <- summary(anova_model)
anova_summary
```

```
## Df Sum Sq Mean Sq F value Pr(>F)

## station 50 289.9 5.797 7.699 <2e-16 ***

## Residuals 2475 1863.6 0.753


## ---

## Signif. codes: 0 '***' 0.001 '**' 0.05 '.' 0.1 ' ' 1
```

The ANOVA analysis shows a high F-value of 7.699, with an associated p-value less that 2e-16, indicating strong statistical significance. We reject the null hypothesis that the mean log trip duration is the same for all stations. This suggests that there is substantial across-station heterogeneity in the log trip duration. The variation in trip duration is significantly different across stations, implying that the station of origin plays an important role n explaining differences in trip lengths.

(b)

Sample Means of Log Duration vs Sample Size



The red dotted line in the middle represent the grand mean of the data. We can see that with small sample size, the means of log trip duration are more spread around the grand mean, indicating high variance. When the sample size increases, the mean log duration tend to be closer to the grand mean.

(c)

Sigma^2: 0.7541967

```
hierarchical_model <- lmer(log_duration ~ (1|station), data = bike_data)
summary_hierarchical <- summary(hierarchical_model)
mu <- fixef(hierarchical_model)[1]
sigma_sq <- attr(VarCorr(hierarchical_model), "sc")^2
tau_sq <- as.data.frame(VarCorr(hierarchical_model))$vcov[1]
cat("Mu:", mu, "\n")

## Mu: 2.657768

cat("Tau^2:", tau_sq, "\n")

## Tau^2: 0.1148483


cat("Sigma^2:", sigma_sq, "\n")
```

summary_hierarchical

```
## Linear mixed model fit by REML ['lmerMod']
## Formula: log_duration ~ (1 | station)
##
     Data: bike_data
##
## REML criterion at convergence: 6561.4
##
## Scaled residuals:
##
       Min
               1Q Median
                               3Q
## -3.2658 -0.6492 -0.0506 0.5580 4.1613
##
## Random effects:
## Groups
                        Variance Std. Dev.
## station (Intercept) 0.1148 0.3389
                        0.7542
                                 0.8684
## Residual
## Number of obs: 2526, groups: station, 51
##
## Fixed effects:
              Estimate Std. Error t value
## (Intercept) 2.65777
                          0.05167
                                   51.44
```

(d)

Sample Means vs Empirical Bayes Estimates




The plot demonstrates the effect of empirical Bayes (EB) shrinkage, where the EB estimates pull station-specific sample means towards the overall grand mean (indicated by the horizontal dashed line). We observe that the line formed by the points in the plot deviates from the 45-degree line (red dashed diagonal), indicating the shrinkage effect. The empirical Bayes estimates have a smaller range compared to the sample means, reflecting the tendency of EB to reduce variance by moving estimates towards the central tendency.

Regarding the amount of shrinkage, the two outliers in the upper-right and lower left of the plot experience significant shrinkage, as their sample means deviate substantially from the grand mean. In contrast, the points in the middle of the plot, where sample means are closer to the grand mean, exhibit less shrinkage.

(e)

```
alpha <- 0.05
sig hat <- as.data.frame(VarCorr(hierarchical model))$sdcor[2]</pre>
ci data <- bike data %>%
  group_by(station) %>%
  summarize(mean_log_duration = mean(log_duration),
            sample size = n() %>%
  rowwise() %>%
  mutate(
    t_{crit} = qt(1 - alpha / 2, df = sample_size - 1),
    margin_error = t_crit * sqrt(sig_hat^2 / sample_size),
    lower_ci = mean_log_duration - margin_error,
    upper_ci = mean_log_duration + margin_error
 )
ci_data <- ci_data %>% arrange(mean_log_duration)
ggplot(ci_data, aes(x = reorder(station, mean_log_duration), y = mean_log_duration)) +
  geom\ point(size = 2) +
  geom_errorbar(aes(ymin = lower_ci, ymax = upper_ci), width = 0.2, color = "blue") +
  labs(title = "95% Confidence Intervals for Each Station's Mean Log Duration",
       x = "Station (ordered by increasing mean)",
       y = "Mean Log Duration") +
  theme minimal() +
  theme(axis.text.x = element_text(angle = 90, hjust = 1))
```

95% Confidence Intervals for Each Station's Mean Log Duration



```
df_samplesize <- bike_data %>%
  group_by(station) %>%
  summarize(sample_size = n()) %>%
  arrange(sample_size)
head(df_samplesize)
```

```
## # A tibble: 6 \times 2
  station sample_size
##
   <chr>
              <int>
##
## 1 DPD-03
                      4
## 2 UW-01
                     12
## 3 UW-10
                     14
## 4 CD-01
                     15
## 5 UW-07
                     15
## 6 UW-04
                     16
```

The two outlying groups are schools 'UW-01' and 'DPD-03', as they have much larger confidence intervals, and their confidence intervals does not overlap with many other schools. We can see from the chart that these two schools have the lowest sample means, which may causes them to be the outliers.