

Introduction to Machine Learning

Lecture 3 Linear Regression - Introduction

Hongteng Xu



中國人民大學
RENMIN UNIVERSITY OF CHINA

高瓴人工智能學院
Gaoling School of Artificial Intelligence

Outline

Review

- ▶ **Linear algebra:** Basic operations of matrix and linear space
- ▶ **Statistics:** Expectation, variance, and the method of moments
- ▶ **Probability:** MLE v.s. MAP, two statistical ML paradigms

Outline

Review

- ▶ **Linear algebra:** Basic operations of matrix and linear space
- ▶ **Statistics:** Expectation, variance, and the method of moments
- ▶ **Probability:** MLE v.s. MAP, two statistical ML paradigms

Today

- ▶ Linear regression: Take polynomial regression as an example
- ▶ Pre-processing, training, and evaluation
- ▶ Model selection: AIC v.s. BIC

Polynomial Regression

A polynomial with $N - 1$ degrees:

$$p(x) = c_0 + c_1x + \dots + c_{N-1}x^{N-1} = \sum_{j=1}^N \underbrace{c_{j-1}}_{\text{Coefficient}} x^{j-1}. \quad (1)$$

Polynomial Regression

A polynomial with $N - 1$ degrees:

$$p(x) = c_0 + c_1x + \dots + c_{N-1}x^{N-1} = \sum_{j=1}^N \underbrace{c_{j-1}}_{\text{Coefficient}} x^{j-1}. \quad (1)$$

Given a sequence of numbers, *i.e.*, x_1, \dots, x_M , the mapping

$$\underbrace{[c_0, \dots, c_{N-1}]^\top}_{\mathbf{c}} \mapsto \underbrace{[p(x_1), \dots, p(x_M)]^\top}_{p(\mathbf{x})}$$

is **linear**.

Polynomial Regression

A polynomial with $N - 1$ degrees:

$$p(x) = c_0 + c_1x + \dots + c_{N-1}x^{N-1} = \sum_{j=1}^N \underbrace{c_{j-1}}_{\text{Coefficient}} x^{j-1}. \quad (1)$$

Given a sequence of numbers, *i.e.*, x_1, \dots, x_M , the mapping

$$\underbrace{[c_0, \dots, c_{N-1}]^\top}_{\mathbf{c}} \mapsto \underbrace{[p(x_1), \dots, p(x_M)]^\top}_{p(\mathbf{x})}$$


is **linear**.

► $p(\mathbf{x}) = \mathbf{A}\mathbf{c}$

► Vandermonde matrix:


$$\begin{bmatrix} 1 & x_1 & x_1^2 & \cdots & x_1^{N-1} \\ 1 & x_2 & x_2^2 & \cdots & x_2^{N-1} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & x_M & x_M^2 & \cdots & x_M^{N-1} \end{bmatrix}$$

Polynomial Regression: An Example




Malthus's principle of population.

Polynomial Regression: An Example



Polynomial Regression: An Example



Could you enumerate more cases suitable for polynomial regression?

The Rationality behind Polynomial Regression

The Rationality behind Polynomial Regression

- **Taylor Expansion:** For $f(x) : \mathbb{R} \mapsto \mathbb{R} \in \mathbb{C}^\infty$ and $a \in \mathbb{R}$,

$$\begin{aligned} f(x) &= f(a) + \frac{f^{(1)}(a)}{1!}(x-a) + \frac{f^{(2)}(a)}{2!}(x-a)^2 + \frac{f^{(3)}(a)}{3!}(x-a)^3 + \dots \\ &= \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!}(x-a)^n. \end{aligned} \tag{2}$$

The Rationality behind Polynomial Regression

- **Taylor Expansion:** For $f(x) : \mathbb{R} \mapsto \mathbb{R} \in \mathbb{C}^\infty$ and $a \in \mathbb{R}$,

$$\begin{aligned} f(x) &= f(a) + \frac{f^{(1)}(a)}{1!}(x-a) + \frac{f^{(2)}(a)}{2!}(x-a)^2 + \frac{f^{(3)}(a)}{3!}(x-a)^3 + \dots \\ &= \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!}(x-a)^n. \end{aligned} \tag{2}$$

- Polynomial function can cover and approximate a large set of functions.

The Rationality behind Polynomial Regression

- **Taylor Expansion:** For $f(x) : \mathbb{R} \mapsto \mathbb{R} \in \mathbb{C}^\infty$ and $a \in \mathbb{R}$,

$$\begin{aligned} f(x) &= f(a) + \frac{f^{(1)}(a)}{1!}(x-a) + \frac{f^{(2)}(a)}{2!}(x-a)^2 + \frac{f^{(3)}(a)}{3!}(x-a)^3 + \dots \\ &= \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!}(x-a)^n. \end{aligned} \tag{2}$$

- Polynomial function can cover and approximate a large set of functions.
- Could you enumerate the functions that cannot be fit well by polynomials?

A Naïve Learning Strategy

Given labeled data $\{(x_n, y_n)\}_{i=1}^N$, train a D -th order polynomial regression model:

$$y = \sum_{d=1}^D w_d x^{d-1} + \epsilon. \quad (3)$$

- ▶ The Vandermonde matrix $\mathbf{X} = [x_n^{d-1}] \in \mathbb{R}^{N \times D}$ and the label vector $\mathbf{y} = [y_n] \in \mathbb{R}^N$.
- ▶ Learning the model via:

$$\min_{\mathbf{w}} \underbrace{\|\mathbf{y} - \mathbf{X}\mathbf{w}\|_p^p}_{L(\mathbf{w}; \mathbf{X}, \mathbf{y})} \quad (4)$$

A Naïve Learning Strategy

Given labeled data $\{(x_n, y_n)\}_{i=1}^N$, train a D -th order polynomial regression model:

$$y = \sum_{d=1}^D w_d x^{d-1} + \epsilon. \quad (3)$$

- ▶ The Vandermonde matrix $\mathbf{X} = [x_n^{d-1}] \in \mathbb{R}^{N \times D}$ and the label vector $\mathbf{y} = [y_n] \in \mathbb{R}^N$.
- ▶ Learning the model via:

$$\min_{\mathbf{w}} \underbrace{\|\mathbf{y} - \mathbf{X}\mathbf{w}\|_p^p}_{L(\mathbf{w}; \mathbf{X}, \mathbf{y})} \quad (4)$$

- ▶ When $p = 2$, we have

$$\frac{\partial L}{\partial \mathbf{w}} = \mathbf{0}$$

A Naïve Learning Strategy

Given labeled data $\{(x_n, y_n)\}_{i=1}^N$, train a D -th order polynomial regression model:

$$y = \sum_{d=1}^D w_d x^{d-1} + \epsilon. \quad (3)$$

- ▶ The Vandermonde matrix $\mathbf{X} = [x_n^{d-1}] \in \mathbb{R}^{N \times D}$ and the label vector $\mathbf{y} = [y_n] \in \mathbb{R}^N$.
- ▶ Learning the model via:

$$\min_{\mathbf{w}} \underbrace{\|\mathbf{y} - \mathbf{X}\mathbf{w}\|_p^p}_{L(\mathbf{w}; \mathbf{X}, \mathbf{y})} \quad (4)$$

- ▶ When $p = 2$, we have

$$\frac{\partial L}{\partial \mathbf{w}} = 0 \quad \Rightarrow \quad 2\mathbf{X}^T(\mathbf{X}\mathbf{w} - \mathbf{y}) = 0$$

A Naïve Learning Strategy

Given labeled data $\{(x_n, y_n)\}_{i=1}^N$, train a D -th order polynomial regression model:

$$y = \sum_{d=1}^D w_d x^{d-1} + \epsilon. \quad (3)$$

- ▶ The Vandermonde matrix $\mathbf{X} = [x_n^{d-1}] \in \mathbb{R}^{N \times D}$ and the label vector $\mathbf{y} = [y_n] \in \mathbb{R}^N$.
- ▶ Learning the model via:

$$\min_{\mathbf{w}} \underbrace{\|\mathbf{y} - \mathbf{X}\mathbf{w}\|_p^p}_{L(\mathbf{w}; \mathbf{X}, \mathbf{y})} \quad (4)$$

- ▶ When $p = 2$, we have

$$\frac{\partial L}{\partial \mathbf{w}} = 0 \quad \Rightarrow \quad 2\mathbf{X}^T(\mathbf{X}\mathbf{w} - \mathbf{y}) = 0 \quad \Rightarrow \quad \mathbf{w}^* = (\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T\mathbf{y} \quad (5)$$

A Frequentist Viewpoint: Maximum Likelihood Estimation (MLE)

- ▶ Recall the model: $y = \underbrace{\sum_{d=1}^D w_d x^{d-1}}_{\mathbf{x}^T \mathbf{w}} + \epsilon$
- ▶ Assume noise $\epsilon \sim \mathcal{N}(0, \sigma^2)$

A Frequentist Viewpoint: Maximum Likelihood Estimation (MLE)

- ▶ Recall the model: $y = \underbrace{\sum_{d=1}^D w_d x^{d-1}}_{\mathbf{x}^T \mathbf{w}} + \epsilon$
- ▶ Assume noise $\epsilon \sim \mathcal{N}(0, \sigma^2)$

$$y - \mathbf{x}^T \mathbf{w} \sim \mathcal{N}(0, \sigma^2) \quad \Rightarrow \quad \underbrace{p(y - \mathbf{x}^T \mathbf{w})}_{p(y|\mathbf{w}, \mathbf{x})} = \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{(y - \mathbf{x}^T \mathbf{w})^2}{2\sigma^2}\right) \quad (6)$$

A Frequentist Viewpoint: Maximum Likelihood Estimation (MLE)

- ▶ Recall the model: $y = \underbrace{\sum_{d=1}^D w_d x^{d-1}}_{\mathbf{x}^T \mathbf{w}} + \epsilon$
- ▶ Assume noise $\epsilon \sim \mathcal{N}(0, \sigma^2)$

$$y - \mathbf{x}^T \mathbf{w} \sim \mathcal{N}(0, \sigma^2) \quad \Rightarrow \quad \underbrace{p(y - \mathbf{x}^T \mathbf{w})}_{p(y|\mathbf{w}, \mathbf{x})} = \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{(y - \mathbf{x}^T \mathbf{w})^2}{2\sigma^2}\right) \quad (6)$$

- ▶ Given $\{(\mathbf{x}_n, y_n)\}_{n=1}^N$, maximum likelihood estimation:

$$\max_{\mathbf{w}} \prod_{n=1}^N p(y_n - \mathbf{x}_n^T \mathbf{w}) \quad (\text{i.i.d. Assumption})$$

A Frequentist Viewpoint: Maximum Likelihood Estimation (MLE)

- ▶ Recall the model: $y = \underbrace{\sum_{d=1}^D w_d x^{d-1}}_{\mathbf{x}^T \mathbf{w}} + \epsilon$
- ▶ Assume noise $\epsilon \sim \mathcal{N}(0, \sigma^2)$

$$y - \mathbf{x}^T \mathbf{w} \sim \mathcal{N}(0, \sigma^2) \quad \Rightarrow \quad \underbrace{p(y - \mathbf{x}^T \mathbf{w})}_{p(y|\mathbf{w}, \mathbf{x})} = \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{(y - \mathbf{x}^T \mathbf{w})^2}{2\sigma^2}\right) \quad (6)$$

- ▶ Given $\{(\mathbf{x}_n, y_n)\}_{n=1}^N$, maximum likelihood estimation:

$$\begin{aligned} \max_{\mathbf{w}} \prod_{n=1}^N p(y_n - \mathbf{x}_n^T \mathbf{w}) & \quad (\text{i.i.d. Assumption}) \\ \Rightarrow \min_{\mathbf{w}} - \sum_{n=1}^N \log p(y_n - \mathbf{x}_n^T \mathbf{w}) & \quad (\text{Negative Log-likelihood}) \end{aligned}$$

A Frequentist Viewpoint: Maximum Likelihood Estimation (MLE)

- ▶ Recall the model: $y = \underbrace{\sum_{d=1}^D w_d x^{d-1}}_{\mathbf{x}^T \mathbf{w}} + \epsilon$
- ▶ Assume noise $\epsilon \sim \mathcal{N}(0, \sigma^2)$

$$y - \mathbf{x}^T \mathbf{w} \sim \mathcal{N}(0, \sigma^2) \quad \Rightarrow \quad \underbrace{p(y - \mathbf{x}^T \mathbf{w})}_{p(y|\mathbf{w}, \mathbf{x})} = \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{(y - \mathbf{x}^T \mathbf{w})^2}{2\sigma^2}\right) \quad (6)$$

- ▶ Given $\{(\mathbf{x}_n, y_n)\}_{n=1}^N$, maximum likelihood estimation:

$$\begin{aligned} & \max_{\mathbf{w}} \prod_{n=1}^N p(y_n - \mathbf{x}_n^T \mathbf{w}) && \text{(i.i.d. Assumption)} \\ \Rightarrow & \min_{\mathbf{w}} - \sum_{n=1}^N \log p(y_n - \mathbf{x}_n^T \mathbf{w}) && \text{(Negative Log-likelihood)} \\ \Rightarrow & \min_{\mathbf{w}} \frac{1}{2\sigma^2} \sum_{n=1}^N (y_n - \mathbf{x}_n^T \mathbf{w})^2 - \text{Const.} \end{aligned}$$

A Frequentist Viewpoint: Maximum Likelihood Estimation (MLE)

- ▶ Recall the model: $y = \underbrace{\sum_{d=1}^D w_d x^{d-1}}_{\mathbf{x}^T \mathbf{w}} + \epsilon$
- ▶ Assume noise $\epsilon \sim \mathcal{N}(0, \sigma^2)$

$$y - \mathbf{x}^T \mathbf{w} \sim \mathcal{N}(0, \sigma^2) \quad \Rightarrow \quad \underbrace{p(y - \mathbf{x}^T \mathbf{w})}_{p(y|\mathbf{w}, \mathbf{x})} = \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{(y - \mathbf{x}^T \mathbf{w})^2}{2\sigma^2}\right) \quad (6)$$

- ▶ Given $\{(\mathbf{x}_n, y_n)\}_{n=1}^N$, maximum likelihood estimation:

$$\begin{aligned} & \max_{\mathbf{w}} \prod_{n=1}^N p(y_n - \mathbf{x}_n^T \mathbf{w}) && \text{(i.i.d. Assumption)} \\ \Rightarrow & \min_{\mathbf{w}} - \sum_{n=1}^N \log p(y_n - \mathbf{x}_n^T \mathbf{w}) && \text{(Negative Log-likelihood)} \\ \Rightarrow & \min_{\mathbf{w}} \frac{1}{2\sigma^2} \sum_{n=1}^N (y_n - \mathbf{x}_n^T \mathbf{w})^2 - \text{Const.} && \\ \Rightarrow & \min_{\mathbf{w}} \|\mathbf{y} - \mathbf{X}^T \mathbf{w}\|_2^2 && \end{aligned} \quad (7)$$

When Facing Big Data: First-order Optimization

- ▶ Closed form solution: $\mathbf{w}^* = (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{y}$
- ▶ How many operations does it involve?

When Facing Big Data: First-order Optimization

- ▶ Closed form solution: $\mathbf{w}^* = (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{y}$
- ▶ How many operations does it involve? $\mathcal{O}(D^2 N + D^3)$

When Facing Big Data: First-order Optimization

- ▶ Closed form solution: $\mathbf{w}^* = (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{y}$
- ▶ How many operations does it involve? $\mathcal{O}(D^2 N + D^3)$

Stochastic gradient descent:

- ▶ Initialize \mathbf{w}_0 , randomly.

When Facing Big Data: First-order Optimization

- ▶ Closed form solution: $\mathbf{w}^* = (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{y}$
- ▶ How many operations does it involve? $\mathcal{O}(D^2 N + D^3)$

Stochastic gradient descent:

- ▶ Initialize \mathbf{w}_0 , randomly.
- ▶ At the t -th iteration:
 - ▶ Sample a batch of data $\mathbf{y}_B, \mathbf{X}_B$ randomly.
 - ▶ Compute the gradient $\frac{\partial L}{\partial \mathbf{w}} = 2\mathbf{X}_B^T (\mathbf{X}_B \mathbf{w}_{t-1} - \mathbf{y}_B)$.

When Facing Big Data: First-order Optimization

- ▶ Closed form solution: $\mathbf{w}^* = (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{y}$
- ▶ How many operations does it involve? $\mathcal{O}(D^2 N + D^3)$


Stochastic gradient descent:

- ▶ Initialize \mathbf{w}_0 , randomly.
- ▶ At the t -th iteration:
 - ▶ Sample a batch of data $\mathbf{y}_B, \mathbf{X}_B$ randomly.
 - ▶ Compute the gradient $\frac{\partial L}{\partial \mathbf{w}} = 2\mathbf{X}_B^T (\mathbf{X}_B \mathbf{w}_{t-1} - \mathbf{y}_B)$.
 - ▶ $\mathbf{w}_t = \mathbf{w}_{t-1} - \tau \frac{\partial L}{\partial \mathbf{w}}$.

Keypoints of The Learning Problem


- ▶ **Data preprocessing**
 - ▶ Suppress the unfairness of features
- ▶ **Evaluation**
 - ▶ Key criteria
 - ▶ Data splitting and cross-validation
- ▶ **Training**
 - ▶ Model selection

Why Do We Need Data Preprocessing?



- ▶ The x here is time (e.g., year).
- ▶ Consider a 3rd-order polynomial $y = w_1 + w_2x + w_3x^2 + w_4x^3$

Why Do We Need Data Preprocessing?



- ▶ The x here is time (e.g., year).
- ▶ Consider a 3rd-order polynomial $y = w_1 + w_2x + w_3x^2 + w_4x^3$
- ▶ $x = \mathcal{O}(10^3)$, while $x^d \mathcal{O}(10^{3d})$.
- ▶ **Numerical issue**

Data Preprocessing: Normalization

Motivation:

- ▶ Make each feature comparable on their ranges.

Data Preprocessing: Normalization

Motivation:

- ▶ Make each feature comparable on their ranges.

Principle: Given $\mathbf{X} = [\mathbf{x}_1, \dots, \mathbf{x}_D]$, for each \mathbf{x}_d

- ▶ $\|\mathbf{x}_d\|_2 = 1 \Rightarrow$ Normalization energy
- ▶ $\|\mathbf{x}_d\|_1 = 1 \Rightarrow$ Normalization absolute sum
- ▶ $\|\mathbf{x}_d\|_\infty = 1 \Rightarrow \max\{|x_{nd}|\}_{n=1}^N = 1$

Data Preprocessing: Shifting and Scaling

Motivation:

- ▶ Same range does not mean same statistics.
- ▶ Make each feature with zero mean and normalized variance.

Data Preprocessing: Shifting and Scaling

Motivation:

- ▶ Same range does not mean same statistics.
- ▶ Make each feature with zero mean and normalized variance.

Principle: Given $\mathbf{X} = [\mathbf{x}_1, \dots, \mathbf{x}_D]$, for each \mathbf{x}_d

- ▶ Estimate expectation $\hat{\mu}_d = \frac{1}{N} \sum_{n=1}^N x_{nd}$
- ▶ Estimate variance $\hat{\sigma}_d = \frac{1}{N-1} \sum_{n=1}^N (x_{nd} - \hat{\mu}_d)^2$.
- ▶ $\tilde{x}_{nd} = \frac{x_{nd} - \hat{\mu}_d}{\hat{\sigma}_d}$

Data Preprocessing: Shifting and Scaling

Motivation:

- ▶ Same range does not mean same statistics.
- ▶ Make each feature with zero mean and normalized variance.

Principle: Given $\mathbf{X} = [\mathbf{x}_1, \dots, \mathbf{x}_D]$, for each \mathbf{x}_d

- ▶ Estimate expectation $\hat{\mu}_d = \frac{1}{N} \sum_{n=1}^N x_{nd}$
- ▶ Estimate variance $\hat{\sigma}_d = \frac{1}{N-1} \sum_{n=1}^N (x_{nd} - \hat{\mu}_d)^2$.
- ▶ $\tilde{x}_{nd} = \frac{x_{nd} - \hat{\mu}_d}{\hat{\sigma}_d}$

$$x \sim \mathcal{N}(\mu, \sigma^2) \quad \Rightarrow \quad \frac{x - \mu}{\sigma} \sim \mathcal{N}(0, 1) \quad (8)$$

Data Preprocessing: Whitening

Motivation:

- ▶ Shifting and scaling assumes uncorrelated features, which is questionable.
- ▶ Make each feature with zero mean, normalized variance, and uncorrelated.

Data Preprocessing: Whitening

Motivation:

- ▶ Shifting and scaling assumes uncorrelated features, which is questionable.
- ▶ Make each feature with zero mean, normalized variance, and uncorrelated.

Principle: Given $\mathbf{X} = [\mathbf{x}_1, \dots, \mathbf{x}_D]$, for each \mathbf{x}_d

- ▶ Estimate expectation $\hat{\mu}_d = \frac{1}{N} \sum_{n=1}^N x_{nd}$ for $d = 1, \dots, D$.
- ▶ Estimate covariance matrix

$$\hat{\Sigma} = \frac{1}{N-1} (\mathbf{X} - \mathbf{1}_N \hat{\boldsymbol{\mu}}^T)^T (\mathbf{X} - \mathbf{1}_N \hat{\boldsymbol{\mu}}^T) \in \mathbb{R}^{D \times D} \quad (9)$$

- ▶ Whitening: $\tilde{\mathbf{X}} = \mathbf{X} \hat{\Sigma}^{-\frac{1}{2}}$.

Evaluation: Loss Functions and Key Criteria

Mean-square error (MSE)

► $|y - \hat{y}|^2$

Evaluation: Loss Functions and Key Criteria

Mean-square error (MSE)

- ▶ $|y - \hat{y}|^2$

Mean absolute error (MAE)

- ▶ $|y - \hat{y}|$

Evaluation: Loss Functions and Key Criteria

Mean-square error (MSE)

- ▶ $|y - \hat{y}|^2$

Mean absolute error (MAE)

- ▶ $|y - \hat{y}|$

The loss function actually is designed based on the evaluation measurement.

- ▶ MSE \Rightarrow Gaussian noise $\Rightarrow \ell_2$ -norm as loss function.

Evaluation: Loss Functions and Key Criteria

Mean-square error (MSE)

- ▶ $|y - \hat{y}|^2$

Mean absolute error (MAE)

- ▶ $|y - \hat{y}|$

The loss function actually is designed based on the evaluation measurement.

- ▶ MSE \Rightarrow Gaussian noise $\Rightarrow \ell_2$ -norm as loss function.
- ▶ MAE?

How To Evaluate The Stability of Learning Methods?

- ▶ When training data change, will we obtain the model with same/similar parameters?
- ▶ When training data change, will we train the model and make it achieve similar performance on the same testing data?

How To Evaluate The Stability of Learning Methods?

- ▶ When training data change, will we obtain the model with same/similar parameters?
- ▶ When training data change, will we train the model and make it achieve similar performance on the same testing data?

Which one is harder?

Confidence Interval and Bootstrapping

Confidence Interval

- ▶ Let X be a random sample from a probability distribution with parameter θ .
- ▶ A confidence interval of θ with confidence level α , is an interval with random endpoints $(l(X), u(X))$, such that

$$P_{\theta, \psi}(l(X) < \theta < u(X)) = \alpha, \quad \forall (\theta, \psi). \quad (10)$$

Confidence Interval and Bootstrapping

Confidence Interval

- ▶ Let X be a random sample from a probability distribution with parameter θ .
- ▶ A confidence interval of θ with confidence level α , is an interval with random endpoints $(l(X), u(X))$, such that


$$P_{\theta, \psi}(l(X) < \theta < u(X)) = \alpha, \quad \forall (\theta, \psi). \quad (10)$$

Note:

- ▶ The random interval covers the unknown θ with probability α , no matter what the true θ is.
- ▶ The true value can be out of the range.

Confidence Interval and Bootstrapping

Bootstrapping has been widely used to estimate confidence interval.



Confidence Interval and Bootstrapping

- ▶ Given bootstrapped parameters $\{\theta_n^*\}_{n=1}^N$ derived by bootstrapping
- ▶ **Percentile bootstrap:**

$$(\theta_{(\alpha/2)}^*, \theta_{(1-\alpha/2)}^*) \quad (11)$$

where $\theta_{(1-\alpha/2)}^*$ denote the $1 - \alpha/2$ percentile of the bootstrapped parameters.

Confidence Interval and Bootstrapping

- ▶ Given bootstrapped parameters $\{\theta_n^*\}_{n=1}^N$ derived by bootstrapping
- ▶ **Percentile bootstrap:**

$$(\theta_{(\alpha/2)}^*, \theta_{(1-\alpha/2)}^*) \quad (11)$$


where $\theta_{(1-\alpha/2)}^*$ denote the $1 - \alpha/2$ percentile of the bootstrapped parameters.

- ▶ **Basic bootstrap:**

$$(2\hat{\theta} - \theta_{(1-\alpha/2)}^*, 2\hat{\theta} - \theta_{(\alpha/2)}^*) \quad (12)$$

The difference between Frequentist statistics and Bayesian statistics is not so obvious as they claimed:)

Cross-validation



How To Compare Models and Select The Best?

- ▶ Good-of-fitness v.s. simplicity of the model
- ▶ Overfitting v.s. Underfitting

Akaike Information Criterion (AIC)

Motivation:

- ▶ Estimate the relative amount of information lost by a given model.
- ▶ Achieve a trade-off between good-of-fitness and model simplicity.

Akaike Information Criterion (AIC)

Motivation:

- ▶ Estimate the relative amount of information lost by a given model.
- ▶ Achieve a trade-off between good-of-fit and model simplicity.

Principle: Suppose that we have a statistical model of some data.

- ▶ Let K be the number of model parameters.
- ▶ Let $\hat{L} = \max p(\mathbf{X}|\hat{\theta})$ be the maximum likelihood for the model.
- ▶ The AIC value of the model:

$$\text{AIC} = 2K - 2 \log \hat{L} \quad (13)$$

Akaike Information Criterion (AIC)

Given M models and their AIC values $\{\text{AIC}_m\}_{m=1}^M$

- ▶ The relative likelihood of model m :

$$\exp\left(\frac{\text{AIC}_{\min} - \text{AIC}_m}{2}\right) \quad (14)$$

- ▶ It is proportional to the probability that the model m minimizes the (estimated) information loss.

Akaike Information Criterion (AIC)

Given M models and their AIC values $\{AIC_m\}_{m=1}^M$

- ▶ The relative likelihood of model m :

$$\exp\left(\frac{AIC_{\min} - AIC_m}{2}\right) \quad (14)$$

- ▶ It is proportional to the probability that the model m minimizes the (estimated) information loss.

Any drawbacks?

Bayesian Information Criterion (BIC)

Motivation:

- ▶ BIC also penalizes model complexity, and with larger penalty term.
- ▶ Consider the influence of data size.

Bayesian Information Criterion (BIC)

Motivation:

- ▶ BIC also penalizes model complexity, and with larger penalty term.
- ▶ Consider the influence of data size.

Principle: Suppose that we have a statistical model of some data.

- ▶ Let K be the number of model parameters.
- ▶ Let N be the number of data points (samples)
- ▶ Let $\hat{L} = \max p(\mathbf{X}|\hat{\theta})$ be the maximum likelihood for the model.
- ▶ The BIC value of the model:

$$\text{BIC} = K \log N - 2 \log \hat{L} \quad (15)$$

What is it reasonable?

Bayesian Information Criterion (BIC)

Suppose that θ are specific parameters of a model \mathcal{M} .

- Consider the 2nd-order Taylor expansion of the log-likelihood $\log p(\mathbf{X}|\theta, \mathcal{M})$ about the MLE $\hat{\theta}$:

$$\log p(\mathbf{X}|\theta, \mathcal{M}) \approx \underbrace{\log p(\mathbf{X}|\hat{\theta})}_{\hat{L}} - \frac{N}{2}(\theta - \hat{\theta})^T \mathbf{I}(\hat{\theta})(\theta - \hat{\theta}) \quad (16)$$

- Fisher Information Matrix:

$$\mathbf{I}(\theta) = -\mathbb{E} \left[\frac{\partial^2 \log p(\mathbf{X}|\theta)}{\partial \theta^2} \right] \quad (17)$$

Bayesian Information Criterion (BIC)

Suppose that θ are specific parameters of a model \mathcal{M} .

- ▶ Consider the 2nd-order Taylor expansion of the log-likelihood $\log p(\mathbf{X}|\theta, \mathcal{M})$ about the MLE $\hat{\theta}$:

$$\log p(\mathbf{X}|\theta, \mathcal{M}) \approx \underbrace{\log p(\mathbf{X}|\hat{\theta})}_{\hat{L}} - \frac{N}{2}(\theta - \hat{\theta})^T \mathbf{I}(\hat{\theta})(\theta - \hat{\theta}) \quad (16)$$

- ▶ Fisher Information Matrix:

$$\mathbf{I}(\theta) = -\mathbb{E} \left[\frac{\partial^2 \log p(\mathbf{X}|\theta)}{\partial \theta^2} \right] \quad (17)$$

- ▶ Why the 1st order derivation term $\frac{\partial \hat{L}}{\partial \theta}(\theta - \hat{\theta})$ is ignored?

Bayesian Information Criterion (BIC)

- We have

$$p(\mathbf{X}|\mathcal{M}) = \int p(\mathbf{X}|\boldsymbol{\theta}, \mathcal{M})p(\boldsymbol{\theta}|\mathcal{M})d\boldsymbol{\theta} \approx \left(\frac{2\pi}{N}\right)^{K/2} \underbrace{\hat{L} |\mathbf{I}(\hat{\boldsymbol{\theta}})|^{-1/2} p(\hat{\boldsymbol{\theta}})}_{\mathcal{O}(1) \text{ as } N \rightarrow \infty}$$

Bayesian Information Criterion (BIC)

- We have

$$\begin{aligned} p(\mathbf{X}|\mathcal{M}) &= \int p(\mathbf{X}|\boldsymbol{\theta}, \mathcal{M})p(\boldsymbol{\theta}|\mathcal{M})d\boldsymbol{\theta} \approx \left(\frac{2\pi}{N}\right)^{K/2} \underbrace{\hat{L} |\mathbf{I}(\hat{\boldsymbol{\theta}})|^{-1/2} p(\hat{\boldsymbol{\theta}})}_{\mathcal{O}(1) \text{ as } N \rightarrow \infty} \\ &= \exp \left(\underbrace{\log \hat{L} - \frac{K}{2} \log N}_{-0.5\text{BIC}} + \mathcal{O}(1) \right) \end{aligned} \tag{18}$$

In Summary

- ▶ Introduction of linear regression model (Take polynomial regression as an example)
- ▶ The keypoints in the whole training and testing pipeline
- ▶ Model selection: AIC and BIC

Next...

- ▶ Generalized linear regression
- ▶ Bias v.s. variance
- ▶ Regularization

Homework 1: DDL — March 17, 2022, Midnight

Python Programming

- ▶ Lab # 1 (3 Pts, Done)
- ▶ Lab # 2 (5 Pts)

Questions for Tech Report (6 Pts, ≤ 3 Pages)

- ▶ Demonstrate the equivalence of the following four claims (3 Pts):

Theorem

\mathbf{A} is of full column rank ($\text{Rank}(\mathbf{A}) = N$).

$\Leftrightarrow \mathbf{A}$ is injective.

$\Leftrightarrow [\mathbf{a}_1, \dots, \mathbf{a}_N]$ is linearly-independent.

$\Leftrightarrow \text{Null}(\mathbf{A}) = \{\mathbf{0}\}$.

- ▶ Demonstrate $\|\mathbf{x}\|_\infty \leq \|\mathbf{x}\|_2 \leq \|\mathbf{x}\|_1 \leq \sqrt{N}\|\mathbf{x}\|_2 \leq N\|\mathbf{x}\|_\infty, \forall \mathbf{x} \in \mathbb{R}^N$, and provide an illustration of the principle in the case of $N = 2$ (3 Pts).