

Introduction to Machine Learning

Lecture 4 Linear Regression - Bias, Variance, and Regularization

Hongteng Xu



中國人民大學
RENMIN UNIVERSITY OF CHINA

高瓴人工智能學院
Gaoling School of Artificial Intelligence

Outline

Review

- ▶ **Polynomial regression:** Formulation, rationality, learning
- ▶ **The devil is in the details:** Data preprocessing, stability of learning methods and models, and evaluation (loss function design)
- ▶ **Model selection:** AIC and BIC

Outline

Review

- ▶ **Polynomial regression:** Formulation, rationality, learning
- ▶ **The devil is in the details:** Data preprocessing, stability of learning methods and models, and evaluation (loss function design)
- ▶ **Model selection:** AIC and BIC

Today

- ▶ Generalized linear regression
- ▶ Bias v.s. variance (underfitting, overfitting, ...)
- ▶ Regularization methods

Revisit Polynomial Regression

$$\min_{\mathbf{w}} \|\mathbf{y} - \mathbf{X}\mathbf{w}\| \quad (1)$$

where the Vandermonde matrix $\mathbf{X} \in \mathbb{R}^{N \times D}$

$$\mathbf{X} = \begin{bmatrix} 1 & x_1 & x_1^2 & \cdots & x_1^{D-1} \\ 1 & x_2 & x_2^2 & \cdots & x_2^{D-1} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & x_N & x_N^2 & \cdots & x_N^{D-1} \end{bmatrix} \quad (2)$$

Revisit Polynomial Regression

$$\min_{\mathbf{w}} \|\mathbf{y} - \mathbf{X}\mathbf{w}\| \quad (1)$$

where the Vandermonde matrix $\mathbf{X} \in \mathbb{R}^{N \times D}$

$$\mathbf{X} = \begin{bmatrix} 1 & x_1 & x_1^2 & \cdots & x_1^{D-1} \\ 1 & x_2 & x_2^2 & \cdots & x_2^{D-1} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & x_N & x_N^2 & \cdots & x_N^{D-1} \end{bmatrix} \quad (2)$$

- The polynomial function works as a feature extractor / data representer, mapping each scalar to a D -dimensional feature vector.

Ordinary Linear Regression

- ▶ Given arbitrary N D -dimensional features $\mathbf{X} \in \mathbb{R}^{N \times D}$ and their labels $\mathbf{y} \in \mathbb{R}^N$, an ordinary linear regression is

$$\min_{\mathbf{w}} \text{loss}(\mathbf{y}, \mathbf{X}\mathbf{w}) \quad (3)$$

Ordinary Linear Regression

- ▶ Given arbitrary N D -dimensional features $\mathbf{X} \in \mathbb{R}^{N \times D}$ and their labels $\mathbf{y} \in \mathbb{R}^N$, an ordinary linear regression is

$$\min_{\mathbf{w}} \text{loss}(\mathbf{y}, \mathbf{X}\mathbf{w}) \quad (3)$$

- ▶ The design of the loss depends on the noise model

$$y = \mathbf{x}^T \mathbf{w} + \epsilon \quad (4)$$

Ordinary Linear Regression

- ▶ Given arbitrary N D -dimensional features $\mathbf{X} \in \mathbb{R}^{N \times D}$ and their labels $\mathbf{y} \in \mathbb{R}^N$, an ordinary linear regression is

$$\min_{\mathbf{w}} \text{loss}(\mathbf{y}, \mathbf{X}\mathbf{w}) \quad (3)$$

- ▶ The design of the loss depends on the noise model

$$y = \mathbf{x}^T \mathbf{w} + \epsilon \quad (4)$$

- ▶ Essentially, the learning task is maximizing $p(\mathbf{y}|\mathbf{X}, \mathbf{w})$ (MLE).

Ordinary Linear Regression

- ▶ Given arbitrary N D -dimensional features $\mathbf{X} \in \mathbb{R}^{N \times D}$ and their labels $\mathbf{y} \in \mathbb{R}^N$, an ordinary linear regression is

$$\min_{\mathbf{w}} \text{loss}(\mathbf{y}, \mathbf{X}\mathbf{w}) \quad (3)$$

- ▶ The design of the loss depends on the noise model

$$y = \mathbf{x}^T \mathbf{w} + \epsilon \quad (4)$$

- ▶ Essentially, the learning task is maximizing $p(\mathbf{y}|\mathbf{X}, \mathbf{w})$ (MLE).
- ▶ \mathbf{X} are random variables, and a linear regression is interested in the expected value of Y conditioned on X based on a linear predictor, i.e., $\mathbb{E}[Y|X]$.

Random Variables and Instances/Samples

- ▶ X : a random variable (r.v.) yielding a distribution P_X , where the random variable can be defined in a D -dimensional space \mathcal{X} .

Random Variables and Instances/Samples

- ▶ X : a random variable (r.v.) yielding a distribution P_X , where the random variable can be defined in a D -dimensional space \mathcal{X} .
- ▶ Accordingly, P_X is defined on the space \mathcal{X} , and sometimes is denoted as $P_{\mathcal{X}}$.

Random Variables and Instances/Samples

- ▶ X : a random variable (r.v.) yielding a distribution P_X , where the random variable can be defined in a D -dimensional space \mathcal{X} .
- ▶ Accordingly, P_X is defined on the space \mathcal{X} , and sometimes is denoted as $P_{\mathcal{X}}$.
- ▶ $\mathbf{x} \in \mathcal{X}$ is a sample in the space, which can be treated as an instance of the r.v. X , i.e., $\mathbf{x} \sim P_X$.

Random Variables and Instances/Samples

- ▶ X : a random variable (r.v.) yielding a distribution P_X , where the random variable can be defined in a D -dimensional space \mathcal{X} .
- ▶ Accordingly, P_X is defined on the space \mathcal{X} , and sometimes is denoted as $P_{\mathcal{X}}$.
- ▶ $\mathbf{x} \in \mathcal{X}$ is a sample in the space, which can be treated as an instance of the r.v. X , i.e., $\mathbf{x} \sim P_X$.
- ▶ Multiple instances \mathbf{x} 's often lead to a matrix \mathbf{X} , and similarly, we denote $\mathbf{X} \sim P_X$.

Random Variables and Instances/Samples

- ▶ X : a random variable (r.v.) yielding a distribution P_X , where the random variable can be defined in a D -dimensional space \mathcal{X} .
- ▶ Accordingly, P_X is defined on the space \mathcal{X} , and sometimes is denoted as $P_{\mathcal{X}}$.
- ▶ $\mathbf{x} \in \mathcal{X}$ is a sample in the space, which can be treated as an instance of the r.v. X , i.e., $\mathbf{x} \sim P_X$.
- ▶ Multiple instances \mathbf{x} 's often lead to a matrix \mathbf{X} , and similarly, we denote $\mathbf{X} \sim P_X$.
- ▶ $\mathbb{E}_{P_X}[X]$ and $\mathbb{V}_{P_X}[X]$ are expectation and variance of the r.v. X .

From Ordinary LR to Generalized Linear Model (GLM)

- ▶ GLM is a natural extension of ordinary linear regression, which consists of
 1. An **exponential family** of probability distributions to generate the output.
 2. A **linear predictor** $\eta = X\beta$
 3. A **link function** g : $\mathbb{E}[Y|X] = \mu = g^{-1}(\eta)$.

From Ordinary LR to Generalized Linear Model (GLM)

- ▶ GLM is a natural extension of ordinary linear regression, which consists of
 1. An **exponential family** of probability distributions to generate the output.
 2. A **linear predictor** $\eta = X\beta$
 3. A **link function** $g: \mathbb{E}[Y|X] = \mu = g^{-1}(\eta)$.
- ▶ The predictor merging input information is linear.
- ▶ The link function connecting the prediction and the conditional expectation can be nonlinear (That is why the model is called GLM).

Exponential Family of Probability Distributions

- ▶ A parametric distribution $p_X(\mathbf{x}|\boldsymbol{\theta})$ having the following form:

$$p_X(\mathbf{x}|\boldsymbol{\theta}) = h(\mathbf{x}) \exp(\langle \boldsymbol{\eta}(\boldsymbol{\theta}), \mathbf{T}(\mathbf{x}) \rangle - A(\boldsymbol{\theta})). \quad (5)$$

- ▶ $\mathbf{T}(\mathbf{x}) : \mathbb{R}^D \mapsto \mathbb{R}^S$: **Sufficient statistic** of the distribution, a function of the data holding all information of the data.

$$I(\boldsymbol{\theta}; \mathbf{T}(\mathbf{x})) = I(\boldsymbol{\theta}; \mathbf{x}) \quad (6)$$

- ▶ For **Likelihood ratio**:

$$\frac{p_X(\mathbf{x}|\boldsymbol{\theta}_1)}{p_X(\mathbf{x}|\boldsymbol{\theta}_2)} = \frac{p_X(\mathbf{y}|\boldsymbol{\theta}_1)}{p_X(\mathbf{y}|\boldsymbol{\theta}_2)} \quad \text{if} \quad \mathbf{T}(\mathbf{x}) = \mathbf{T}(\mathbf{y}) \quad (7)$$

- ▶ $S = \dim(\mathbf{T}(\mathbf{x})) = \dim(\boldsymbol{\theta})$.

Exponential Family of Probability Distributions

- ▶ A parametric distribution $p_X(\mathbf{x}|\boldsymbol{\theta})$ having the following form:

$$p_X(\mathbf{x}|\boldsymbol{\theta}) = h(\mathbf{x}) \exp(\langle \boldsymbol{\eta}(\boldsymbol{\theta}), \mathbf{T}(\mathbf{x}) \rangle - A(\boldsymbol{\theta})). \quad (8)$$

- ▶ $\boldsymbol{\eta}(\boldsymbol{\theta}) : \mathbb{R}^S \mapsto \mathbb{R}^S$ is **natural parameter**.
- ▶ The natural parameter space, $\{\boldsymbol{\eta} | p_X(\mathbf{x}|\boldsymbol{\theta}) \leq \infty\}$, is a convex set.

Exponential Family of Probability Distributions

- ▶ A parametric distribution $p_X(\mathbf{x}|\boldsymbol{\theta})$ having the following form:

$$p_X(\mathbf{x}|\boldsymbol{\theta}) = h(\mathbf{x}) \exp(\langle \boldsymbol{\eta}(\boldsymbol{\theta}), \mathbf{T}(\mathbf{x}) \rangle - A(\boldsymbol{\theta})). \quad (8)$$

- ▶ $\boldsymbol{\eta}(\boldsymbol{\theta}) : \mathbb{R}^S \mapsto \mathbb{R}^S$ is **natural parameter**.
- ▶ The natural parameter space, $\{\boldsymbol{\eta} | p_X(\mathbf{x}|\boldsymbol{\theta}) \leq \infty\}$, is a convex set.
- ▶ $A(\boldsymbol{\theta}) : \mathbb{R}^S \mapsto \mathbb{R}$ is called the **log-partition function** because it is the logarithm of a normalization factor

$$A(\boldsymbol{\theta}) = \log \left(\int_{\mathbf{x} \in \mathcal{X}} h(\mathbf{x}) \exp(\langle \boldsymbol{\eta}(\boldsymbol{\theta}), \mathbf{T}(\mathbf{x}) \rangle) d\mathbf{x} \right) \quad (9)$$

- ▶ The moments (including mean and variance) of $\mathbf{T}(\mathbf{x})$ can be derived simply by differentiating $A(\boldsymbol{\theta})$.

Exponential Family of Probability Distributions

- ▶ A parametric distribution $p_X(\mathbf{x}|\boldsymbol{\theta})$ having the following form:

$$p_X(\mathbf{x}|\boldsymbol{\theta}) = h(\mathbf{x}) \exp(\langle \boldsymbol{\eta}(\boldsymbol{\theta}), \mathbf{T}(\mathbf{x}) \rangle - A(\boldsymbol{\theta})). \quad (8)$$

- ▶ $\boldsymbol{\eta}(\boldsymbol{\theta}) : \mathbb{R}^S \mapsto \mathbb{R}^S$ is **natural parameter**.
- ▶ The natural parameter space, $\{\boldsymbol{\eta} | p_X(\mathbf{x}|\boldsymbol{\theta}) \leq \infty\}$, is a convex set.
- ▶ $A(\boldsymbol{\theta}) : \mathbb{R}^S \mapsto \mathbb{R}$ is called the **log-partition function** because it is the logarithm of a normalization factor

$$A(\boldsymbol{\theta}) = \log \left(\int_{\mathbf{x} \in \mathcal{X}} h(\mathbf{x}) \exp(\langle \boldsymbol{\eta}(\boldsymbol{\theta}), \mathbf{T}(\mathbf{x}) \rangle) d\mathbf{x} \right) \quad (9)$$

- ▶ The moments (including mean and variance) of $\mathbf{T}(\mathbf{x})$ can be derived simply by differentiating $A(\boldsymbol{\theta})$.
- ▶ $h(\mathbf{x}) : \mathbb{R}^D \mapsto \mathbb{R}$ is a non-negative integratable function.

Exponential Family of Probability Distributions

Useful Properties

- ▶ For i.i.d. data $\mathbf{X} = \{\mathbf{x}_n\}_{n=1}^N$, $\mathbf{T}(\mathbf{X}) = \sum_{n=1}^N \mathbf{T}(\mathbf{x}_n)$.

Exponential Family of Probability Distributions

Useful Properties

- For i.i.d. data $\mathbf{X} = \{\mathbf{x}_n\}_{n=1}^N$, $\mathbf{T}(\mathbf{X}) = \sum_{n=1}^N \mathbf{T}(\mathbf{x}_n)$.

$$\begin{aligned} p(\mathbf{X}|\boldsymbol{\theta}) &= \prod_{n=1}^N p(\mathbf{x}_n|\boldsymbol{\theta}) \\ &= \prod_{n=1}^N h(\mathbf{x}_n) \exp(\langle \boldsymbol{\eta}(\boldsymbol{\theta}), \mathbf{T}(\mathbf{x}_n) \rangle - A(\boldsymbol{\theta})) \end{aligned}$$

Exponential Family of Probability Distributions

Useful Properties

- For i.i.d. data $\mathbf{X} = \{\mathbf{x}_n\}_{n=1}^N$, $\mathbf{T}(\mathbf{X}) = \sum_{n=1}^N \mathbf{T}(\mathbf{x}_n)$.

$$\begin{aligned} p(\mathbf{X}|\boldsymbol{\theta}) &= \prod_{n=1}^N p(\mathbf{x}_n|\boldsymbol{\theta}) \\ &= \prod_{n=1}^N h(\mathbf{x}_n) \exp(\langle \boldsymbol{\eta}(\boldsymbol{\theta}), \mathbf{T}(\mathbf{x}_n) \rangle - A(\boldsymbol{\theta})) \\ &= \left(\prod_{n=1}^N h(\mathbf{x}_n) \right) \exp \left(\langle \boldsymbol{\eta}(\boldsymbol{\theta}), \sum_{n=1}^N \mathbf{T}(\mathbf{x}_n) \rangle - NA(\boldsymbol{\theta}) \right) \end{aligned} \tag{10}$$

Exponential Family of Probability Distributions

Useful Properties

- ▶ Exponential families have **conjugate priors**
 - ▶ In Bayesian probability theory, if the posterior distribution $p(\theta|\mathbf{x})$ is in the same distribution family as the prior distribution $p(\theta)$, the prior and posterior are called **conjugate distributions**, and the prior is called a **conjugate prior** for the likelihood function $p(\mathbf{x}|\theta)$.

Exponential Family of Probability Distributions

Useful Properties

- ▶ Exponential families have **conjugate priors**
 - ▶ In Bayesian probability theory, if the posterior distribution $p(\theta|\mathbf{x})$ is in the same distribution family as the prior distribution $p(\theta)$, the prior and posterior are called **conjugate distributions**, and the prior is called a **conjugate prior** for the likelihood function $p(\mathbf{x}|\theta)$.
- ▶ The posterior distribution of an exponential-family random variable with a conjugate prior can always be written in closed form. (Important for efficient Bayesian machine learning)

Typical Exponential Families and Their Conjugate Priors

- ▶ Normal distribution
- ▶ Exponential distribution
- ▶ Gamma distribution
- ▶ Bernoulli distribution
- ▶ Beta distribution
- ▶ Poisson distribution
- ▶ Categorical distribution
- ▶ Geometric distribution
- ▶ Multinormal distribution

https://en.wikipedia.org/wiki/Exponential_family

Typical Exponential Families and Their Conjugate Priors

- ▶ Normal distribution \Rightarrow Normal/Gamma/Normal-Gamma
- ▶ Exponential distribution \Rightarrow Gamma
- ▶ Gamma distribution \Rightarrow Gamma
- ▶ Bernoulli distribution \Rightarrow Beta
- ▶ Poisson distribution \Rightarrow Gamma
- ▶ Categorical distribution \Rightarrow Dirichlet
- ▶ Geometric distribution \Rightarrow Beta
- ▶ Multinomial distribution \Rightarrow Dirichlet

https://en.wikipedia.org/wiki/Conjugate_prior

Revisit Ordinary LR from A Viewpoint of GLM

$$y = \mathbf{x}^T \mathbf{w} + \epsilon \quad (11)$$

- ▶ Exponential family: $y \sim \mathcal{N}(\mathbf{x}^T \mathbf{w}, \sigma^2)$
- ▶ Linear predictor: $\eta = \mathbf{x}^T \mathbf{w}$
- ▶ Identity link function: $g^{-1}(\eta) = \eta$

Revisit Ordinary LR from A Viewpoint of GLM

$$y = \mathbf{x}^T \mathbf{w} + \epsilon \quad (11)$$

- ▶ Exponential family: $y \sim \mathcal{N}(\mathbf{x}^T \mathbf{w}, \sigma^2)$
- ▶ Linear predictor: $\eta = \mathbf{x}^T \mathbf{w}$
- ▶ Identity link function: $g^{-1}(\eta) = \eta$

The selection of link function is highly relevant to the distribution type: for $y = g^{-1}(\mathbf{x}^T \mathbf{w}) \sim P$

- ▶ Poisson distribution $\Leftrightarrow g(\mu) = \log \mu$
- ▶ Gamma distribution $\Leftrightarrow g(\mu) = \frac{1}{\mu}$
- ▶ Bernoulli, Categorical, Multinomial $\Leftrightarrow g(\mu) = \log \frac{\mu}{1-\mu}$ (Logit)

https://en.wikipedia.org/wiki/Generalized_linear_model

The Bias of Estimation

- ▶ Given a statistical model with parameter θ .
- ▶ Given a set of observed data x 's and each $x \sim P_\theta(x) = P(x|\theta)$.
- ▶ The estimation of θ based on the data points is denoted as $\hat{\theta}$.

The Bias of Estimation

- ▶ Given a statistical model with parameter θ .
- ▶ Given a set of observed data x 's and each $x \sim P_\theta(x) = P(x|\theta)$.
- ▶ The estimation of θ based on the data points is denoted as $\hat{\theta}$.
- ▶ The **bias** of $\hat{\theta}$ relative to θ is

$$\text{Bias}(\hat{\theta}, \theta) = \text{Bias}_\theta[\hat{\theta}] = \mathbb{E}_{x|\theta}[\hat{\theta}] - \theta = \mathbb{E}_{x|\theta}[\hat{\theta} - \theta], \quad (12)$$

which measures the difference between the estimator's expected value and the ground truth.

The Bias of Estimation

- ▶ Given a statistical model with parameter θ .
- ▶ Given a set of observed data x 's and each $x \sim P_\theta(x) = P(x|\theta)$.
- ▶ The estimation of θ based on the data points is denoted as $\hat{\theta}$.
- ▶ The **bias** of $\hat{\theta}$ relative to θ is

$$\text{Bias}(\hat{\theta}, \theta) = \text{Bias}_\theta[\hat{\theta}] = \mathbb{E}_{x|\theta}[\hat{\theta}] - \theta = \mathbb{E}_{x|\theta}[\hat{\theta} - \theta], \quad (12)$$

which measures the difference between the estimator's expected value and the ground truth.

- ▶ How to understand the notation $\mathbb{E}_{x|\theta}$?

The Bias of Estimation

- ▶ Given a statistical model with parameter θ .
- ▶ Given a set of observed data x 's and each $x \sim P_\theta(x) = P(x|\theta)$.
- ▶ The estimation of θ based on the data points is denoted as $\hat{\theta}$.
- ▶ The **bias** of $\hat{\theta}$ relative to θ is

$$\text{Bias}(\hat{\theta}, \theta) = \text{Bias}_\theta[\hat{\theta}] = \mathbb{E}_{x|\theta}[\hat{\theta}] - \theta = \mathbb{E}_{x|\theta}[\hat{\theta} - \theta], \quad (12)$$

which measures the difference between the estimator's expected value and the ground truth.

- ▶ How to understand the notation $\mathbb{E}_{x|\theta}$?
- ▶ $\text{Bias}(\hat{\theta}, \theta) = 0 \Leftrightarrow$ The estimator $\hat{\theta}$ is unbiased.

Toy Example 1: Is average an unbiased estimation of mean?

Given i.i.d. random variables $\{X_n\}_{n=1}^N$, with expectation μ and variance σ^2 .

- ▶ Sample average $\hat{\mu} = \frac{1}{N} \sum_{n=1}^N X_n$.

Toy Example 1: Is average an unbiased estimation of mean?

Given i.i.d. random variables $\{X_n\}_{n=1}^N$, with expectation μ and variance σ^2 .

- ▶ Sample average $\hat{\mu} = \frac{1}{N} \sum_{n=1}^N X_n$.
- ▶ We have

$$\begin{aligned}\text{Bias}(\hat{\mu}, \mu) &= \mathbb{E}_{X|\mu}[\hat{\mu}] - \mu \\ &= \mathbb{E}_{X|\mu}\left[\frac{1}{N} \sum_{n=1}^N X_n\right] - \mu \\ &= \frac{1}{N} \sum_{n=1}^N \mathbb{E}_{X|\mu}[X_n] - \mu \\ &= \frac{N\mu}{N} - \mu = 0\end{aligned}\tag{13}$$

Toy Example 2: What is the unbiased estimation of variance?

- ▶ Sample variance $\hat{\sigma}^2 = \frac{1}{N} \sum_{n=1}^N (X_n - \hat{\mu})^2$.

Toy Example 2: What is the unbiased estimation of variance?

- Sample variance $\hat{\sigma}^2 = \frac{1}{N} \sum_{n=1}^N (X_n - \hat{\mu})^2$.

$$\text{Bias}(\hat{\sigma}^2, \sigma^2) = \mathbb{E}_{X|\sigma^2}[\hat{\sigma}^2] - \sigma^2 = \mathbb{E}_{X|\sigma^2}\left[\frac{1}{N} \sum_{n=1}^N (X_n - \hat{\mu})^2\right] - \sigma^2$$

Toy Example 2: What is the unbiased estimation of variance?

- Sample variance $\hat{\sigma}^2 = \frac{1}{N} \sum_{n=1}^N (X_n - \hat{\mu})^2$.

$$\begin{aligned}\text{Bias}(\hat{\sigma}^2, \sigma^2) &= \mathbb{E}_{X|\sigma^2}[\hat{\sigma}^2] - \sigma^2 = \mathbb{E}_{X|\sigma^2}\left[\frac{1}{N} \sum_{n=1}^N (X_n - \hat{\mu})^2\right] - \sigma^2 \\ &= \mathbb{E}_{X|\sigma^2}\left[\frac{1}{N} \sum_{n=1}^N [(X_n - \mu) - (\hat{\mu} - \mu)]^2\right] - \sigma^2 \\ &= \mathbb{E}_{X|\sigma^2}\left[\frac{1}{N} \sum_{n=1}^N [(X_n - \mu)^2 - 2(X_n - \mu)(\hat{\mu} - \mu) + (\hat{\mu} - \mu)^2]\right] - \sigma^2\end{aligned}$$

Toy Example 2: What is the unbiased estimation of variance?

- ▶ Sample variance $\hat{\sigma}^2 = \frac{1}{N} \sum_{n=1}^N (X_n - \hat{\mu})^2$.

$$\begin{aligned}\text{Bias}(\hat{\sigma}^2, \sigma^2) &= \mathbb{E}_{X|\sigma^2}[\hat{\sigma}^2] - \sigma^2 = \mathbb{E}_{X|\sigma^2}\left[\frac{1}{N} \sum_{n=1}^N (X_n - \hat{\mu})^2\right] - \sigma^2 \\&= \mathbb{E}_{X|\sigma^2}\left[\frac{1}{N} \sum_{n=1}^N [(X_n - \mu) - (\hat{\mu} - \mu)]^2\right] - \sigma^2 \\&= \mathbb{E}_{X|\sigma^2}\left[\frac{1}{N} \sum_{n=1}^N [(X_n - \mu)^2 - 2(X_n - \mu)(\hat{\mu} - \mu) + (\hat{\mu} - \mu)^2]\right] - \sigma^2 \\&= \mathbb{E}_{X|\sigma^2}\left[\frac{1}{N} \sum_{n=1}^N (X_n - \mu)^2 - \frac{2}{N}(\hat{\mu} - \mu) \sum_{n=1}^N (X_n - \mu) + (\hat{\mu} - \mu)^2\right] - \sigma^2 \\&= \mathbb{E}_{X|\sigma^2}\left[\frac{1}{N} \sum_{n=1}^N (X_n - \mu)^2 - 2(\hat{\mu} - \mu)^2 + (\hat{\mu} - \mu)^2\right] - \sigma^2 \\&= \underbrace{\mathbb{E}_{X|\sigma^2}\left[\frac{1}{N} \sum_{n=1}^N (X_n - \mu)^2\right]}_{\sigma^2} - \underbrace{\mathbb{E}_{X|\sigma^2}[(\hat{\mu} - \mu)^2]}_{\frac{1}{N}\sigma^2} - \sigma^2.\end{aligned}$$

Toy Example 2: What is the unbiased estimation of variance?

- ▶ Sample variance $\hat{\sigma}^2 = \frac{1}{N} \sum_{n=1}^N (X_n - \hat{\mu})^2$.

$$\begin{aligned} \text{Bias}(\hat{\sigma}^2, \sigma^2) &= \mathbb{E}_{X|\sigma^2}[\hat{\sigma}^2] - \sigma^2 = \mathbb{E}_{X|\sigma^2}\left[\frac{1}{N} \sum_{n=1}^N (X_n - \hat{\mu})^2\right] - \sigma^2 \\ &= \mathbb{E}_{X|\sigma^2}\left[\frac{1}{N} \sum_{n=1}^N [(X_n - \mu) - (\hat{\mu} - \mu)]^2\right] - \sigma^2 \\ &= \mathbb{E}_{X|\sigma^2}\left[\frac{1}{N} \sum_{n=1}^N [(X_n - \mu)^2 - 2(X_n - \mu)(\hat{\mu} - \mu) + (\hat{\mu} - \mu)^2]\right] - \sigma^2 \\ &= \mathbb{E}_{X|\sigma^2}\left[\frac{1}{N} \sum_{n=1}^N (X_n - \mu)^2 - \frac{2}{N}(\hat{\mu} - \mu) \sum_{n=1}^N (X_n - \mu) + (\hat{\mu} - \mu)^2\right] - \sigma^2 \quad (14) \\ &= \mathbb{E}_{X|\sigma^2}\left[\frac{1}{N} \sum_{n=1}^N (X_n - \mu)^2 - 2(\hat{\mu} - \mu)^2 + (\hat{\mu} - \mu)^2\right] - \sigma^2 \\ &= \underbrace{\mathbb{E}_{X|\sigma^2}\left[\frac{1}{N} \sum_{n=1}^N (X_n - \mu)^2\right]}_{\sigma^2} - \underbrace{\mathbb{E}_{X|\sigma^2}[(\hat{\mu} - \mu)^2]}_{\frac{1}{N}\sigma^2} - \sigma^2. \end{aligned}$$

- ▶ The unbiased estimation is $\hat{\sigma}^2 = \frac{1}{N-1} \sum_{n=1}^N (X_n - \hat{\mu})^2$

The Variance of Estimation

- ▶ Given a statistical model with parameter θ .
- ▶ Given a set of observed data x 's and each $x \sim P_\theta(x) = P(x|\theta)$.
- ▶ The estimation of θ based on the data points is denoted as $\hat{\theta}$.

The Variance of Estimation

- ▶ Given a statistical model with parameter θ .
- ▶ Given a set of observed data x 's and each $x \sim P_\theta(x) = P(x|\theta)$.
- ▶ The estimation of θ based on the data points is denoted as $\hat{\theta}$.
- ▶ The **variance** of $\hat{\theta}$ is

$$\mathbb{V}[\hat{\theta}] = \mathbb{E}_{x|\theta}[(\hat{\theta} - \mathbb{E}_{x|\theta}[\hat{\theta}])^2] \quad (15)$$

The Trade-off Between Bias and Variance

- ▶ Suppose that \mathbf{w} is the ground truth parameter of a linear model
- ▶ A set of data (\mathbf{X}, \mathbf{y}) are observed and yield

$$y = \underbrace{\mathbf{x}^T \mathbf{w}}_{f_{\mathbf{w}}(\mathbf{x})} + \epsilon, \quad \epsilon \sim \mathcal{N}(0, \sigma^2) \quad (16)$$

- ▶ $\hat{\mathbf{w}}$ is the estimator obtained based on the data.

$$\text{MSE} = \mathbb{E}_{\mathbf{y}|\mathbf{x}, \mathbf{w}}[(y - \hat{y})^2] = \mathbb{E}[(f_{\mathbf{w}}(\mathbf{x}) + \epsilon - f_{\hat{\mathbf{w}}}(\mathbf{x}))^2] = \sigma^2 + \mathbb{E}[(f_{\mathbf{w}}(\mathbf{x}) - f_{\hat{\mathbf{w}}}(\mathbf{x}))^2]$$

The Trade-off Between Bias and Variance

- ▶ Suppose that \mathbf{w} is the ground truth parameter of a linear model
- ▶ A set of data (\mathbf{X}, \mathbf{y}) are observed and yield

$$y = \underbrace{\mathbf{x}^T \mathbf{w}}_{f_{\mathbf{w}}(\mathbf{x})} + \epsilon, \quad \epsilon \sim \mathcal{N}(0, \sigma^2) \quad (16)$$

- ▶ $\hat{\mathbf{w}}$ is the estimator obtained based on the data.

$$\begin{aligned} \text{MSE} &= \mathbb{E}_{\mathbf{y}|\mathbf{x}, \mathbf{w}}[(y - \hat{y})^2] = \mathbb{E}[(f_{\mathbf{w}}(\mathbf{x}) + \epsilon - f_{\hat{\mathbf{w}}}(\mathbf{x}))^2] = \sigma^2 + \mathbb{E}[(f_{\mathbf{w}}(\mathbf{x}) - f_{\hat{\mathbf{w}}}(\mathbf{x}))^2] \\ &= \sigma^2 + \mathbb{E}[(f_{\mathbf{w}}(\mathbf{x}) - \mathbb{E}[f_{\hat{\mathbf{w}}}(\mathbf{x})] + \mathbb{E}[f_{\hat{\mathbf{w}}}(\mathbf{x})] - f_{\hat{\mathbf{w}}}(\mathbf{x}))^2] \\ &= \sigma^2 + \mathbb{E}[(f_{\mathbf{w}}(\mathbf{x}) - \mathbb{E}[f_{\hat{\mathbf{w}}}(\mathbf{x})])^2] + \underbrace{\mathbb{E}[2(f_{\mathbf{w}}(\mathbf{x}) - \mathbb{E}[f_{\hat{\mathbf{w}}}(\mathbf{x})])(\mathbb{E}[f_{\hat{\mathbf{w}}}(\mathbf{x})] - f_{\hat{\mathbf{w}}}(\mathbf{x}))]}_{=0} \\ &\quad + \mathbb{E}[(\mathbb{E}[f_{\hat{\mathbf{w}}}(\mathbf{x})] - f_{\hat{\mathbf{w}}}(\mathbf{x}))^2] \end{aligned}$$

The Trade-off Between Bias and Variance





- ▶ Suppose that \mathbf{w} is the ground truth parameter of a linear model
- ▶ A set of data (\mathbf{X}, \mathbf{y}) are observed and yield

$$y = \underbrace{\mathbf{x}^T \mathbf{w}}_{f_{\mathbf{w}}(\mathbf{x})} + \epsilon, \quad \epsilon \sim \mathcal{N}(0, \sigma^2) \quad (16)$$





- ▶ $\hat{\mathbf{w}}$ is the estimator obtained based on the data.

$$\begin{aligned} \text{MSE} &= \mathbb{E}_{\mathbf{y}|\mathbf{x}, \mathbf{w}}[(y - \hat{y})^2] = \mathbb{E}[(f_{\mathbf{w}}(\mathbf{x}) + \epsilon - f_{\hat{\mathbf{w}}}(\mathbf{x}))^2] = \sigma^2 + \mathbb{E}[(f_{\mathbf{w}}(\mathbf{x}) - f_{\hat{\mathbf{w}}}(\mathbf{x}))^2] \\ &= \sigma^2 + \mathbb{E}[(f_{\mathbf{w}}(\mathbf{x}) - \mathbb{E}[f_{\hat{\mathbf{w}}}(\mathbf{x})] + \mathbb{E}[f_{\hat{\mathbf{w}}}(\mathbf{x})] - f_{\hat{\mathbf{w}}}(\mathbf{x}))^2] \\ &= \sigma^2 + \underbrace{\mathbb{E}[(f_{\mathbf{w}}(\mathbf{x}) - \mathbb{E}[f_{\hat{\mathbf{w}}}(\mathbf{x})])^2] + \mathbb{E}[2(f_{\mathbf{w}}(\mathbf{x}) - \mathbb{E}[f_{\hat{\mathbf{w}}}(\mathbf{x})])(\mathbb{E}[f_{\hat{\mathbf{w}}}(\mathbf{x})] - f_{\hat{\mathbf{w}}}(\mathbf{x}))]}_{=0} \\ &\quad + \mathbb{E}[(\mathbb{E}[f_{\hat{\mathbf{w}}}(\mathbf{x})] - f_{\hat{\mathbf{w}}}(\mathbf{x}))^2] \\ &= \underbrace{\sigma^2}_{\text{Irreducible Noise}} + \underbrace{\mathbb{E}[(f_{\mathbf{w}}(\mathbf{x}) - \mathbb{E}[f_{\hat{\mathbf{w}}}(\mathbf{x})])^2]}_{\text{Bias}^2(f_{\hat{\mathbf{w}}}(\mathbf{x})|f_{\mathbf{w}}(\mathbf{x}))} + \underbrace{\mathbb{E}[(f_{\hat{\mathbf{w}}}(\mathbf{x}) - \mathbb{E}[f_{\hat{\mathbf{w}}}(\mathbf{x})])^2]}_{\text{Variance}} \end{aligned} \quad (17)$$

Bias-Variance Trade-off to Avoid Overfitting and Underfitting



Bias-Variance Trade-off to Avoid Overfitting and Underfitting



Can We Learn Complicated Models from Sparse Data?

- ▶ Overfitting: Model complexity \gg data complexity
 - ▶ The number of model parameters is larger than that of data points
 - ▶ **Case 1:** The model is wrongly complicated \Rightarrow we need to simplify the model
 - ▶ **Case 2:** The model is with reasonable complexity but the data are insufficient \Rightarrow more common, and we need to introduce more side information.

Can We Learn Complicated Models from Sparse Data?

- ▶ Overfitting: Model complexity \gg data complexity
 - ▶ The number of model parameters is larger than that of data points
 - ▶ **Case 1:** The model is wrongly complicated \Rightarrow we need to simplify the model
 - ▶ **Case 2:** The model is with reasonable complexity but the data are insufficient \Rightarrow more common, and we need to introduce more side information.
- ▶ Underfitting: Model complexity \ll data complexity
 - ▶ The number of model parameters is smaller than that of data points
- ▶ To learn complicated models from sparse data, we need to impose side information on the model parameters (as **regularizers**)

Ridge Regression: MSE with L2 Regularization

- Ridge regression:

$$\min_{\mathbf{w}} \underbrace{\|\mathbf{y} - \mathbf{X}\mathbf{w}\|_2^2 + \lambda \|\mathbf{w}\|_2^2}_{L(\mathbf{w})} \quad (18)$$

Ridge Regression: MSE with L2 Regularization

- ▶ Ridge regression:

$$\min_{\mathbf{w}} \underbrace{\|\mathbf{y} - \mathbf{X}\mathbf{w}\|_2^2 + \lambda \|\mathbf{w}\|_2^2}_{L(\mathbf{w})} \quad (18)$$

- ▶ Consider the data fidelity and penalize the energy of parameters.

Ridge Regression: MSE with L2 Regularization

- ▶ Ridge regression:

$$\min_{\mathbf{w}} \underbrace{\|\mathbf{y} - \mathbf{X}\mathbf{w}\|_2^2 + \lambda \|\mathbf{w}\|_2^2}_{L(\mathbf{w})} \quad (18)$$

- ▶ Consider the data fidelity and penalize the energy of parameters.
- ▶ Closed form solution:

$$\frac{\partial L}{\partial \mathbf{w}} = 2\mathbf{X}^T(\mathbf{X}\mathbf{w} - \mathbf{y}) + 2\lambda\mathbf{w} = \mathbf{0}$$

Ridge Regression: MSE with L2 Regularization

- ▶ Ridge regression:

$$\min_{\mathbf{w}} \underbrace{\|\mathbf{y} - \mathbf{X}\mathbf{w}\|_2^2 + \lambda \|\mathbf{w}\|_2^2}_{L(\mathbf{w})} \quad (18)$$

- ▶ Consider the data fidelity and penalize the energy of parameters.
- ▶ Closed form solution:

$$\frac{\partial L}{\partial \mathbf{w}} = 2\mathbf{X}^T(\mathbf{X}\mathbf{w} - \mathbf{y}) + 2\lambda\mathbf{w} = \mathbf{0} \quad \Rightarrow \quad \mathbf{w} = (\mathbf{X}^T\mathbf{X} + \lambda\mathbf{I})^{-1}\mathbf{X}^T\mathbf{y}. \quad (19)$$

The name “ridge” refers to the shape along the diagonal of \mathbf{I} .

Ridge Regression: MSE with L2 Regularization

- ▶ Ridge regression:

$$\min_{\mathbf{w}} \underbrace{\|\mathbf{y} - \mathbf{X}\mathbf{w}\|_2^2 + \lambda \|\mathbf{w}\|_2^2}_{L(\mathbf{w})} \quad (18)$$

- ▶ Consider the data fidelity and penalize the energy of parameters.
- ▶ Closed form solution:

$$\frac{\partial L}{\partial \mathbf{w}} = 2\mathbf{X}^T(\mathbf{X}\mathbf{w} - \mathbf{y}) + 2\lambda\mathbf{w} = \mathbf{0} \quad \Rightarrow \quad \mathbf{w} = (\mathbf{X}^T\mathbf{X} + \lambda\mathbf{I})^{-1}\mathbf{X}^T\mathbf{y}. \quad (19)$$

The name “ridge” refers to the shape along the diagonal of \mathbf{I} .

- ▶ Stochastic gradient descent:

$$\mathbf{w}_{t+1} = \mathbf{w}_t - \tau \nabla_{\mathbf{w}_t} L \quad (20)$$

Ridge Regression: A Bayesian Viewpoint

- Data model:

$$y = \mathbf{x}^T \mathbf{w} + \epsilon, \quad \epsilon \sim \mathcal{N}(0, \sigma^2) \quad (21)$$

- Model prior:

$$\mathbf{w} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_D \gamma^2) \quad (22)$$

Ridge Regression: A Bayesian Viewpoint

- Data model:

$$y = \mathbf{x}^T \mathbf{w} + \epsilon, \quad \epsilon \sim \mathcal{N}(0, \sigma^2) \quad (21)$$

- Model prior:

$$\mathbf{w} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_D \gamma^2) \quad (22)$$

- Maximizing a posterior (MAP): (**Derive It**)

$$\max_{\mathbf{w}} p(\mathbf{w} | \mathbf{y}, \mathbf{X}) \propto \max_{\mathbf{w}} p(\mathbf{y} | \mathbf{X}, \mathbf{w}) \underbrace{p(\mathbf{w} | \mathbf{X})}_{p(\mathbf{w})}$$

Ridge Regression: A Bayesian Viewpoint

- Data model:

$$y = \mathbf{x}^T \mathbf{w} + \epsilon, \quad \epsilon \sim \mathcal{N}(0, \sigma^2) \quad (21)$$

- Model prior:

$$\mathbf{w} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_D \gamma^2) \quad (22)$$

- Maximizing a posterior (MAP): (**Derive It**)

$$\max_{\mathbf{w}} p(\mathbf{w} | \mathbf{y}, \mathbf{X}) \propto \max_{\mathbf{w}} p(\mathbf{y} | \mathbf{X}, \mathbf{w}) \underbrace{p(\mathbf{w} | \mathbf{X})}_{p(\mathbf{w})} \Rightarrow \max_{\mathbf{w}} \prod_n p(y_n | \mathbf{x}_n, \mathbf{w}) p(\mathbf{w})$$

$$\Rightarrow \min_{\mathbf{w}} - \sum_n \log p(y_n | \mathbf{x}_n, \mathbf{w}) - \log p(\mathbf{w})$$

Ridge Regression: A Bayesian Viewpoint

- Data model:

$$y = \mathbf{x}^T \mathbf{w} + \epsilon, \quad \epsilon \sim \mathcal{N}(0, \sigma^2) \quad (21)$$

- Model prior:

$$\mathbf{w} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_D \gamma^2) \quad (22)$$

- Maximizing a posterior (MAP): **(Derive It)**

$$\begin{aligned} \max_{\mathbf{w}} p(\mathbf{w} | \mathbf{y}, \mathbf{X}) &\propto \max_{\mathbf{w}} p(\mathbf{y} | \mathbf{X}, \mathbf{w}) \underbrace{p(\mathbf{w} | \mathbf{X})}_{p(\mathbf{w})} \Rightarrow \max_{\mathbf{w}} \prod_n p(y_n | \mathbf{x}_n, \mathbf{w}) p(\mathbf{w}) \\ &\Rightarrow \min_{\mathbf{w}} - \sum_n \log p(y_n | \mathbf{x}_n, \mathbf{w}) - \log p(\mathbf{w}) \Rightarrow \min_{\mathbf{w}} \frac{1}{2\sigma^2} \|\mathbf{y} - \mathbf{X}\mathbf{w}\|_2^2 + \frac{D}{2\gamma^2} \|\mathbf{w}\|_2^2 + C. \end{aligned} \quad (23)$$

Some Variants of Ridge Regression

Tikhonov regularization:

$$\min_{\mathbf{w}} \|\mathbf{y} - \mathbf{X}\mathbf{w}\|_2^2 + \lambda \|\mathbf{\Gamma}\mathbf{w}\|_2^2 \quad (24)$$

- ▶ $\mathbf{\Gamma}$: Tikhonov matrix
- ▶ Derive its closed form solution.

Some Variants of Ridge Regression

Tikhonov regularization:

$$\min_{\mathbf{w}} \|\mathbf{y} - \mathbf{X}\mathbf{w}\|_2^2 + \lambda \|\mathbf{\Gamma}\mathbf{w}\|_2^2 \quad (24)$$

- ▶ $\mathbf{\Gamma}$: Tikhonov matrix
- ▶ Derive its closed form solution.

Generalized Tikhonov regularization:

$$\min_{\mathbf{w}} \|\mathbf{y} - \mathbf{X}\mathbf{w}\|_P^2 + \lambda \|\mathbf{w} - \mathbf{w}_0\|_Q^2 \quad (25)$$

- ▶ $\|\mathbf{w}\|_A^2 = \mathbf{w}^T \mathbf{A} \mathbf{w}$
- ▶ Derive its closed form solution.

Some Variants of Ridge Regression

Tikhonov regularization:

$$\min_{\mathbf{w}} \|\mathbf{y} - \mathbf{X}\mathbf{w}\|_2^2 + \lambda \|\mathbf{\Gamma}\mathbf{w}\|_2^2 \quad (24)$$

- ▶ $\mathbf{\Gamma}$: Tikhonov matrix
- ▶ Derive its closed form solution.

Generalized Tikhonov regularization:

$$\min_{\mathbf{w}} \|\mathbf{y} - \mathbf{X}\mathbf{w}\|_{\mathbf{P}}^2 + \lambda \|\mathbf{w} - \mathbf{w}_0\|_{\mathbf{Q}}^2 \quad (25)$$

- ▶ $\|\mathbf{w}\|_{\mathbf{A}}^2 = \mathbf{w}^T \mathbf{A} \mathbf{w}$
- ▶ Derive its closed form solution.
- ▶ What if $\mathbf{P} = \Sigma_y^{-1}$, $\mathbf{Q} = \Sigma_w^{-1}$, and $\mathbf{w}_0 = \mathbb{E}[\mathbf{w}]$?

Lasso: MSE with L1 Regularization

Lasso (Least Absolute Shrinkage and Selection Operator)

$$\min_{\mathbf{w}} \|\mathbf{y} - \mathbf{X}\mathbf{w}\|_2^2 + \lambda \|\mathbf{w}\|_1 \quad (26)$$

- It is also called “Basis pursuit” in the field of signal processing.

Lasso: MSE with L1 Regularization

Lasso (Least Absolute Shrinkage and Selection Operator)

$$\min_{\mathbf{w}} \|\mathbf{y} - \mathbf{X}\mathbf{w}\|_2^2 + \lambda \|\mathbf{w}\|_1 \quad (26)$$

- ▶ It is also called “Basis pursuit” in the field of signal processing.
- ▶ A Bayesian Viewpoint of Lasso: $\mathbf{w} \sim \text{Laplace}(0, b\mathbf{I}_D)$, so that $p(\mathbf{w}) = \frac{1}{(2b)^D} \exp(-\frac{\|\mathbf{w}\|_1}{b})$. (Derive the MAP optimization problem)

Lasso: MSE with L1 Regularization

Lasso (Least Absolute Shrinkage and Selection Operator)

$$\min_{\mathbf{w}} \|\mathbf{y} - \mathbf{X}\mathbf{w}\|_2^2 + \lambda \|\mathbf{w}\|_1 \quad (26)$$

- ▶ It is also called “Basis pursuit” in the field of signal processing.
- ▶ A Bayesian Viewpoint of Lasso: $\mathbf{w} \sim \text{Laplace}(0, b\mathbf{I}_D)$, so that $p(\mathbf{w}) = \frac{1}{(2b)^D} \exp(-\frac{\|\mathbf{w}\|_1}{b})$. (Derive the MAP optimization problem)
- ▶ MAP:

$$\begin{aligned} \max_{\mathbf{w}} p(\mathbf{w}|\mathbf{y}, \mathbf{X}) &\propto \max_{\mathbf{w}} p(\mathbf{y}|\mathbf{X}, \mathbf{w}) \underbrace{p(\mathbf{w}|\mathbf{X})}_{p(\mathbf{w})} \Rightarrow \max_{\mathbf{w}} \prod_n p(y_n|\mathbf{x}_n, \mathbf{w}) p(\mathbf{w}) \\ &\Rightarrow \min_{\mathbf{w}} - \sum_n \log p(y_n|\mathbf{x}_n, \mathbf{w}) - \log p(\mathbf{w}) \Rightarrow \min_{\mathbf{w}} \frac{1}{2\sigma^2} \|\mathbf{y} - \mathbf{X}\mathbf{w}\|_2^2 + \frac{1}{b} \|\mathbf{w}\|_1 + C. \end{aligned} \quad (27)$$

Ridge Regression v.s. Lasso


Ridge regression:

- ▶ Penalize the energy of parameters.
- ▶ Strictly convex and easy to solve with linear convergence.

Lasso:

- ▶ Penalize the sparsity of parameters (benefits for model and feature selection).
- ▶ Convex but nonsmooth, relatively hard to solve with sublinear convergence.

Ridge Regression v.s. Lasso



Optimization Methods of Lasso Regression

$$\min_{\mathbf{w}} \|\mathbf{y} - \mathbf{X}\mathbf{w}\|_2^2 + \lambda \|\mathbf{w}\|_1 \quad (28)$$

Soft-thresholding: When $\mathbf{X} = [\mathbf{x}_1, \dots, \mathbf{x}_D] \in \mathbb{R}^{N \times D}$ are orthonormal ($\mathbf{X}^T \mathbf{X} = \mathbf{I}_D$):

- The solution of ordinary least squares (OLS) is

$$\hat{\mathbf{w}}^{(OLS)} = \arg \min_{\mathbf{w}} \|\mathbf{y} - \mathbf{X}\mathbf{w}\|_2^2 = (\mathbf{X}^T \mathbf{X}) \mathbf{X}^T \mathbf{y} = \mathbf{I}_D \mathbf{X}^T \mathbf{y} = \mathbf{X}^T \mathbf{y}.$$

Optimization Methods of Lasso Regression

$$\min_{\mathbf{w}} \|\mathbf{y} - \mathbf{X}\mathbf{w}\|_2^2 + \lambda \|\mathbf{w}\|_1 \quad (28)$$


Soft-thresholding: When $\mathbf{X} = [\mathbf{x}_1, \dots, \mathbf{x}_D] \in \mathbb{R}^{N \times D}$ are orthonormal ($\mathbf{X}^T \mathbf{X} = \mathbf{I}_D$):

- The solution of ordinary least squares (OLS) is

$$\hat{\mathbf{w}}^{(OLS)} = \arg \min_{\mathbf{w}} \|\mathbf{y} - \mathbf{X}\mathbf{w}\|_2^2 = (\mathbf{X}^T \mathbf{X}) \mathbf{X}^T \mathbf{y} = \mathbf{I}_D \mathbf{X}^T \mathbf{y} = \mathbf{X}^T \mathbf{y}. \quad (29)$$

- The solution of lasso also has a closed form:

$$\hat{w}_d = S_\lambda(\hat{w}_d^{(OLS)}) = \text{sign}(\hat{w}_d^{(OLS)}) \max\{0, |\hat{w}_d^{(OLS)}| - \lambda\}, \quad \forall d = 1, \dots, D. \quad (30)$$



Optimization Methods of Lasso Regression

$$\min_{\mathbf{w}} \|\mathbf{y} - \mathbf{X}\mathbf{w}\|_2^2 + \lambda \|\mathbf{w}\|_1, \quad \text{where } \mathbf{X} = [\mathbf{x}_1, \dots, \mathbf{x}_D] \quad (31)$$

Iterative soft-thresholding for general situations: Although $\mathbf{X}^T \mathbf{X} \neq \mathbf{I}_D$, we can **construct orthonormal vectors column-wisely and update parameters iteratively.**

- In the t -th iteration, for $d = 1, \dots, D$:

$$\begin{aligned} \hat{w}_d^{(t+1)} &= \arg \min_w \left\| \mathbf{y} - \underbrace{\sum_{i \neq d} \mathbf{x}_i w_i^{(t)}}_{\mathbf{X}_{-d} \mathbf{w}_{-d}^{(t)}} - \mathbf{x}_d w_d \right\|_2^2 + \lambda |w| \\ &= \arg \min_w \left\| \frac{1}{\|\mathbf{x}_d\|_2} (\mathbf{y} - \mathbf{X}_{-d} \mathbf{w}_{-d}^{(t)}) - w \underbrace{\frac{\mathbf{x}_d}{\|\mathbf{x}_d\|_2}}_{\text{orthonormal}} \right\|_2^2 + \frac{\lambda}{\|\mathbf{x}_d\|_2^2} |w| \quad (32) \\ &= S_{\frac{\lambda}{\|\mathbf{x}_d\|_2^2}} \left(\frac{\mathbf{x}_d^T (\mathbf{y} - \mathbf{X}_{-d} \mathbf{w}_{-d}^{(t)})}{\|\mathbf{x}_d\|_2^2} \right) \end{aligned}$$

Other Methods for Sparse Model Parameters

Lasso

- ▶ ADMM (Alternating Direction Method of Multiplier)
- ▶ LARS (Least Angle Regression) ...

Other Methods for Sparse Model Parameters

Lasso

- ▶ ADMM (Alternating Direction Method of Multiplier)
- ▶ LARS (Least Angle Regression) ...

Stronger sparsity: **L0 Regularization and Hard Thresholding**

$$\min_{\mathbf{w}} \|\mathbf{y} - \mathbf{X}\mathbf{w}\|_2^2 + \lambda \|\mathbf{w}\|_0 \quad (33)$$

Other Methods for Sparse Model Parameters

Lasso

- ▶ ADMM (Alternating Direction Method of Multiplier)
- ▶ LARS (Least Angle Regression) ...

Stronger sparsity: **Lo Regularization and Hard Thresholding**

$$\min_{\mathbf{w}} \|\mathbf{y} - \mathbf{X}\mathbf{w}\|_2^2 + \lambda \|\mathbf{w}\|_0 \quad (33)$$

Weaker sparsity: **Elastic net Regularization**

$$\min_{\mathbf{w}} \|\mathbf{y} - \mathbf{X}\mathbf{w}\|_2^2 + \lambda_1 \|\mathbf{w}\|_1 + \lambda_2 \|\mathbf{w}\|_2^2 \quad (34)$$

Other Methods for Sparse Model Parameters

Lasso

- ▶ ADMM (Alternating Direction Method of Multiplier)
- ▶ LARS (Least Angle Regression) ...

Stronger sparsity: **Lo Regularization and Hard Thresholding**

$$\min_{\mathbf{w}} \|\mathbf{y} - \mathbf{X}\mathbf{w}\|_2^2 + \lambda \|\mathbf{w}\|_0 \quad (33)$$

Weaker sparsity: **Elastic net Regularization**

$$\min_{\mathbf{w}} \|\mathbf{y} - \mathbf{X}\mathbf{w}\|_2^2 + \lambda_1 \|\mathbf{w}\|_1 + \lambda_2 \|\mathbf{w}\|_2^2 \quad (34)$$

How to interpret it from a Bayesian viewpoint?

Extensions: How to Deal With Outliers?

MAE

$$\min_{\mathbf{w}} \|\mathbf{y} - \mathbf{X}\mathbf{w}\|_1 \quad (35)$$

Extensions: How to Deal With Outliers?

MAE

$$\min_{\mathbf{w}} \|\mathbf{y} - \mathbf{X}\mathbf{w}\|_1 \quad (35)$$

IRLS (Iteratively Reweighted Least Squares)

$$\begin{aligned} \mathbf{w}^{(t+1)} &= \arg \min_{\mathbf{w}} \sum_{n=1}^N \alpha_n(\mathbf{w}^{(t)}) |y_n - \mathbf{x}_n^T \mathbf{w}|^2 \\ &= \arg \min_{\mathbf{w}} \left\| \underbrace{\text{diag}^{\frac{1}{2}}(\boldsymbol{\alpha}(\mathbf{w}^{(t)}))}_{\mathbf{A}^{(t)\frac{1}{2}}} (\mathbf{y} - \mathbf{X}\mathbf{w}) \right\|_2^2 \end{aligned}$$

Extensions: How to Deal With Outliers?

MAE

$$\min_{\mathbf{w}} \|\mathbf{y} - \mathbf{X}\mathbf{w}\|_1 \quad (35)$$

IRLS (Iteratively Reweighted Least Squares)

$$\begin{aligned} \mathbf{w}^{(t+1)} &= \arg \min_{\mathbf{w}} \sum_{n=1}^N \alpha_n(\mathbf{w}^{(t)}) |y_n - \mathbf{x}_n^T \mathbf{w}|^2 \\ &= \arg \min_{\mathbf{w}} \left\| \underbrace{\text{diag}^{\frac{1}{2}}(\boldsymbol{\alpha}(\mathbf{w}^{(t)}))}_{\mathbf{A}^{(t)\frac{1}{2}}} (\mathbf{y} - \mathbf{X}\mathbf{w}) \right\|_2^2 = (\mathbf{X}^T \mathbf{A}^{(t)} \mathbf{X})^{-1} \mathbf{X}^T \mathbf{A}^{(t)} \mathbf{y} \end{aligned}$$

Extensions: How to Deal With Outliers?

MAE

$$\min_{\mathbf{w}} \|\mathbf{y} - \mathbf{X}\mathbf{w}\|_1 \quad (35)$$

IRLS (Iteratively Reweighted Least Squares)

$$\begin{aligned} \mathbf{w}^{(t+1)} &= \arg \min_{\mathbf{w}} \sum_{n=1}^N \alpha_n(\mathbf{w}^{(t)}) |y_n - \mathbf{x}_n^T \mathbf{w}|^2 \\ &= \arg \min_{\mathbf{w}} \left\| \underbrace{\text{diag}^{\frac{1}{2}}(\boldsymbol{\alpha}(\mathbf{w}^{(t)}))}_{\mathbf{A}^{(t)\frac{1}{2}}} (\mathbf{y} - \mathbf{X}\mathbf{w}) \right\|_2^2 = (\mathbf{X}^T \mathbf{A}^{(t)} \mathbf{X})^{-1} \mathbf{X}^T \mathbf{A}^{(t)} \mathbf{y} \\ \alpha_n^{(0)} &= 1, \quad \alpha_n^{(t)} = |y_n - \mathbf{x}_n^T \mathbf{w}^{(t)}|^{-1} \end{aligned} \quad (36)$$

Extensions: How to Deal With Outliers?

MAE

$$\min_{\mathbf{w}} \|\mathbf{y} - \mathbf{X}\mathbf{w}\|_1 \quad (35)$$

IRLS (Iteratively Reweighted Least Squares)

$$\begin{aligned} \mathbf{w}^{(t+1)} &= \arg \min_{\mathbf{w}} \sum_{n=1}^N \alpha_n(\mathbf{w}^{(t)}) |y_n - \mathbf{x}_n^T \mathbf{w}|^2 \\ &= \arg \min_{\mathbf{w}} \left\| \underbrace{\text{diag}^{\frac{1}{2}}(\boldsymbol{\alpha}(\mathbf{w}^{(t)}))}_{\mathbf{A}^{(t)\frac{1}{2}}} (\mathbf{y} - \mathbf{X}\mathbf{w}) \right\|_2^2 = (\mathbf{X}^T \mathbf{A}^{(t)} \mathbf{X})^{-1} \mathbf{X}^T \mathbf{A}^{(t)} \mathbf{y} \end{aligned} \quad (36)$$

$$\alpha_n^{(0)} = 1, \quad \alpha_n^{(t)} = |y_n - \mathbf{x}_n^T \mathbf{w}^{(t)}|^{-1}$$

- ▶ It works for p -norm with $p \leq 2$, i.e., $\alpha_n^{(t)} = |y_n - \mathbf{x}_n^T \mathbf{w}^{(t)}|^{p-2}$
- ▶ It works as the MLE of GLM, i.e., $y = f_{\mathbf{w}}(\mathbf{x})$.

In Summary

- ▶ Introduce generalized linear regression problem
- ▶ Theoretical analysis of linear regression models and some key concepts of statistical machine learning (bias and variance)
- ▶ Typical regularization methods and their Bayesian interpretability

Next...

- ▶ Non-linear regression
- ▶ Duality and kernelization
- ▶ Gaussian process