

axionbloch: an Open-Source Python Package for Simulating Axion-Induced Spin Dynamics

Yuzhe Zhang*

Johannes Gutenberg-Universität Mainz, 55128 Mainz, Germany

Helmholtz Institute Mainz, 55099 Mainz, Germany and

GSI Helmholtzzentrum für Schwerionenforschung GmbH, 64291 Darmstadt, Germany

And more people?

The interaction of ultralight bosonic dark matter with spins can be interpreted as a pseudo-magnetic field acting on normal matter. We model such interaction along with the usual magnetic interactions using spin-evolution (Bloch) equations. We present **axionbloch**, an open-source Python package for simulating spin dynamics induced by both usual and exotic interactions. The numerical simulations serve as a tool for deriving axion signal signatures, which are crucial for designing experimental searches and data analysis. Simulations are calibrated against theoretical expectation, ensuring the accuracy of the simulated signals. **axionbloch** is available at <https://github.com/Yuzhe98/AxionBloch>, allowing researchers to simulate axion pseudomagnetic signals under specific configurations of the axion models and experimental setups. The package is documented at <http://axionbloch.readthedocs.io/> and includes example scripts for application.

I. INTRODUCTION

The axion was originally proposed to resolve the strong CP problem in quantum chromodynamics [1–4]. Here C denotes charge conjugation and P denotes parity. Later, more general axionlike particles (ALPs) were proposed in various extensions of the Standard Model associated with spontaneously broken global symmetries [5–8]. Concerning the cosmological implications, axions and ALPs are also well-motivated dark matter candidates [9–11]. In this paper, we refer to axions and ALPs as “axions” for simplicity.

Axions may possess couplings to ordinary Standard Model particles[?], including photons, gluons and fermions. Crucially, axion field gradients (∇a) may couple to fermionic spins through a pseudoscalar interaction. The Hamiltonian for this interaction can be expressed as

$$\mathcal{H}_{\text{int}} = -g_{\text{aNN}} \nabla a \cdot \mathbf{S}, \quad (1)$$

where g_{aNN} is the axion-spin coupling strength, and $\mathbf{S}$ is the spin operator. The axion field gradient can be regarded as a pseudomagnetic field that mimics the behavior of a real magnetic field, where the Hamiltonian can be written as

$$\mathcal{H} = -\hbar\gamma \mathbf{B} \cdot \mathbf{S}, \quad (2)$$

where γ is the gyromagnetic ratio of the spin, and $\mathbf{B}$ is the magnetic field. The pseudomagnetic field induces spin dynamics, providing a pathway for detection in precision measurement experiments such as nuclear magnetic resonance (NMR) and comagnetometer-based searches [? ?]

[17–19]. There have been dedicated studies of the axion-induced spin dynamics [20–22], providing axion signal signatures and experimental sensitivity to axion fields. However, to our knowledge, there have not been studies taking advantage of numerical methods to verify the derived axion signal signatures. On the other hand, as more axion models and experiments are being proposed, such as cosmic axion background [23], fine-grained axion [24], and quadratically coupled axion [25], a tool is needed to systematically simulate the axion-induced spin dynamics under various physical conditions and axion field configurations.

In this work, we present **axionbloch**, an open-source package designed to simulate spin dynamics induced by axion fields. The interface is implemented in Python, a programming language known for its simplicity and readability, while the underlying numerical integration is implemented in C++ for efficiency. The package provides a flexible framework for modeling axion fields and experimental configurations. By numerically solving the relevant spin evolution equations, **axionbloch** enables the prediction and analysis of experimentally observable signals.

II. CAPABILITIES

In this section, we describe the main components of **axionbloch**. We start with the units, physical quantities, and constants (Sec. II A). Then, we continue with the physics and the algorithm for the numerical simulation (Sec. II B), after which the calibration for the simulation follows (??). In the end, we provide an example simulation of axion-induced signals (??).

* Contact author: yuhzhang@uni-mainz.de

```

Define a quantity from scalar and unit
>>> from astropy import units as u
>>> B = 1.0 * u.gauss
>>> B
<Quantity 1. G>
>>> B.si
<Quantity 0.0001 T>

Import constants and use them with scalars
>>> from axionbloch.constants import gamma_p
>>> import numpy as np
find the 90 degree pulse duration
>>> t90 = np.pi / 2 / (gamma_p * B)
>>> t90.to(u.microsecond)
<Quantity 58.71648793 us>

Operation on an array of quantities with numpy
>>> tStamps = np.array([0, 1 / 3, 1]) * t90
>>> phases = gamma_p * B * tStamps * u.radian
>>> np.sin(phases)
<Quantity [0. , 0.5, 1. ]>

```

FIG. 1. Examples of operations of physical quantities using `astropy.units` and `axionbloch.constants`.

A. Units, Physical Quantities, and Constants

We take advantage of the packages `astropy`¹ [?] to deal with units and physical quantities. Using `astropy.units`, we can create physical quantities by attaching units to Python scalars or NumPy² [?] arrays. Quantities can be converted between dimensionally equivalent units using the `to()` method. In `astropy`, unit definitions are included in both the International System of Units (SI) and the Centimeter-Gram-Second (CGS) systems. Values in SI or CGS units can be found in the quantity's properties `.si` and `.cgs`. A code example can be found in Fig. 1.

In `axionbloch.constants`, we defined a few commonly-used constants in axion and NMR research, like gyromagnetic ratios. The constants are instances of `astropy.units.Quantity`; therefore, all operations on physical quantities are supported for the constants.

B. Algorithm

1. Spin Dynamics

Considering the current experiments looking for axion-induced spin dynamics, `axionbloch` focuses on spin-1/2 magnetic resonance. Different spins may be addressed in future updates. We use magnetization as the observable to characterize the spin ensemble. In the presence of

magnetic fields, including both bias fields applied in experiments and the axion-induced pseudomagnetic fields, the magnetization $\mathbf{M}$ evolves as

$$\frac{dM_x}{dt} = \gamma (\mathbf{M} \times \mathbf{B}_{\text{eff}})_x - \frac{M_x}{T_2}, \quad (3)$$

$$\frac{dM_y}{dt} = \gamma (\mathbf{M} \times \mathbf{B}_{\text{eff}})_y - \frac{M_y}{T_2}, \quad (4)$$

$$\frac{dM_z}{dt} = \gamma (\mathbf{M} \times \mathbf{B}_{\text{eff}})_z - \frac{M_z - M_0}{T_1}, \quad (5)$$

where γ is the gyromagnetic ratio, $\mathbf{B}_{\text{eff}}$ is the total effective field, T_1 and T_2 are relaxation times. The bias field is usually much stronger than the pseudomagnetic field. Therefore, we adopt the rotating coordinate frame (RCF) to remove the Larmor precession due to the bias field, allowing efficient simulation of the slower axion-induced dynamics of interest.

The transverse relaxation time, characterized by time constant T_2^* , includes the T_2 relaxation and the dephasing due to the inhomogeneity of the bias field. Such inhomogeneity of the bias field is simulated by sampling the spread of the bias fields in the `Magnet` instance, and solving the Bloch equations for individual bias fields.

2. Numerical Integration

To derive the time evolution of the magnetization, `axionbloch` numerically integrates the Bloch equations using the fourth-order Runge-Kutta (RK4) method. Given a time step Δt , effective magnetic field, and initial magnetization $\mathbf{M}_0$, the magnetization at the $(n+1)$ -th step can be found via

$$\mathbf{M}^{n+1} = \mathbf{M}^n + \frac{\Delta t}{6} (\mathbf{k}_1 + 2\mathbf{k}_2 + 2\mathbf{k}_3 + \mathbf{k}_4). \quad (6)$$

Here

$$\mathbf{k}_1 = \left. \frac{d\mathbf{M}}{dt} \right|_{\mathbf{M}=\mathbf{M}^n}, \quad (7)$$

$$\mathbf{k}_2 = \left. \frac{d\mathbf{M}}{dt} \right|_{\mathbf{M}=\mathbf{M}^n + \frac{1}{2}\mathbf{k}_1\Delta t}, \quad (8)$$

$$\mathbf{k}_3 = \left. \frac{d\mathbf{M}}{dt} \right|_{\mathbf{M}=\mathbf{M}^n + \frac{1}{2}\mathbf{k}_2\Delta t}, \quad (9)$$

$$\mathbf{k}_4 = \left. \frac{d\mathbf{M}}{dt} \right|_{\mathbf{M}=\mathbf{M}^n + \mathbf{k}_3\Delta t}, \quad (10)$$

where $d\mathbf{M}/dt$ can be found using the Bloch equations [Eqs. (3–5)]. This fourth-order method yields a local truncation error of $\mathcal{O}(\Delta t^5)$, while the accumulated error over $N = 1/\Delta t$ steps is of order $\mathcal{O}(\Delta t^4)$. The time step Δt is usually set to be at least one order of magnitude smaller than the characteristic timescales of the system, such as relaxation times.

¹ <https://astropy.org/>

² <https://numpy.org/>

3. Implementation

Using object-oriented programming, we implement the algorithm in a modular way to separate experimental configuration, axion field modeling, and numerical integration, allowing for flexible configuration of simulations and axion field models. A diagram of the architecture of `axionbloch` is shown in ??.

On the Python side, classes `Sample`, `Magnet`, and `MagField` encapsulate the relevant experimental parameters, including sample relaxation times, magnetic fields strength and homogeneity. The axion field models are implemented by dedicated classes, such as `MilkyWayAxionHalo`, which carry the properties of the axion fields. Together with the method `MagField.setAxionFields()`, the pseudomagnetic fields can be generated. Different axion models can be incorporated as separate classes without modifying the simulation framework. Simulation is handled within the `axionbloch.SimuTools` module. In particular, the classes `Simulation` and `Simulations` manage the interaction between experimental configurations and axion field models, organizing the parameters and execution of single or multiple simulation runs.

The RK4 algorithm is implemented in a dedicated C++ library. The application programming interface (API) for this library is `blochSimulation`. This C++ backend is interfaced with Python using `pybind11`,³ enabling cooperation between high-level Python workflow management and low-level efficient numerical computation.

C. Calibration

Accurate calibration is essential to ensure that the simulated signals reliably reflect the underlying physical system and can be compared with experimental observables. In particular, calibration examines the correspondence between the NMR signal and model parameters, including magnetic fields, nuclear spin gyromagnetic ratio γ and relaxation times (T_1 , T_2 and T_2^*). It allows one to verify the correctness and stability of the numerical implementation. In this section, examples of calibration simulations are presented along with theoretical predictions, demonstrating the methodology of calibration. A quantitative analysis is included by the end of the section. For the examples in the section, an ethanol sample and an axial bias field $\mathbf{B}_0 = B_0 \hat{\mathbf{e}}_z$ are used.

1. Pulsed-NMR: Free Decay and Spin Echo

In the pulsed-NMR calibration, we apply one 90° pulse and one 180° magnetic-field pulse in the simulation. The

magnetization is expected to[? ?]:

- oscillate at the Larmor frequency;
- be tipped by 90° to the x-y plane by the 90° pulse;
- free decay with relaxation time T_2^* in the absence of pulses;
- refocus (echo) after 180° pulses;
- show decaying amplitude of echoes with relaxation time T_2 .

The behaviors described above can be found in the calibration example illustrated in ?? (a), indicating a rational simulation. With such benchmarks, magnetic pulses, Larmor precession, and transverse relaxations (T_2 and T_2^*) can be calibrated.

2. Continuous-Wave NMR

In the continuous-wave (CW) NMR calibration, an oscillating magnetic field is applied. When the magnetic field strength B is weak ($\gamma B T_2^* \ll 1$) and on resonance, the transverse magnetization grows with $\gamma B T_2^* (1 - e^{-t/T_2^*})$ [?]. One such example is illustrated in ?? (b). The simulation goes as expected. With such benchmarks, the Larmor precession and T_2^* relaxation can be calibrated.

3. Hyperpolarized sample

To achieve better sensitivity to the axion field, hyperpolarization is favored in experiments. Compared to thermal polarization at room temperature, hyperpolarization techniques can boost the polarization by orders of magnitude to the order of unity. The `axionbloch` package handles the polarization by the `Sample` class. We use a 1 %-polarization ethanol sample as an example demonstrating the evolution of hyperpolarization due to the T_1 relaxation. At first ($t \ll T_1$), the polarization exponentially decays with time constant T_1 , until it reaches equilibrium after many T_1 . As can be seen in ??, the T_1 relaxation followed the expectation. More of such simulation can be applied in calibrating T_1 relaxation.

4. Quantitative evaluation

Accuracy of simulations is evaluated by the discrepancy between theoretically-expected magnetization and simulation results, characterized by

$$\chi^2 = \frac{\sum |M_{\text{expected}} - M_{\text{simu}}|^2}{\sum |M_{\text{simu}}|^2}. \quad (11)$$

Here M_{expected} and M_{simu} are magnetization expected and simulated, respectively. $\sum$ stands for the summary

³ <http://pybind11.readthedocs.io/>




FIG. 2. Architecture of `axionbloch`. Each `Simulation` instance imports parameters from `Magnet`, `Sample`, and `MagField` instances. For axion simulations, axion-model objects are required both for `Simulation` to configure the simulation and for `MagField` to generate the pseudomagnetic field. `Simulations` manages one or multiple `Simulation` instances. Numerical integration is performed via the `blochSimulation` API, which provides Python access to the underlying C++ implementation.




FIG. 3. Simulated magnetic fields in x and y directions (upper row) and induced NMR signals (lower row). The magnetizations plotted are normalized by equilibrium magnetization. (a) After 90° and 180° magnetic-field pulses in x-y plane, free decay (relaxation time T_2^*) and echo signals (relaxation time T_2) are generated. (b) Excited by a weak CW magnetic field, the NMR signals increase with $\gamma B T_2^* (1 - e^{-t/T_2^*})$, where $B = \sqrt{B_x^2 + B_y^2}$ is the magnitude of the excitation.

over integration steps. Well-calibrated simulation should give $\chi^2 \ll 1$. Calibration scripts are stored in the test script set of `axionbloch`. In each calibration, parameters such as Larmor frequency and relaxation times are varied over orders of magnitude to cover all possible cases in simulation.

D. Axion Simulation

Similar to the calibration examples, we simulate the axion-induced NMR signals by solving the Bloch equations with the axion-induced pseudomagnetic field as part of the effective magnetic field. We present an example of a commonly considered model – axion consti-

tuting the Milky Way galactic dark matter halo[? ?], implemented in the class `MilkyWayAxionHalo`. In this model, the axion field is treated as an oscillating field with a frequency determined by the axion mass, and a linewidth (characterizing the stochasticity) determined by the velocity dispersion of the virialized dark matter halo. The sample is chosen to be ^{129}Xe , polarized to 50 % initially, and the bias field is applied along the z direction. The details of the simulation configuration can be found in the code example in ???. We refer readers to ??? which illustrates the relationship between the objects in the code example. The simulated pseudomagnetic fields and the amplitude of the induced transverse magnetization are shown in ???, in which we can find the signal increasing with time at the beginning of the simulation

TABLE I. Summary of calibration simulations. Varied parameters were swept over orders of magnitude to cover all simulation regimes relevant to CASPER-class experiments. χ^2 is defined in ??; $\chi^2 \ll 1$ indicates a well-calibrated simulation.

Calibration	Parameters calibrated	Varied parameters	χ^2	Error analysis
Pulsed NMR: free decay & spin echo (??)	Larmor frequency, RF flip angle, T_2^* , T_2	ω_0 , T_2^* , T_2 , pulse amplitude B_1	...	Dominated by integration step size relative to the Larmor period; accuracy degrades when simulation rate $\lesssim \omega_0/2\pi$
CW-NMR (??)	Larmor frequency, T_2^* , on-resonance response amplitude	ω_0 , T_2^* , driving field B_1	...	Requires weak-field regime $\gamma B_1 T_2^* \ll 1$; error scales with integration step size
Hyperpolarized T_1 relaxation (??)	T_1 relaxation timescale, initial polarization	T_1 , initial polarization M_{init}	...	Discretization error scales as $(\Delta t/T_1)^2$; accurate when $\Delta t \ll T_1$




FIG. 4. Simulated T_1 relaxation of a hyperpolarized sample in a static bias field. The axial magnetization M_z is normalized by the equilibrium magnetization M_0 .

due to axion field gradient, and then decaying due to the T_1 relaxation. The efficiency of the simulation is limited by time consumption for the numerical integration of the Bloch equations. The total number of integration steps can be found by

$$N = \text{Simulation.rate} \times \text{Simulation.duration} \times \text{MagField.numFields} \times \text{Magnet.numPt}, \quad (12)$$

where `Simulation.rate` is the rate of the simulation, `Simulation.duration` is the duration of the simulation, `MagField.numFields` is the number of random pseudo-magnetic fields generated to account for the stochasticity of the axion field, and `Magnet.numPt` is the number of bias fields sampled to account for the inhomogeneity of the bias field. For the example in ??, the total number of integration steps is 4.064×10^9 , and the runtime is 10s on a personal computer (CPU: Intel[®] Core[™] Ultra 7 155H; base speed: 1.40 GHz). Such moderate runtime allows users to explore different parameter regimes with many simulations.

III. DEVELOPMENT

Development of `axionbloch` is managed using `git`,⁴ with the code hosted on a public GitHub repository.⁵ Changes to the main codebase follow a *pull request* workflow, allowing contributions to be reviewed before being merged. Users can participate by forking the repository, making modifications in their fork, and submitting pull requests for review by the developers. As `axionbloch` is intended as a collaborative resource, user contributions and suggestions are encouraged and appreciated.

`axionbloch` includes a suite of automated tests to ensure correctness and reliability. The tests are implemented using the `pytest` framework,⁶ which allows for calibration of physical processes and testing of functions, classes, and modules. The test suite covers core functionalities such as:

- initialization and manipulation of classes;
- numerical integration of the Bloch equations;
- magnetic fields and relaxations;
- handling of each axion field configurations.

Tests can be executed locally. In the future, we may add tools for online testing considering the demand of testing. Regular execution of the test suite ensures that new changes do not introduce regressions, maintaining the stability and correctness of the package.

Documentation for `axionbloch` is available on Read the Docs.⁷ It is built using `Sphinx`⁸ and provides detailed explanations of the API, along with instructions for installation, configuration, and usage.

⁴ <http://git-scm.com>

⁵ <https://github.com/Yuzhe98/AxionBloch>

⁶ <https://docs.pytest.org/>

⁷ <http://axionbloch.readthedocs.io>

⁸ <http://www.sphinx-doc.org>

```

# numpy, matplotlib, astropy dependency
from axionbloch.dependency import *
# Gyromagnetic ratio and magnetic dipole moment of Xe-129
from axionbloch.constants import gamma_Xe129, mu_Xe129
# classes for simulations
from axionbloch.SimuTools import MagField, Simulations
from axionbloch.SimuTypes import SimuParams
from axionbloch.Sample import Sample
from axionbloch.Apparatus import Magnet
from axionbloch.MilkyWayAxionHalo import MilkyWayAxionHalo
# Define the Xe-129 sample with gyromagnetic ratio,
# mass density, molar mass, number of spins per molecule,
# relaxation times, volume, magnetic dipole moment,
# temperature, and polarization.
sample = Sample(
    name="Liquid Xe-129",
    gamma=gamma_Xe129,
    massDensity=3.1 * unit.g * unit.cm ** (-3),
    molarMass=131.29 * unit.g / unit.mol,
    numofSpinsPerMolecule=1 * unit.one,
    T2=10 * unit.minute,
    T1=15 * unit.minute,
    vol=1 * unit.cm**3,
    mu=mu_Xe129, # magnetic dipole moment
    temp=163 * unit.K,
    pol=1 * unit.percent,
    verbose=False,
)
# Define the axion field with axion Compton frequency
# and axion-nucleon coupling strength
axion = MilkyWayAxionHalo(
    name="Milky Way Axion Halo",
    nu_a=1 * unit.kHz,
    g_aNN=1.0e-9 * unit.GeV ** (-1),
    verbose=False,
)
# Set the bias field strength, direction, and homogeneity
magnet = Magnet(
    B0=axion.nu_a_eff / (sample.gamma / (2 * PI)),
    FWHM=2 * ppm,
)
# rms amplitude of pseudomagnetic field
B_a_rms = (axion.getRabiFreq() / (sample.gamma / (2 * PI)))**0.5
# Bundle all inputs into one dictionary
params: SimuParams = {
    "key_info": {"nu_a": axion.nu_a},
    "axion": axion,
    "sample": sample,
    "magnet": magnet,
    "excField": MagField(),
    "B_a_rms": B_a_rms,
    # Number of random field realizations.
    "numFields": 1000,
    "rand_seed": 10, # random seed
    # amplitude, polar and azimuthal angle
    # of the initial magnetization
    "init_M": None, # use None when setting M automatically
    "init_M_theta": 0 * unit.rad,
    "init_M_phi": 0 * unit.rad,
    # sampling rate and duration of the time series
    "rate": 1 * unit.Hz,
    "duration": 4000 * unit.s,
}
# Create and execute the simulation job collection
simulations = Simulations(all_params=[params])
# run the simulation
simulations.run(verbose=True)
# plot results
for i, item in enumerate(simulations.pool):
    item.simu.displayTrjries()
# Save to .pkl file for later analysis
simulations.saveToPkl(dir="path_to_save")

```

FIG. 5. Example code of an axion simulation.

As we are having numerous new axion field models and experiments emerging, we are planning to extend the functionality, so that axionbloch can be a tool for updating our knowledge of the induced spin dynamics.

IV. CONCLUSIONS

We have presented axionbloch, a package providing a framework to simulate spin dynamics induced by magnetic fields or axion fields under a variety of physical scenarios. The package adopts SI units so that the inputs and outputs are directly comparable to experimental parameters and observables. Numerical integration of the Bloch equations is performed using a fourth-order Runge-Kutta method in the rotating coordinate frame, enabling stable and efficient simulation. Comprehensive calibrations have been done to ensure the accuracy of the simulations. An example simulation of axion-induced signals from a local virialized axion dark matter halo is presented for demonstration of package's capabilities.

By providing an open-source, well-documented, and tested tool, axionbloch aims to facilitate the study of axion-induced spin dynamics in precision measurement experiments and to serve as a platform for further method development and incorporation of new axion models.

We note that this package can be of interest to the NMR community for its applications in, for example, educational purposes. For example, students can specify pulse sequences for the simulation and see the visualized simulation results.

DATA AVAILABILITY

The data and code that generate this manuscript are available at <https://github.com/Yuzhe98/AxionBloch-paper>.

ACKNOWLEDGMENTS

The authors acknowledge helpful discussions with [names]. We are grateful to the communities supporting the open-source software and tools: git, NumPy, SciPy⁹ [?], matplotlib¹⁰, astropy, pybind11, IPython¹¹ [?], Jupyter¹² [?], pytest, Sphinx, GitHub¹³ and Read the Docs¹⁴.

⁹ <https://docs.scipy.org/>

¹⁰ <https://matplotlib.org/>

¹¹ <https://ipython.org/>

¹² <https://jupyter.org/>

¹³ <https://github.com/>

¹⁴ <https://about.readthedocs.com/>




FIG. 6. Simulations of local virialized axion dark matter halo. From top to bottom: pseudomagnetic fields in the transverse plane (B_x and B_y) and the transverse magnetization induced (M_{xy}). Left: one single simulation. Right: 1000 simulations to account for the stochasticity. The solid lines indicate the value (left column) or mean value (right column) of B_x , B_y , and M_{xy} , while the gray shadows indicate $\pm\sigma$ (standard deviation) range. The sample for the simulations was assumed to be polarized to 50 % initially.

This work has been supported by the Cluster of Excellence “Precision Physics, Fundamental Interactions, and Structure of Matter” (PRISMA++ EXC 2118/2) funded by the German Research Foundation (DFG) within the German Excellence Strategy (Project ID 390831469).

Yuzhe: Check acknowledgments!

AUTHOR CONTRIBUTION STATEMENT

Yuzhe: I propose we have such a section.

Yuzhe Zhang: Conceptualization, Software, Visualization, Writing – original draft and Writing – review & editing

Dmitry Budker: Funding acquisition, Supervision, Writing – review & editing

Author contribution roles and definitions

Conceptualization: Ideas; formulation or evolution of overarching research goals and aims.

Data curation: Management activities to annotate (produce metadata), scrub data, and maintain research data (including software code, where necessary for interpreting the data itself) for initial use and later re-use.

Formal analysis: Application of statistical, mathematical, computational, or other formal techniques to analyze or synthesize study data.

Funding acquisition: Acquisition of the financial support for the project leading to this publication.

Investigation: Conducting a research and investigation process, specifically performing the experiments or data/evidence collection.

Methodology: Development or design of methodology; creation of models.

Project administration: Management and coordination responsibility for the research activity planning and execution.

Resources: Provision of study materials, reagents, materials, laboratory samples, instrumentation, computing resources, or other analysis tools.

Software: Programming, software development; designing computer programs; implementation of the computer code and supporting algorithms; testing of existing code components.

Supervision: Oversight and leadership responsibility for the research activity planning and execution, including mentorship external to the core team.

Validation: Verification, whether as a part of the activity or separate, of the overall replication or reproducibility of results, experiments, and other research outputs.

Visualization: Preparation, creation, and/or presentation of the published work, specifically visualization or data presentation.

Writing – original draft: Preparation, creation, and/or presentation of the published work, specifically writing the initial draft (including substantive translation).

Writing – review & editing: Preparation, creation, and/or presentation of the published work by members of the original research group, specifically critical review, commentary, or revision, including pre- or post-publication stages.

-
- [1] R. D. Peccei and Helen R. Quinn. Constraints imposed by CP conservation in the presence of pseudoparticles. *Phys. Rev. D*, 16:1791–1797, Sep 1977.
- [2] Steven Weinberg. A New Light Boson? *Phys. Rev. Lett.*, 40:223–226, Jan 1978.
- [3] R. D. Peccei and Helen R. Quinn. CP Conservation in the Presence of Pseudoparticles. *Phys. Rev. Lett.*, 38:1440–1443, Jun 1977.
- [4] Frank Wilczek. Problem of Strong P and T Invariance in the Presence of Instantons. *Phys. Rev. Lett.*, 40:279–282, Jan 1978.
- [5] Peter Svrcek and Edward Witten. Axions in string theory. *Journal of High Energy Physics*, 2006(06):051, 2006.
- [6] Jihn E. Kim. Light pseudoscalars, particle physics and cosmology. *Physics Reports*, 150(1-2):1–177, 1987.
- [7] Joerg Jaeckel and Andreas Ringwald. The low-energy frontier of particle physics. *Annual Review of Nuclear and Particle Science*, 60(1):405–437, 2010.
- [8] Asimina Arvanitaki, Savas Dimopoulos, Sergei Dubovsky, Nemanja Kaloper, and John March-Russell. String axiverse. *Phys. Rev. D*, 81:123530, Jun 2010.
- [9] John Preskill, Mark B. Wise, and Frank Wilczek. Cosmology of the invisible axion. *Physics Letters B*, 120(1):127–132, 1983.
- [10] L.F. Abbott and P. Sikivie. A cosmological bound on the invisible axion. *Physics Letters B*, 120(1):133–136, 1983.
- [11] David J. E. Marsh. Axion cosmology. *Physics Reports*, 643:1–79, 2016.
- [12] Peter W. Graham and Surjeet Rajendran. New observables for direct detection of axion dark matter. *Phys. Rev. D*, 88:035023, Aug 2013.
- [13] Jonathan L. Ouellet, Chiara P. Salemi, Joshua W. Foster, Reyco Henning, Zachary Bogorad, Janet M. Conrad, Joseph A. Formaggio, Yonatan Kahn, Joe Minervini, Alexey Radovinsky, Nicholas L. Rodd, Benjamin R. Safdi, Jesse Thaler, Daniel Winklehner, and Lindley Winslow. First results from abracadabra-10 cm: A search for sub- μeV axion dark matter. *Phys. Rev. Lett.*, 122:121802, Mar 2019.
- [14] N. Du, N. Force, R. Khatriwada, E. Lentz, R. Ottens, L. J. Rosenberg, G. Rybka, G. Carosi, N. Woollett, D. Bowring, A. S. Chou, A. Sonnenschein, W. Wester, C. Boutan, N. S. Oblath, R. Bradley, E. J. Daw, A. V. Dixit, J. Clarke, S. R. O’Kelley, N. Crisosto, J. R. Gleason, S. Jois, P. Sikivie, I. Stern, N. S. Sullivan, D. B. Tanner, and G. C. Hilton. Search for invisible axion dark matter with the axion dark matter experiment. *Phys. Rev. Lett.*, 120:151301, Apr 2018.
- [15] Alexander J. Millar, Steven M. Anlage, Rustam Balafendiev, Pavel Belov, Karl van Bibber, Jan Conrad, Marcel Demarteau, Alexander Droster, Katherine Dunne, Andrea Gallo Rosso, Jon E. Gudmundsson, Heather Jackson, Gagandeep Kaur, Tove Klaesson, Nolan Kowitt, Matthew Lawson, Alexander Leder, Akira Miyazaki, Sid Morampudi, Hiranya V. Peiris, Henrik S. Røising, Gaganpreet Singh, Dajie Sun, Jacob H. Thomas, Frank Wilczek, Stafford Withington, Mackenzie Wooten, Jens Dilling, Michael Febbraro, Stefan Knirck, and Claire Marvinney. Searching for dark matter with plasma haloscopes. *Phys. Rev. D*, 107:055013, Mar 2023.
- [16] Beatriz Aja, Sergio Arguedas Cuendis, Ivan Arregui, Eduardo Artal, R. Belén Barreiro, Francisco J Casas, Marina C de Ory, Alejandro Díaz-Morcillo, Luisa de la Fuente, Juan Daniel Gallego, et al. The canfranc axion detection experiment (CADEx): Search for axions at 90 GHz with kinetic inductance detectors. *Journal of Cosmology and Astroparticle Physics*, 2022(11):044, 2022.
- [17] Saebyeok Ahn, JinMyeong Kim, Boris I. Ivanov, Ohjoon Kwon, HeeSu Byun, Arjan F. van Loo, SeongTae Park, Junu Jeong, Soohyung Lee, Jinsu Kim, Çağlar Kutlu, Andrew K. Yi, Yasunobu Nakamura, Seonjeong Oh, Danho Ahn, SungJae Bae, Hyoungsoon Choi, Jihoon Choi, Yonuk Chong, Woohyun Chung, Violeta Gkika, Jihn E. Kim, Younggeun Kim, Byeong Rok Ko, Lino Miceli, Doyu Lee, Jiwon Lee, Ki Woong Lee, Myeong-Jae Lee, Andrei Matlashov, Pallavi Parashar, Taehyeon Seong, Yun Chang Shin, Sergey V. Uchaikin, SungWoo Youn, and Yannis K. Semertzidis. Extensive Search for Axion Dark Matter over 1 GHz with CAPP’S Main Axion Experiment. *Phys. Rev. X*, 14:031023, Aug 2024.
- [18] L. Brouwer, S. Chaudhuri, H.-M. Cho, J. Corbin, W. Craddock, C. S. Dawson, A. Droster, J. W. Foster, J. T. Fry, P. W. Graham, R. Henning, K. D. Irwin, F. Kadribasic, Y. Kahn, A. Keller, R. Kolevator, S. Kuenstner, A. F. Leder, D. Li, J. L. Ouellet, K. M. W. Pappas, A. Phipps, N. M. Rapidis, B. R. Safdi, C. P. Salemi, M. Simanovskaia, J. Singh, E. C. van Assendelft, K. van Bibber, K. Wells, L. Winslow, W. J. Wisniewski, and B. A. Young. Projected sensitivity of dmradio- m^3 : A search for the QCD axion below 1 μeV . *Phys. Rev. D*, 106:103008, Nov 2022.
- [19] David Alesini, Danilo Babusci, Paolo Beltrame, Fabio Bossi, Paolo Ciambrone, Alessandro D’Elia, Daniele Di Gioacchino, Giampiero Di Pirro, Babette Döbrich, Paolo Falferi, et al. The future search for low-frequency axions and new physics with the FLASH resonant cavity experiment at Frascati National Laboratories. *Physics of the Dark Universe*, 42:101370, 2023.
- [20] L. Zhong, S. Al Kenany, K. M. Backes, B. M. Brubaker, S. B. Cahn, G. Carosi, Y. V. Gurevich, W. F. Kindel, S. K. Lamoreaux, K. W. Lehnert, S. M. Lewis, M. Malnou, R. H. Maruyama, D. A. Palken, N. M. Rapidis, J. R. Root, M. Simanovskaia, T. M. Shokair, D. H. Speller, I. Urdinaran, and K. A. van Bibber. Results from phase 1 of the haystac microwave cavity axion experiment. *Phys. Rev. D*, 97:092001, May 2018.

- [21] J. Egge, D. Leppla-Weber, S. Knirck, B. Ary dos Santos Garcia, D. Bergermann, A. Caldwell, V. Dabhi, C. Diaconu, J. Diehl, G. Dvali, M. Ekmedžić, F. Gallo, E. Garutti, S. Heyminck, F. Hubaut, A. Ivanov, J. Jochum, P. Karst, M. Kramer, D. Kreikemeyer-Lorenzo, C. Krieger, C. Lee, A. Lindner, J. P. A. Maldonado, B. Majorovits, S. Martens, A. Martini, A. Miyazaki, E. Öz, P. Pralavorio, G. Raffelt, A. Ringwald, J. Redondo, S. Roset, N. Salama, J. Schaffran, A. Schmidt, F. Steffen, C. Strandhagen, I. Usherov, H. Wang, G. Wieching, G. Cancelo, M. Di Federico, G. Hoshino, and L. Stefanazzi. First search for dark photon dark matter with a madmax prototype. *Phys. Rev. Lett.*, 134:151004, Apr 2025.
- [22] Aaron P. Quiskamp, Graeme R. Flower, Steven Samuels, Ben T. McAllister, Paul Altin, Eugene N. Ivanov, Maxim Goryachev, and Michael E. Tobar. Near-quantum-limited axion dark matter search with the ORGAN experiment around 26 μeV . *Phys. Rev. D*, 111:095007, May 2025.
- [23] A. Rettaroli, D. Alesini, D. Babusci, C. Braggio, G. Carugno, D. D’Agostino, A. D’Elia, D. Di Gioacchino, R. Di Vora, P. Falferi, U. Gambardella, A. Gardikiotis, C. Gatti, C. Ligi, A. Lombardi, G. Maccarrone, A. Ortolan, G. Ruoso, S. Tocci, and G. Vidali. Search for axion dark matter with the QUAX-LNF tunable haloscope. *Phys. Rev. D*, 110:022008, Jul 2024.
- [24] Alexander V. Gramolin, Deniz Aybas, Dorian Johnson, Janos Adam, and Alexander O. Sushkov. Search for axion-like dark matter with ferromagnets. *Nature Physics*, 17(1):79–84, Jan 2021.
- [25] Daniel Gavilan-Martin, Grzegorz Lukasiewicz, Mikhail Padniuk, Emmanuel Klinger, Magdalena Smolis, Nataniel L Figueroa, Derek F Jackson Kimball, Alexander O Sushkov, Szymon Pustelny, Dmitry Budker, et al. Searching for dark matter with a spin-based interferometer. *Nature Communications*, 16(1):4953, May 2025.
- [26] Itay M. Bloch, Gil Ronen, Roy Shaham, Ori Katz, Tomer Volansky, and Or Katz. New constraints on axion-like dark matter using a Floquet quantum detector. *Science Advances*, 8(5):eabl8919, 2022.
- [27] C. Abel, N. J. Ayres, G. Ban, G. Bison, K. Bodek, V. Bondar, E. Chanel, C. B. Crawford, M. Daum, B. Dechenaux, S. Emmenegger, P. Flaux, W. C. Griffith, P. G. Harris, Y. Kermaidic, K. Kirch, S. Komposch, P. A. Koss, J. Krempel, B. Lauss, T. Lefort, Prajwal Mohan-Murthy, Oscar Naviliat Cuncic, D. Pais, F. M. Piegsa, Guillaume Pignol, M. Rawlik, D. Ries, Stephanie Roccia, D. Rozpedzik, Philipp Schmidt-Wellenburg, N. Severijns, Y. V. Stadnik, J. A. Thorne, A. Weis, E. Wursten, Jacek Zejma, and Geza Zsigmond. Search for ultralight axion dark matter in a side-band analysis of a ^{199}Hg free-spin precession signal. *SciPost Phys.*, 15:058, 2023.
- [28] S. Karanth, E. J. Stephenson, S. P. Chang, V. Hejny, S. Park, J. Pretz, Y. K. Semertzidis, A. Wirzba, A. Wrońska, F. Abusaif, A. Aggarwal, A. Aksentev, B. Alberdi, A. Andres, L. Barion, I. Bekman, M. Beyß, C. Böhme, B. Breitkreutz, C. von Byern, N. Canale, G. Ciullo, S. Dymov, N.-O. Fröhlich, R. Gebel, K. Grigoryev, D. Grzonka, J. Hetzel, O. Javakhishvili, H. Jeong, A. Kacharava, V. Kamerdzhiyev, I. Keshe-lashvili, A. Kononov, K. Laihem, A. Lehrach, P. Lenisa, N. Lomidze, B. Lorentz, A. Magiera, D. Mchedlishvili, F. Müller, A. Nass, N. N. Nikolaev, A. Pesce, V. Poncza, D. Prasuhn, F. Rathmann, A. Saleev, D. Shergelashvili, V. Shmakova, N. Shurkhno, S. Siddique, J. Slim, H. Soltner, R. Stassen, H. Ströher, M. Tabidze, G. Tagliente, Y. Valdau, M. Vitz, T. Wagner, and P. Wüstner. First search for axionlike particles in a storage ring using a polarized deuteron beam. *Phys. Rev. X*, 13:031004, Jul 2023.
- [29] Junyi Lee, Mariangela Lisanti, William A. Terrano, and Michael Romalis. Laboratory constraints on the neutron-spin coupling of fev-scale axions. *Phys. Rev. X*, 13:011050, Mar 2023.
- [30] Francesco Capozzi, Bhaskar Dutta, Gajendra Gurung, Wooyoung Jang, Ian M. Shoemaker, Adrian Thompson, and Jaehoon Yu. New constraints on alp couplings to electrons and photons from argoneut and the miniboone beam dump. *Phys. Rev. D*, 108:075019, Oct 2023.
- [31] Dmitry Budker, Peter W. Graham, Micah Ledbetter, Surjeet Rajendran, and Alexander O. Sushkov. Proposal for a Cosmic Axion Spin Precession Experiment (CASPER). *Phys. Rev. X*, 4:021030, May 2014.
- [32] Derek F. Jackson Kimball, S. Afach, D. Aybas, J. W. Blanchard, D. Budker, G. Centers, M. Engler, N. L. Figueroa, A. Garcon, P. W. Graham, H. Luo, S. Rajendran, M. G. Sendra, A. O. Sushkov, T. Wang, A. Wickenbrock, A. Wilzewski, and T. Wu. Overview of the cosmic axion spin precession experiment (CASPER). In Gianpaolo Carosi and Gray Rybka, editors, *Microwave Cavities and Detectors for Axion Research*, pages 105–121, Cham, 2020. Springer International Publishing.
- [33] Tao Wang, Derek F Jackson Kimball, Alexander O Sushkov, Deniz Aybas, John W Blanchard, Gary Centers, Sean R O’Kelley, Arne Wickenbrock, Jiancheng Fang, and Dmitry Budker. Application of spin-exchange relaxation-free magnetometry to the cosmic axion spin precession experiment. *Physics of the dark universe*, 19:27–35, 2018.
- [34] Deniz Aybas, Janos Adam, Emmy Blumenthal, Alexander V. Gramolin, Dorian Johnson, Annalies Kleyheeg, Samer Afach, John W. Blanchard, Gary P. Centers, Antoine Garcon, Martin Engler, Nataniel L. Figueroa, Marina Gil Sendra, Arne Wickenbrock, Matthew Lawson, Tao Wang, Teng Wu, Haosu Luo, Hamdi Mani, Philip Mauskopf, Peter W. Graham, Surjeet Rajendran, Derek F. Jackson Kimball, Dmitry Budker, and Alexander O. Sushkov. Search for axionlike dark matter using solid-state nuclear magnetic resonance. *Phys. Rev. Lett.*, 126:141802, Apr 2021.
- [35] Antoine Garcon, John W. Blanchard, Gary P. Centers, Nataniel L. Figueroa, Peter W. Graham, Derek F. Jackson Kimball, Surjeet Rajendran, Alexander O. Sushkov, Yevgeny V. Stadnik, Arne Wickenbrock, Teng Wu, and Dmitry Budker. Constraints on bosonic dark matter from ultralow-field nuclear magnetic resonance. *Science Advances*, 5(10):eaax4539, 2019.
- [36] Julian Walter, Olympia Maliaka, Yuzhe Zhang, John W. Blanchard, Gary Centers, Arian Dogan, Martin Engler, Nataniel L. Figueroa, Younggeun Kim, Derek F. Jackson Kimball, Matthew Lawson, Declan W. Smith, Alexander O. Sushkov, Dmitry Budker, Hendrik Bekker, and Arne Wickenbrock. Search for axionlike dark matter using liquid-state nuclear magnetic resonance. *Phys. Rev. D*, 112:052008, Sep 2025.
- [37] Deniz Aybas, Hendrik Bekker, John W. Blanchard,

- Dmitry Budker, Gary P. Centers, Nataniel L. Figueroa, Alexander V. Gramolin, Derek F. Jackson Kimball, Arne Wickenbrock, and Alexander O. Sushkov. Quantum sensitivity limits of nuclear magnetic resonance experiments searching for new fundamental physics. *Quantum Science and Technology*, 6(3):034007, jun 2021.
- [38] Jeff A. Dror, Stefania Gori, Jacob M. Leedom, and Nicholas L. Rodd. Sensitivity of spin-precession axion experiments. *Phys. Rev. Lett.*, 130:181801, May 2023.
- [39] Yuzhe Zhang, Deniz Aybas Tunturk, Hendrik Bekker, Dmitry Budker, Derek F. Jackson Kimball, Alexander O. Sushkov, and Arne Wickenbrock. Frequency-Scanning Considerations in Axionlike Dark Matter Spin-Precession Experiments. *Annalen der Physik*, 536(1):2300223, 2024.
- [40] Jeff A. Dror, Hitoshi Murayama, and Nicholas L. Rodd. Cosmic axion background. *Phys. Rev. D*, 103:115004, Jun 2021.
- [41] T. Nitta, T. Braine, N. Du, M. Guzzetti, C. Hanretty, G. Leum, L. J. Rosenberg, G. Rybka, J. Sinnis, John Clarke, I. Siddiqi, M. H. Awida, A. S. Chou, M. Hollister, S. Knirck, A. Sonnenschein, W. Wester, J. R. Gleason, A. T. Hipp, P. Sikivie, N. S. Sullivan, D. B. Tanner, R. Khatiwada, G. Carosi, N. Robertson, L. D. Duffy, C. Boutan, E. Lentz, N. S. Oblath, M. S. Taubman, J. Yang, E. J. Daw, M. G. Perry, C. Bartram, J. H. Buckley, C. Gaikwad, J. Hoffman, K. W. Murch, M. Goryachev, E. Hartman, B. T. McAllister, A. Quiskamp, C. Thomson, M. E. Tobar, J. A. Dror, H. Murayama, and N. L. Rodd. Search for a dark-matter-induced cosmic axion background with ADMX. *Phys. Rev. Lett.*, 131:101002, Sep 2023.
- [42] Ciaran AJ O'Hare and Giovanni Pierobon. Fine-grained dark matter substructure and axion haloscopes, 2025.
- [43] Xucheng Gan, Hyungjin Kim, and Andrea Mitridate. Probing quadratically coupled ultralight dark matter with pulsar timing arrays. *Phys. Rev. D*, 113:063034, Mar 2026.
- [44] Xiaofei Huang, Xiaolin Ma, Zitong Xu, Itay M. Bloch, and Kai Wei. Earth matter enhanced axion dark matter search, 2026.
- [45] The Astropy Collaboration, Robitaille, Thomas P., Tollerud, Erik J., Greenfield, Perry, Droettboom, Michael, Bray, Erik, Aldcroft, Tom, Davis, Matt, Ginsburg, Adam, Price-Whelan, Adrian M., Kerzendorf, Wolfgang E., Conley, Alexander, Crighton, Neil, Barbary, Kyle, Muna, Dimitri, Ferguson, Henry, Grollier, Frédéric, Parikh, Madhura M., Nair, Prasanth H., Günther, Hans M., Deil, Christoph, Woillez, Julien, Conseil, Simon, Kramer, Roban, Turner, James E. H., Singer, Leo, Fox, Ryan, Weaver, Benjamin A., Zalza, Victor, Edwards, Zachary I., Azalee Bostroem, K., Burke, D. J., Casey, Andrew R., Crawford, Steven M., Dencheva, Nadia, Ely, Justin, Jenness, Tim, Labrie, Kathleen, Lim, Pey Lian, Pierfederici, Francesco, Pontzen, Andrew, Ptak, Andy, Refsdal, Brian, Servillat, Mathieu, and Streicher, Ole. Astropy: A community python package for astronomy. *A&A*, 558:A33, 2013.
- [46] Charles R. Harris, K. Jarrod Millman, Stéfan J. van der Walt, Ralf Gommers, Pauli Virtanen, David Cournapeau, Eric Wieser, Julian Taylor, Sebastian Berg, Nathaniel J. Smith, Robert Kern, Matti Picus, Stephan Hoyer, Marten H. van Kerkwijk, Matthew Brett, Allan Haldane, Jaime Fernández del Río, Mark Wiebe, Pearu Peterson, Pierre Gérard-Marchant, Kevin Sheppard, Tyler Reddy, Warren Weckesser, Hameer Abbasi, Christoph Gohlke, and Travis E. Oliphant. Array programming with numpy. *Nature*, 585(7825):357–362, Sep 2020.
- [47] H. Y. Carr and E. M. Purcell. Effects of Diffusion on Free Precession in Nuclear Magnetic Resonance Experiments. *Phys. Rev.*, 94:630–638, May 1954.
- [48] Anatole Abragam and Herman Y. Carr. The principles of nuclear magnetism. *Physics Today*, 14:56–58, 1961.
- [49] N. Wyn Evans, Ciaran A. J. O'Hare, and Christopher McCabe. Refinement of the standard halo model for dark matter searches in light of the Gaia Sausage. *Phys. Rev. D*, 99:023012, Jan 2019.
- [50] Alexander V. Gramolin, Arne Wickenbrock, Deniz Aybas, Hendrik Bekker, Dmitry Budker, Gary P. Centers, Nataniel L. Figueroa, Derek F. Jackson Kimball, and Alexander O. Sushkov. Spectral signatures of axionlike dark matter. *Phys. Rev. D*, 105:035029, Feb 2022.
- [51] Pauli Virtanen, Ralf Gommers, Travis E. Oliphant, Matt Haberland, Tyler Reddy, David Cournapeau, Evgeni Burovski, Pearu Peterson, Warren Weckesser, Jonathan Bright, Stéfan J. van der Walt, Matthew Brett, Joshua Wilson, K. Jarrod Millman, Nikolay Mayorov, Andrew R. J. Nelson, Eric Jones, Robert Kern, Eric Larson, C J Carey, İlhan Polat, Yu Feng, Eric W. Moore, Jake VanderPlas, Denis Laxalde, Josef Perktold, Robert Cimrman, Ian Henriksen, E. A. Quintero, Charles R. Harris, Anne M. Archibald, Antônio H. Ribeiro, Fabian Pedregosa, Paul van Mulbregt, and SciPy 1.0 Contributors. SciPy 1.0: Fundamental Algorithms for Scientific Computing in Python. *Nature Methods*, 17:261–272, 2020.
- [52] Fernando Perez and Brian E. Granger. Ipython: A system for interactive scientific computing. *Computing in Science & Engineering*, 9(3):21–29, 2007.
- [53] Brian E. Granger and Fernando Pérez. Jupyter: Thinking and storytelling with code and data. *Computing in Science & Engineering*, 23(2):7–14, 2021.