Q0) I have seen and studied most of these models in a fair amount of detail in our differential equations class and some other statistics class. Except the Lanchester's model of war and Lorentz system which are new to me, and the three dimensional gravitation equations which we didn't cover thoroughly. I am familiar with most of the other equations.

Q1) - For both modelling decisions regarding human and rabbits population, the exponential growth/decay equations would be reasonable but limited because human population does not grow without bounds because of living resources and rabbits are hunted sometimes so we need some form of harvesting rate in the equation.
- I am not really familiar with yeast and dough but since we are talking about cells I think of exponential growth because cells multiply/divide a considerable amount of time so exponential growth/decay since reasonable.

Q2) I would be looking at the SIR model for this question.
Assumptions:
↳ People can recover from any disease modelled
↳ No acquired immunity from the disease
↳ No seclusion/confinement, interactions are always possible
↳ once recovered you can't be infected again or become susceptible
↳ Everyone has the same parameters for each group regardless of heterogeneity in the population
↳ There is no death

These assumptions might fail if we try to correctly model AIDS, once you are infected you can not recover, you can't have immunity so you are always susceptible to different variations of the disease, people are supposed to be secluded or they are often ostracized once they are infected so they can't infect, people get infected and die at different rates so AIDS challenge the limit of the SIR model.

Q3) My top priority would be to bring as much heterogeneity as I can in the model, especially the fact that women and men and adults and child, might not be affected the same way, I would also try to include some spatial dimension to model how the disease will spread in a metro-city with a higher capacity than some rural or sub-urban areas. To account for this I would need to know how much interaction there is among people, the distribution of ages, sexual orientation and city capacity. The transmission rate of the disease under different conditions such as weather and demography and socio-economic stability.

Q5) It would take the form:

$$\dot{r} = \boxed{rA\left(1-\frac{r}{C}\right)} - BrW$$

$$\dot{w} = CrW - dW$$

The system is no longer cyclic, the rabbit population first declines between $t=0$ and $t=2.5$ then increases between $t=2.5$ and $t=5$ to settle below the carrying capacity at time $t=10$ around 300 rabbits. The population of wolf declines minimally to remain constant at time $t=2$ around at around 0.

Q6) When r is much smaller than P, r doesn't contribute much to the expression so we get $-BP\frac{r}{P}W \approx -BrW$, so when rabbits are abundant and $r \ll P$ we get the same classical model, where wolf eat rabbit at rate Br.
When r is much larger than P, P doesn't contribute much to the expression so we get $-BP\frac{r}{r}W \approx -BPW$
When rabbits are abundant, wolf will consume that at a rate BP independent of r.

Q7) $\dot{w} = sW\left(1-\frac{W}{L(r)}\right)$

$L(r) = ar$, $a>0$

We see a similar dynamic as the previous model, except in this new model there is no temporary growth, we have a slight decline in term of rabbit population then it stabilizes at around 200. We get similar dynamic in term of wolf population as the previous model.

Q8)

$$\dot{F} = aF\left(1-\frac{F}{L(t)}\right) - H(F,t)$$ Account for seasonality

$L(t) = \alpha + \beta\sin(\pi t)$, $\alpha > \beta > 0$

$H(F,t) = d\,E(t)F$ ← effort (how much work do you put into fishing, labor)

↳ efficiency
more efficient means they good at catching fishes caught

Q9)

$$m\ddot{y} = -r\dot{y} - Ky$$

$$V(t) = y'(t),\ V'(t) = y''(t)$$

$$mV'(t) = -rV(t) - Ky(t)$$

$$V'(t) = -\frac{r}{m}V(t) - \frac{K}{m}y(t)$$

$$= -\frac{1}{m}\left(rV(t) + Ky(t)\right)$$