

Group Task 3: Hurricane Risk Assessment for Gulf of Mexico Cities

Yves-Langston Mays¹, Uyen Vi Phan², and Ny Dang³

¹Department of Natural Sciences & Mathematics, University of Houston Main Campus,
Houston, USA, ymmays@cougarnet.uh.edu

²Department of Natural Sciences & Mathematics, University of Houston Main Campus,
Houston, USA, uphan2@uh.edu

³Department of Natural Sciences & Mathematics, University of Houston Main Campus,
Houston, USA, tndang8@cougarnet.uh.edu

November 8, 2024

Abstract

Task 3 centers on hurricane risk for 25 cities in the Gulf of Mexico, using data from the Atlantic hurricane database (HURDAT2) from 1851 to 2023, provided by the National Hurricane Center. To analyze and assess this risk, we will perform 3 analyses in R to assess the hurricane risk on the Gulf of Mexico. First, we visualize and note our findings on the storm tracks over the last 25 years (1999-2024), focusing on storm paths, intensity, and duration at each location. Spatial correlation analysis will be used to explore the relationship between hurricane occurrences and contributing environmental factors, while Non-Parametric Density Estimation will estimate location-specific risk based on historical hurricane trajectories. These analyses collectively aim to identify the cities at highest risk of hurricane impact and gauge potential severity.

Contents

1.0	Introduction	3
2.0	Background	4
3.0	Methodology	5
3.1	3.1 Data Collection and Preparation	5
3.2	3.2 Geographic Data Setup	5
3.3	3.3 Visualization	8
3.4	3.4 Statistical Analysis	9
4.0	Results	13
5.0	Discussion	14
6.0	Conclusion	15
	References	15

1.0 Introduction

One of the most common natural disaster plaguing the Gulf of Mexico are Hurricanes. Just this year (2024) there has been 9 hurricanes in the Atlantic Ocean including Beryl, Helen and Milton. These storms can have lasting impacts to people's lives, the environment, infrastructure and to the economy. Since 1980, hurricane damage has costed over \$1.3 trillion in damages with an average of \$22.8 billion dollars per event and 6,890 deaths [1]. Learning to predict where, when, and the intensity of hurricanes can not only save us billions of dollars but thousands of lives as well. This is especially important for cities that are at high risk such as cities like New Orleans that exists at extremely low elevations.

This task aims to use historical data on hurricanes to predict future hurricane activity and highlight cities that are at most risk. The historical hurricane data from the National Hurricane Center's HURDAT2 database, contains hurricane data in the Atlantic from 1851 to 2023. HURDAT2 records the six-hourly information on the location, maximum winds, central pressure, and (beginning in 2004) size of all known tropical cyclones and subtropical cyclones[2]. Along with a list of 25 cities and their locations in the Gulf of Mexico, 3 analyses will be performed oh HURDAT2 to analyze and predict the storm tracks in the Gulf of Mexico.

The last 25 (1999-2024) years worth of storm tracks will be first visualized and analyzed on the over the Gulf of Mexico to identify common patterns and trends of the storm tracks. The visualization and the manual analysis provides a good general overview of how these storms move, where they are the most intense, and what places receive the most storms.

The second analysis examines how certain factors like sea surface temperatures and El Nino/La Nina patterns affect hurricane activity. Spatial correlation analysis between these factors and hurricane activity will be used to achieve this. The correlation analysis will show how certain weather phenomena can affect hurricane activity to better predict hurricane activity.

The last analysis uses non-parametric density estimation to assess the hurricane risk

based on past trajectories and severity. This can provide insight on where hurricanes are more likely to appear and what routes they take. It can assess what regions will receive the most intense storms. Paired with the spatial correlation analysis, the information provided from the non-parametric density estimation can highlight what cities are most at risk in the Gulf of Mexico.

These 3 analyses can provide valuable insight to the behavior and activity of future hurricanes. Knowing the behavior and patterns of the hurricanes, knowing what cities and regions are at the most risk from hurricane activity and how different factors play a role in hurricane activity can help people better prepare for hurricanes and minimize the losses caused by these storms. Meteorologists can use these predictions to better inform people about the route, severity of storms and what to expect as it passes. Governments can use this information to predict what cities will require the most aid. Insurance companies can use this data to decide what services are best suited to a specific location.

2.0 Background

The data source, HURDAT2, short for Hurricane Database 2 from the National Hurricane Center, provides detailed hurricane tracks from 1851 to 2023. This database records measurements of critical parameters including location coordinates, maximum wind speeds, and central pressure. Since 2004, the database has also included storm size data, enhancing our understanding of hurricane characteristics.

Our report encompasses 25 cities across the Gulf of Mexico region, representing diverse geographical and political jurisdictions. Ten U.S. cities form the northern boundary of the study area, including major metropolitan centers like New Orleans, Houston, and Miami, alongside significant coastal cities such as Tampa, Corpus Christi, Pensacola, Mobile, Galveston, Biloxi, and Key West. Along the western and southern Gulf coast, nine Mexican cities are included: Veracruz, Tampico, and Campeche serve as major ports, while Cancún repre-

sents a vital tourism center. The study also includes Mérida, Ciudad del Carmen, Progreso, Coatzacoalcos, and Tuxpan, which are crucial to Mexico’s coastal economy. The Caribbean region is represented by Havana, Varadero, and Cienfuegos, along with key island locations including Belize City, George Town, and Nassau.

Natural factors significantly influence hurricane activity in the Gulf region, with varying degrees of impact severity. Sea surface temperatures directly affect hurricane formation and intensification, while El Niño/La Niña patterns influence atmospheric conditions and storm frequency. The Atlantic Multidecadal Oscillation affects long-term hurricane activity cycles, and Saharan dust levels can suppress hurricane development. Upper-level wind patterns play a crucial role in determining storm trajectories and development potential.

Each city’s unique geographical position and environmental characteristics contribute to its specific hurricane risk profile. This research aims to quantify specific hurricane risks for each city, identify correlations between environmental factors and hurricane patterns, and develop location-specific risk assessment models.

The integration of these methods below will support and inform decision-making processes for protecting coastal communities from future hurricane impacts.

3.0 Methodology

3.1 3.1 Data Collection and Preparation

3.2 3.2 Geographic Data Setup

Study Area Boundaries:

Latitude: 16.5 to 31.7 °N


Longitude: -98.9 to -76.4 °W

Total storm observations: 13669

Unique storms: 961


Date range: 18510625 to 20230830

3.3 3.3 Visualization




3.4 3.4 Statistical Analysis


```
ggplot(yearly_storms, aes(x = YEAR, y = storm_count)) +
  geom_line(color = "blue") +
  geom_point() +
  labs(title = "year storm count from 99-23",
        x = "Year",
        y = "# of Storms")
```



```
ggplot(monthly_storms, aes(x = MONTH, y = storm_count)) +
  geom_col(fill = "steelblue") +
  labs(title = "monthly storm freq",
        x = "Month",
        y = "# of Storms") +
  scale_x_continuous(breaks = 1:12,
                     labels = c("Jan", "Feb", "Mar", "Apr", "May", "Jun",
                                "Jul", "Aug", "Sep", "Oct", "Nov", "Dec"))
```




```
ggplot(status_count, aes(x = STATUS, y = count, fill = STATUS)) +
  geom_bar(stat = "identity") +
  labs(title = "strm count by status",
        x = "Status",
        y = "# of Storms") +
  scale_fill_manual(values = c("HU" = "red", "TS" = "orange", "TD" = "blue"))
```



```
intensity_trend <- recent_gulf_storms %>%
  group_by(YEAR) %>%
  summarize(avg_windspeed = mean(WINDSPEED_KT, na.rm = TRUE))

ggplot(intensity_trend, aes(x = YEAR, y = avg_windspeed)) +
  geom_line(color = "darkred") +
  geom_point() +
  labs(title = "Avg Storm Intensity Over Time 99-23",
       x = "Year",
       y = "Avg Wind Speed (kt)")
```



4.0 Results

Discuss results of analysis

5.0 Discussion

Discuss the implications and significance here

6.0 Conclusion

Conclusion Here

References