

**A FIRST COURSE  
IN  
ANALYSIS**



---

# A FIRST COURSE IN ANALYSIS

## MAT2006 Notebook

---

### Lecturer

*Prof. Weiming Ni*

*The Chinese University of Hongkong, Shenzhen*

### Tex Written By

*Mr. Jie Wang*

*The Chinese University of Hongkong, Shenzhen*



香港中文大學(深圳)

The Chinese University of Hong Kong, Shenzhen



# Contents

Acknowledgments	ix
Notations	xi
<b>1 Week1</b>	<b>1</b>
1.1 Wednesday	1
1.1.1 Introduction to Set	1
1.2 Quiz	5
1.3 Friday	6
1.3.1 Proof of Schroder-Bernstein Theorem	6
1.3.2 Connectedness of Real Numbers	10
<b>2 Week2</b>	<b>13</b>
2.1 Wednesday	13
2.1.1 Review and Announcement	13
2.1.2 Irrational Number Analysis	13
2.2 Friday	21
2.2.1 Set Analysis	21
2.2.2 Set Analysis Meets Sequence	22
2.2.3 Completeness of Real Numbers	23
<b>3 Week3</b>	<b>27</b>
3.1 Tuesday	27
3.1.1 Application of Heine-Borel Theorem	27
3.1.2 Set Structure Analysis	29
3.1.3 Reviewing	31

<b>3.2</b>	<b>Friday</b>	<b>33</b>
3.2.1	Review . . . . .	33
3.2.2	Continuity Analysis . . . . .	34
<b>4</b>	<b>Week4 . . . . .</b>	<b>41</b>
<b>4.1</b>	<b>Wednesday</b>	<b>41</b>
4.1.1	Function Analysis . . . . .	41
4.1.2	Continuity Analysis . . . . .	46
<b>4.2</b>	<b>Friday</b>	<b>50</b>
4.2.1	Continuity Analysis . . . . .	50
4.2.2	Monotone Analysis . . . . .	53
4.2.3	Cantor Set . . . . .	55
<b>5</b>	<b>Week5 . . . . .</b>	<b>59</b>
<b>5.1</b>	<b>Wednesdays</b>	<b>59</b>
5.1.1	Differentiation . . . . .	59
5.1.2	Basic Rules of Differentiation . . . . .	61
5.1.3	Analysis on Differential Calculus . . . . .	62
<b>5.2</b>	<b>Friday</b>	<b>67</b>
5.2.1	Analysis on Derivative . . . . .	67
5.2.2	Analysis on Mean-Value Theorem . . . . .	68
<b>5.3</b>	<b>Saturday: Comments on Quiz 1</b>	<b>73</b>
5.3.1	First Question . . . . .	73
5.3.2	Second Question . . . . .	73
5.3.3	Third Question . . . . .	74
5.3.4	Fourth Question . . . . .	75
5.3.5	Fifth Question . . . . .	75
5.3.6	Grading policy . . . . .	76

<b>6</b>	<b>Week6</b>	<b>77</b>
<b>6.1</b>	<b>Wednesday</b>	<b>77</b>
6.1.1	Reviewing	77
6.1.2	Convergence Analysis	78
<b>6.2</b>	<b>Friday</b>	<b>83</b>
6.2.1	Recap	83
6.2.2	Riemann Integration	84
<b>7</b>	<b>Week7</b>	<b>89</b>
<b>7.1</b>	<b>Wednesday</b>	<b>89</b>
7.1.1	Integrable Analysis	89
7.1.2	Elementary Calculus Analysis	92
<b>7.2</b>	<b>Friday</b>	<b>97</b>
7.2.1	Improper Integrals	97
<b>8</b>	<b>Week8</b>	<b>105</b>
<b>8.1</b>	<b>Friday</b>	<b>105</b>
8.1.1	Introduction to metric space	105
<b>9</b>	<b>Week9</b>	<b>111</b>
<b>9.1</b>	<b>Friday</b>	<b>111</b>
9.1.1	Preliminaries	112
9.1.2	Differentiation	114





# Acknowledgments

This book is taken notes from the MAT2006 in fall semester, 2018. These lecture notes were taken and compiled in  $\text{\LaTeX}$  by Jie Wang, an undergraduate student in Fall 2018. Prof. Weiming Ni has not edited this document. Students taking this course may use the notes as part of their reading and reference materials. This version of the lecture notes were revised and extended for many times, but may still contain many mistakes and typos, including English grammatical and spelling errors, in the notes. It would be greatly appreciated if those students, who will use the notes as their reading or reference material, tell any mistakes and typos to Jie Wang for improving this notebook.



# Notations and Conventions

$\mathbb{R}^n$	$n$ -dimensional real space
$\mathbb{C}^n$	$n$ -dimensional complex space
$\mathbb{R}^{m \times n}$	set of all $m \times n$ real-valued matrices
$\mathbb{C}^{m \times n}$	set of all $m \times n$ complex-valued matrices
$x_i$	$i$ th entry of column vector $\mathbf{x}$
$a_{ij}$	$(i, j)$ th entry of matrix $\mathbf{A}$
$\mathbf{a}_i$	$i$ th column of matrix $\mathbf{A}$
$\mathbf{a}_i^T$	$i$ th row of matrix $\mathbf{A}$
$\mathbb{S}^n$	set of all $n \times n$ real symmetric matrices, i.e., $\mathbf{A} \in \mathbb{R}^{n \times n}$ and $a_{ij} = a_{ji}$ for all $i, j$
$\mathbb{H}^n$	set of all $n \times n$ complex Hermitian matrices, i.e., $\mathbf{A} \in \mathbb{C}^{n \times n}$ and $\bar{a}_{ij} = a_{ji}$ for all $i, j$
$\mathbf{A}^T$	transpose of $\mathbf{A}$ , i.e, $\mathbf{B} = \mathbf{A}^T$ means $b_{ji} = a_{ij}$ for all $i, j$
$\mathbf{A}^H$	Hermitian transpose of $\mathbf{A}$ , i.e, $\mathbf{B} = \mathbf{A}^H$ means $b_{ji} = \bar{a}_{ij}$ for all $i, j$
$\text{trace}(\mathbf{A})$	sum of diagonal entries of square matrix $\mathbf{A}$
$\mathbf{1}$	A vector with all 1 entries
$\mathbf{0}$	either a vector of all zeros, or a matrix of all zeros
$\mathbf{e}_i$	a unit vector with the nonzero element at the $i$ th entry
$\mathcal{C}(\mathbf{A})$	the column space of $\mathbf{A}$
$\mathcal{R}(\mathbf{A})$	the row space of $\mathbf{A}$
$\mathcal{N}(\mathbf{A})$	the null space of $\mathbf{A}$
$\text{Proj}_{\mathcal{M}}(\mathbf{A})$	the projection of $\mathbf{A}$ onto the set $\mathcal{M}$



# Chapter 9

## Week9

### 9.1. Friday

We are now in the multi-variate differentiation part.

**Comments on question in last lecture.** The question left in the last lecture is that

In dimension  $\mathbb{R}^k$  with  $k$  to be determined, is it possible to find the smallest  $k$  such that a sphere  $S^2$  and a circle  $S^1$  have a way of putting to make each point from  $S^2$  to  $S^1$  have the same distance?

The answer is  $k = 5$ , Let's give an example. We define the sphere and the circle to be

$$S^2 = \{(x, y, z, 0, 0) \mid x^2 + y^2 + z^2 = 1\},$$

$$S^1 = \{(0, 0, 0, u, v) \mid u^2 + v^2 = 1\}$$

Therefore, the distance between any two points on the sphere and the circle, respectively, is

$$d = \sqrt{x^2 + y^2 + z^2 + u^2 + v^2} \equiv \sqrt{2}$$

Why  $k \leq 4$  is not ok? Now we give a instructive proof:

*Proof.* For the case  $k \leq 4$ , let  $\mathbf{c}_2$  denote the center of sphere  $S^2$  with radius  $r_2$ ;  $\mathbf{c}_3$  denote the center of circle  $S^1$  with radius  $r_1$ . Any point on  $S^2$  can be written as  $\mathbf{c}_2 + \mathbf{x}$  with  $\mathbf{x} \in M := \{\mathbf{x} \in \mathbb{R}^3 : \|\mathbf{x}\| = r_2\}$ ; and any point on  $S^1$  can be written as  $\mathbf{c}_1 + \mathbf{y}$  with  $\mathbf{x} \in N := \{\mathbf{y} \in \mathbb{R}^2 : \|\mathbf{y}\| = r_1\}$ . It follows that the distance between any two points can

be expressed as:

$$d(\mathbf{c}_2 + \mathbf{x}, \mathbf{c}_1 + \mathbf{y}) = \|(\mathbf{c}_2 + \mathbf{x}) - (\mathbf{c}_1 + \mathbf{y})\| = \|\mathbf{x}\|^2 + \|\mathbf{c}_2 - \mathbf{y} - \mathbf{c}_1\|^2 + 2\langle \mathbf{x}, \mathbf{c}_2 - \mathbf{y} - \mathbf{c}_1 \rangle$$

Note that the distance between any point in  $S^2$  and any point in  $S^1$  is the same. In particular,  $d(\mathbf{c}_2 + \mathbf{x}, \mathbf{c}_1 + \mathbf{y}) = d(\mathbf{c}_2 - \mathbf{x}, \mathbf{c}_1 + \mathbf{y})$ , which implies  $\langle \mathbf{x}, \mathbf{c}_2 - \mathbf{y} - \mathbf{c}_1 \rangle = 0, \forall \mathbf{x} \in M, \mathbf{y} \in N$ , i.e.,

$$\mathbf{c}_2 - \mathbf{c}_1 + N \subseteq M^\perp,$$

and therefore

$$\begin{aligned} \dim(\mathbf{c}_2 - \mathbf{c}_1 + N) &= \dim(N) = 2 \\ &\leq \dim(M^\perp) = k - 3, \end{aligned}$$

i.e.,  $k \leq 5$  is the sufficient condition of the problem. ■

### 9.1.1. Preliminaries

**Notations.** Here we use lower-case and bolded alphabet to denote a vector, e.g.,  $\mathbf{x}$ ; upper-case and bolded alphabet to denote a matrix, e.g.,  $\mathbf{A}$ ; unbolded alphabet to denote a scalar, e.g.,  $x_1$  or  $A_1$

Define  $\mathbf{x} := (x_1, \dots, x_m)$  and  $\mathbf{y} := (y_1, \dots, y_m)$ , we define the  $L_2$  norm

$$\begin{aligned} d(\mathbf{x}, \mathbf{y}) &:= \|\mathbf{x} - \mathbf{y}\| \\ &= [(x_1 - y_1)^2 + \dots + (x_m - y_m)^2]^{1/2} \end{aligned}$$

and the  $L_1, L_S$  (sup) norm:

$$\begin{aligned} d_1(\mathbf{x}, \mathbf{y}) &:= \|\mathbf{x} - \mathbf{y}\| := |x_1 - y_1| + \dots + |x_m - y_m| \\ d_S(\mathbf{x}, \mathbf{y}) &:= \|\mathbf{x} - \mathbf{y}\|_S := \max_{1 \leq i \leq m} (x_i - y_i) \end{aligned}$$

**Definition 9.1** [Vector Norm] A norm  $\|\cdot\|$  is a function from a vector space  $X$  to  $\mathbb{R}$  such that

1.  $\|\mathbf{x}\| \geq 0$ ,  $\forall \mathbf{x} \in X$ ; and  $\|\mathbf{x}\| = 0$  iff  $\mathbf{x} = \mathbf{0}$ ;
2.  $\|\lambda \mathbf{x}\| = |\lambda| \|\mathbf{x}\|$ , for  $\forall \mathbf{x} \in X, \lambda \in \mathbb{R}$ ;
3.  $\|\mathbf{x}_1 + \mathbf{x}_2\| \leq \|\mathbf{x}_1\| + \|\mathbf{x}_2\|$

**R** Any norm defines a metric:  $d(\mathbf{x}, \mathbf{y}) = \|\mathbf{x} - \mathbf{y}\|$ . From now on, we pre-assume the norm to be  $L_2$  norm and the metric to be  $L_2$  metric, unless specified.

The norm are used to measure the length of a vector, or the distance of two vectors. Correspondingly, the inner product can be used to measure the angle between two angles:

**Definition 9.2** [Inner Product] An **inner product** is a binary operation function  $\langle \cdot, \cdot \rangle : X \times X \mapsto \mathbb{R}$  such that

1.  $\langle x, x \rangle \geq 0$  for  $\forall x \in X$  and  $\langle x, x \rangle = 0$  iff  $\mathbf{x} = \mathbf{0}$
2.  $\langle \mathbf{x}, \mathbf{y} \rangle = \langle \mathbf{y}, \mathbf{x} \rangle$ , for  $\forall \mathbf{x}, \mathbf{y} \in X$
3.  $\langle \lambda \mathbf{x}, \mathbf{y} \rangle = \lambda \langle \mathbf{x}, \mathbf{y} \rangle$  for  $\forall \mathbf{x}, \mathbf{y} \in X$  and  $\forall \lambda \in \mathbb{R}$
4.  $\langle \mathbf{x} + \mathbf{y}, \mathbf{z} \rangle = \langle \mathbf{x}, \mathbf{z} \rangle + \langle \mathbf{y}, \mathbf{z} \rangle$

**R**

1. Any inner product always defines a norm, i.e.,  $\|\mathbf{x}\| = \sqrt{\langle \mathbf{x}, \mathbf{x} \rangle}$ .
2. The angle  $\theta$  between vector  $\mathbf{x}, \mathbf{y}$  is defined as:

$$\theta = \cos^{-1} \left( \frac{\langle \mathbf{x}, \mathbf{y} \rangle}{\|\mathbf{x}\| \|\mathbf{y}\|} \right)$$

In particular,  $\mathbf{x}$  and  $\mathbf{y}$  are orthogonal, i.e.,  $\langle \mathbf{x}, \mathbf{y} \rangle = 0$  if  $\theta = \pm \frac{\pi}{2}$ ;  $\mathbf{x}, \mathbf{y}$  are parallel if  $\theta = 0$ , or  $\theta = \pm \pi$ .

## 9.1.2. Differentiation

**Review for One-dimension.** Given a function  $f : \mathbb{R} \mapsto \mathbb{R}$ , the derivative is defined as

$$f'(x_0) = \lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0}$$

or equivalently, the value  $f'(x_0)$  is said to be the derivative of  $f$  if it satisfies the equality:

$$\begin{aligned} 0 &= \lim_{x \rightarrow x_0} \left[ \frac{f(x) - f(x_0)}{x - x_0} - f'(x_0) \right] \\ &= \lim_{x \rightarrow x_0} \frac{f(x) - f(x_0) - f'(x_0)(x - x_0)}{x - x_0} \\ &= \lim_{x \rightarrow x_0} \left| \frac{f(x) - [f(x_0) + f'(x_0)(x - x_0)]}{x - x_0} \right| \end{aligned}$$

The interpretation is that the affine (linear function)  $f(x_0) + f'(x_0)(x - x_0)$  approximates  $f$  near  $x_0$  in at least first order. We can use the similar way to define the high-dimension derivative.

### High-Dimension Derivative.

**Definition 9.3** [Differentiable] A map  $f : U \mapsto \mathbb{R}^n$ , where  $U$  is open in  $\mathbb{R}^m$ , is **differentiable** at  $\mathbf{x}_0 \in U$  if

$$\lim_{\mathbf{x} \rightarrow \mathbf{x}_0} \frac{\|f(\mathbf{x}) - f(\mathbf{x}_0) - \mathbf{L}(\mathbf{x}_0)(\mathbf{x} - \mathbf{x}_0)\|}{\|\mathbf{x} - \mathbf{x}_0\|} = 0, \quad (9.1)$$

where  $\mathbf{L}(\mathbf{x}_0)$  is said to be the derivative of  $f(\mathbf{x})$  at  $\mathbf{x} = \mathbf{x}_0$ , which is often denoted as  $Df(\mathbf{x}_0)$ , or  $f'(\mathbf{x}_0)$ . ■

**R** Note that  $f(\mathbf{x}), f(\mathbf{x}_0) \in \mathbb{R}^n$ , but  $(\mathbf{x} - \mathbf{x}_0) \in \mathbb{R}^m$ , thus  $\mathbf{L}(\mathbf{x}_0)$  is a **linear transformation** from  $\mathbb{R}^m$  to  $\mathbb{R}^n$ , i.e., a  $n \times m$  matrix.

**Interpretation.** We re-write  $f(\mathbf{x}) : \mathbb{R}^m \mapsto \mathbb{R}^n$  (a  $n$ -vector valued function of a  $m$ -vector argument) as:

$$f(\mathbf{x}) = \begin{pmatrix} f_1(\mathbf{x}) & \cdots & f_n(\mathbf{x}) \end{pmatrix},$$



with each  $f_i : \mathbb{R}^m \mapsto \mathbb{R}$  being a scalar-valued function of a  $m$ -vector argument. Let's study only one component with 2-argument function first, i.e.,  $n = 1, m = 2$ .




Figure 9.1: Diagram for partial derivatives

Given a surface  $S : z = f(x_1, x_2)$ , we study a point on the surface  $S$ , say  $M_0(x_1, x_2, f(x_1, x_2))$ . If the surface  $S$  intersects the plane  $y = x_2$  with the point  $M_0$ , then the **partial derivative**  $\frac{\partial f}{\partial x_1}(x_1, x_2)$  denotes the slope of the tangent line at  $M_0$ ; the tangent line  $T_{x_1}$  can be denoted as a vector  $T_{x_1} := (1, 0, \frac{\partial f}{\partial x_1}(x_1, x_2))$ .

Similarly, if the surface  $S$  intersects the plane  $x = x_1$  with the point  $M_0$ , then  $\frac{\partial f}{\partial x_2}(x_1, x_2)$  denotes the slope of the tangent line at  $M_0$ ; the tangent line  $T_{x_2}$  can be denoted as a vector  $T_{x_2} := (0, 1, \frac{\partial f}{\partial x_2}(x_1, x_2))$ .

Furthermore, the plane  $\text{span}\{T_{x_1}, T_{x_2}\}$  denotes the tangent plane at point  $M_0$ .

**Corollary 9.1**  $f$  is differentiable at  $\mathbf{x}_0$  implies all partial derivatives of  $f$  at  $\mathbf{x}_0$  exist.

The converse is not true. Let's raise a counter-example to explain that.

■ **Example 9.1**

$$f(x_1, x_2) = \begin{cases} \frac{x_1 x_2}{x_1^2 + x_2^2}, & (x_1, x_2) \neq (0, 0) \\ 0, & (x_1, x_2) = (0, 0) \end{cases}$$

When  $x_2 = mx_1$ , we have

$$f(x_1, mx_1) = \frac{mx_1^2}{x_1^2 + mx_1^2} = \frac{m}{1 + m^2},$$

i.e.,  $f$  is not differentiable at the origin as it is not continuous at the origin.

However, we can verify that  $\frac{\partial f}{\partial x_1}(0, 0) = 0 = \frac{\partial f}{\partial x_2}(0, 0)$ .

What guarantees  $f$  to be differentiable if all partial derivatives exist?

$$\frac{\|f(\mathbf{x}) - f(\mathbf{x}_0) - Df(\mathbf{x}_0)(\mathbf{x} - \mathbf{x}_0)\|}{\|\mathbf{x} - \mathbf{x}_0\|} \rightarrow 0,$$

where  $f : \mathbb{R}^m \mapsto \mathbb{R}^n$  and  $Df(\mathbf{x}_0)$  be a  $n \times m$  matrix. We re-write the formula for the derivative of  $f$  at  $\mathbf{x} = \mathbf{x}_0$  first. Here we write  $f(\mathbf{x}_0)$  in column vector form:

$$\begin{pmatrix} f_1(\mathbf{x}) \\ f_2(\mathbf{x}) \\ \vdots \\ f_n(\mathbf{x}) \end{pmatrix} = \begin{pmatrix} f_1(\mathbf{x}_0) \\ f_2(\mathbf{x}_0) \\ \vdots \\ f_n(\mathbf{x}_0) \end{pmatrix} + \begin{pmatrix} \frac{\partial f_1}{\partial x_1}(\mathbf{x}_0) & \frac{\partial f_1}{\partial x_2}(\mathbf{x}_0) & \cdots & \frac{\partial f_1}{\partial x_m}(\mathbf{x}_0) \\ \frac{\partial f_2}{\partial x_1}(\mathbf{x}_0) & \frac{\partial f_2}{\partial x_2}(\mathbf{x}_0) & \cdots & \frac{\partial f_2}{\partial x_m}(\mathbf{x}_0) \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial f_n}{\partial x_1}(\mathbf{x}_0) & \frac{\partial f_n}{\partial x_2}(\mathbf{x}_0) & \cdots & \frac{\partial f_n}{\partial x_m}(\mathbf{x}_0) \end{pmatrix} \begin{pmatrix} x_1 - x_{0,1} \\ x_2 - x_0 \\ \vdots \\ x_m - x_0 \end{pmatrix} + o(\|\mathbf{x} - \mathbf{x}_0\|), \quad (9.2)$$

where  $o(\|\mathbf{x} - \mathbf{x}_0\|)$  is a  $m \times 1$  vector that has order less than  $\|\mathbf{x} - \mathbf{x}_0\|$ .

Each entry in LHS of (9.2) can be expressed as:

$$f_i(\mathbf{x}) = f_i(\mathbf{x}_0) + \nabla^T f_i(\mathbf{x}_0) \cdot (\mathbf{x} - \mathbf{x}_0) + o(\|\mathbf{x} - \mathbf{x}_0\|),$$

with  $\nabla^T f_1(\mathbf{x}_0) = \left( \frac{\partial f_1}{\partial x_1}(\mathbf{x}_0) \quad \cdots \quad \frac{\partial f_1}{\partial x_m}(\mathbf{x}_0) \right)$  to be a row vector, and  $(\mathbf{x} - \mathbf{x}_0)$  to be a column vector.

**Sufficient Condition for differentiability.** Recall that we have faced a function that is everywhere differentiable but nowhere monotone, which is very counter-intuitive. If adding the condition that such function is continuously differentiable, then such function is monotone. Similarly, the gap for corollary(9.1) is continuous differentiability.

**Theorem 9.1** Let  $f : U \mapsto \mathbb{R}^n$ , where  $U \subseteq \mathbb{R}^m$  is open. If all partial derivatives of  $f$  are **continuous** in  $U$ , then  $f$  is differentiable in  $U$ .

*Proof for  $n = 2, m = 1$  case.* Consider  $f : \mathbb{R}^2 \mapsto \mathbb{R}$ , for  $x_0 \in U$  with small  $h, k$ , we have:

$$f(x_0 + h, y_0 + k) - f(x_0, y_0) = [f(x_0 + h, y_0 + k) - f(x_0 + h, y_0)] + [f(x_0 + h, y_0) - f(x_0, y_0)] \quad (9.3a)$$

$$\text{(by Mean Value Theorem, } \exists(c, d) \in [0, h] \times [0, k] \text{ such that)} \quad (9.3b)$$

$$= k \frac{\partial f}{\partial y}(x_0 + h, y_0 + c) + h \frac{\partial f}{\partial x}(x_0 + d, y_0) \quad (9.3c)$$

$$= k \frac{\partial f}{\partial y}(x_0, y_0 + c) + o(h) + h \frac{\partial f}{\partial x}(x_0, y_0) + o(h) \quad (9.3d)$$

$$= k \frac{\partial f}{\partial y}(x_0, y_0) + h \frac{\partial f}{\partial x}(x_0, y_0) + o(h) + o(k) \quad (9.3e)$$

Or we write it in compact matrix form:

$$f(x_0 + h, y_0 + k) = f(x_0, y_0) + \begin{pmatrix} \frac{\partial f}{\partial x}(x_0, y_0) & \frac{\partial f}{\partial y}(x_0, y_0) \end{pmatrix} \begin{pmatrix} h \\ k \end{pmatrix} + o(\|(h, k)\|),$$

i.e., the derivative of  $f$  at  $x = x_0$  exists, say  $(\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y})$ . Thus  $f$  is differentiable. ■

The same proof can easily extend to  $f : \mathbb{R}^n \mapsto \mathbb{R}$  by adding and subtracting  $n - 1$  terms and then apply mean value theorem for  $n$  times; the extension to  $f : \mathbb{R}^n \mapsto \mathbb{R}^m$  is clear by the same proof for each component.

*Proof.* Consider in general  $f : \mathbb{R}^n \mapsto \mathbb{R}^m$ , and it suffices to show (9.2), and moreover, it

suffices to show each entry  $f_i(\mathbf{x})$  in (9.2) can be expressed as

$$f_i(\mathbf{x}) = f_i(\mathbf{x}_0) + \nabla^T f_i(\mathbf{x}_0) \cdot (\mathbf{x} - \mathbf{x}_0) + o(\|\mathbf{x} - \mathbf{x}_0\|).$$

Let's abuse notation to refer  $g$  as the  $i$ -th entry of  $f_i$ , i.e.,  $g : \mathbb{R}^n \mapsto \mathbb{R}$ .

Let  $\varepsilon > 0$ , by continuity of partial derivatives, we can pick  $\delta > 0$  such that for any  $\|\mathbf{h}\| < \delta$ , we have

$$\left\| \frac{\partial g}{\partial x_k}(\mathbf{x} + \mathbf{h}) - \frac{\partial g}{\partial x_k}(\mathbf{x}) \right\| < \varepsilon, \quad k = 1, 2, \dots, n \quad (9.4)$$

Let  $\mathbf{L} = (\frac{\partial g}{\partial x_1}(\mathbf{x}), \dots, \frac{\partial g}{\partial x_n}(\mathbf{x}))$ ;  $\phi_k(\mathbf{h}) = (h_1, h_2, \dots, h_k, 0, \dots, 0)$ ,  $\phi_0(\mathbf{h}) = \mathbf{0}$ , and suppose  $\|\mathbf{h}\| < \delta$ , then we have:

$$|g(\mathbf{x} + \mathbf{h}) - g(\mathbf{x}) - \mathbf{L}\mathbf{h}| = \left| \sum_{k=1}^n [g(\mathbf{x} + \phi_k(\mathbf{h})) - g(\mathbf{x} + \phi_{k-1}(\mathbf{h})) - \frac{\partial g}{\partial x_k}(\mathbf{x})h_k] \right| \quad (9.5)$$

$$\leq \sum_{k=1}^n \left| g(\mathbf{x} + \phi_k(\mathbf{h})) - g(\mathbf{x} + \phi_{k-1}(\mathbf{h})) - \frac{\partial g}{\partial x_k}(\mathbf{x})h_k \right| \quad (9.6)$$

By the Mean Value Theorem, there exists  $\mathbf{c}_k \in [\mathbf{x} + \phi_{k-1}(\mathbf{h}), \mathbf{x} + \phi_k(\mathbf{h})]$  such that  $g(\mathbf{x} + \phi_k(\mathbf{h})) - g(\mathbf{x} + \phi_{k-1}(\mathbf{h})) = \frac{\partial g}{\partial x_k}(\mathbf{c}_k)h_k$ , and therefore

$$|g(\mathbf{x} + \mathbf{h}) - g(\mathbf{x}) - \mathbf{L}\mathbf{h}| \leq \sum_{k=1}^n \left| \frac{\partial g}{\partial x_k}(\mathbf{c}_k)h_k - \frac{\partial g}{\partial x_k}(\mathbf{x})h_k \right| \quad (9.7)$$

$$= \sum_{k=1}^n \left| \frac{\partial g}{\partial x_k}(\mathbf{c}_k) - \frac{\partial g}{\partial x_k}(\mathbf{x}) \right| |h_k| \quad (9.8)$$

$$< \varepsilon \sum_{k=1}^n |h_k| \quad (9.9)$$

where (9.9) is due to the continuity of partial derivatives. To make life easier, we specify the norm to be  $L_1$  norm, which follows that

$$|g(\mathbf{x} + \mathbf{h}) - g(\mathbf{x}) - \mathbf{L}\mathbf{h}| \leq \varepsilon \|\mathbf{h}\|_1,$$

which implies  $g$  is differentiable at  $\mathbf{x}$ . In particular, if every component  $f_i$  is differentiable, then  $f$  is differentiable.



**Planning for following weeks.** The next lecture will talk about the chain rule, the inverse, the derivative of inverse, the directional derivative, the gradient. Next Friday will talk about Inverse function theorem and Implicit function theorem, which may last for two weeks.

