

**A FIRST COURSE
IN
ANALYSIS**

A FIRST COURSE IN ANALYSIS

MAT2006 Notebook

Lecturer

Prof. Weiming Ni

The Chinese University of Hongkong, Shenzhen

Tex Written By

Mr. Jie Wang

The Chinese University of Hongkong, Shenzhen



香港中文大學(深圳)

The Chinese University of Hong Kong, Shenzhen

Contents

Acknowledgments	vii
Notations	ix
1 Week1	1
1.1 Wednesday	1
1.1.1 Introduction to Set	1
1.2 Quiz	5
1.3 Friday	6
1.3.1 Proof of Schroder-Berstein Theorem	6
1.3.2 Connectedness of Real Numbers	10
2 Week2	13
2.1 Wednesday	13
2.1.1 Review and Announcement	13
2.1.2 Irrational Number Analysis	13
2.2 Friday	21
2.2.1 Set Analysis	21
2.2.2 Set Analysis Meets Sequence	22
2.2.3 Completeness of Real Numbers	23
3 Week3	27
3.1 Tuesday	27
3.1.1 Application of Heine-Borel Theorem	27
3.1.2 Set Structure Analysis	29
3.1.3 Reviewing	31

3.2	Friday	33
3.2.1	Review	33
3.2.2	Continuity Analysis	34
4	Week4	41
4.1	Wednesday	41
4.1.1	Function Analysis	41
4.1.2	Continuity Analysis	46

Acknowledgments

This book is taken notes from the MAT2006 in fall semester, 2018. These lecture notes were taken and compiled in \LaTeX by Jie Wang, an undergraduate student in Fall 2018. Prof. Weiming Ni has not edited this document.

Notations and Conventions

\mathbb{R}^n	n -dimensional real space
\mathbb{C}^n	n -dimensional complex space
$\mathbb{R}^{m \times n}$	set of all $m \times n$ real-valued matrices
$\mathbb{C}^{m \times n}$	set of all $m \times n$ complex-valued matrices
x_i	i th entry of column vector \mathbf{x}
a_{ij}	(i, j) th entry of matrix \mathbf{A}
\mathbf{a}_i	i th column of matrix \mathbf{A}
\mathbf{a}_i^T	i th row of matrix \mathbf{A}
\mathbb{S}^n	set of all $n \times n$ real symmetric matrices, i.e., $\mathbf{A} \in \mathbb{R}^{n \times n}$ and $a_{ij} = a_{ji}$ for all i, j
\mathbb{H}^n	set of all $n \times n$ complex Hermitian matrices, i.e., $\mathbf{A} \in \mathbb{C}^{n \times n}$ and $\bar{a}_{ij} = a_{ji}$ for all i, j
\mathbf{A}^T	transpose of \mathbf{A} , i.e., $\mathbf{B} = \mathbf{A}^T$ means $b_{ji} = a_{ij}$ for all i, j
\mathbf{A}^H	Hermitian transpose of \mathbf{A} , i.e., $\mathbf{B} = \mathbf{A}^H$ means $b_{ji} = \bar{a}_{ij}$ for all i, j
$\text{trace}(\mathbf{A})$	sum of diagonal entries of square matrix \mathbf{A}
$\mathbf{1}$	A vector with all 1 entries
$\mathbf{0}$	either a vector of all zeros, or a matrix of all zeros
\mathbf{e}_i	a unit vector with the nonzero element at the i th entry
$\mathcal{C}(\mathbf{A})$	the column space of \mathbf{A}
$\mathcal{R}(\mathbf{A})$	the row space of \mathbf{A}
$\mathcal{N}(\mathbf{A})$	the null space of \mathbf{A}
$\text{Proj}_{\mathcal{M}}(\mathbf{A})$	the projection of \mathbf{A} onto the set \mathcal{M}

Chapter 4

Week4

4.1. Wednesday

This lecture will talk about the applications of Baire-Category Theorem and continuity analysis.

4.1.1. Function Analysis


In last lecture we have studied that given a analytic function f , if f can be expressed as a partial sum of its series for each x , then f is a polynomial:

Proposition 4.1 Let $f(x) = \sum_{k=0}^{\infty} a_k x^k$ in $|x| < 1$. If for every $x \in (-1, 1)$, there exists $n(= n(x))$ such that $\sum_{k=n+1}^{\infty} a_k x^k = 0$, then f is a polynomial, i.e., n is independent of x .

The idea of the proof is to construct a sequence of points such that f coincide with a polynomial over these points, which implies f is indeed a polynomial.

Now we study its stronger version, i.e., f may not be analytic, it only needs to be infinitely differentiable:

Proposition 4.2 Suppose $f \in C^{\infty}[-1, 1]$. If for every $|x| \leq 1$, there exists $n(= n(x))$ such that $f^{(n)}(x) = 0$, then f is a polynomial.

 Note that an analytic function (i.e., can be expressed as power series) is always infinitely differentiable, but the reverse direction is necessarily not.

For example, recall we have learnt a function

$$f(x) = \begin{cases} \exp\left(-\frac{1}{x^2}\right), & x \neq 0 \\ 0, & x = 0 \end{cases},$$

such that it is infinitely differentiable but $f^{(n)} = 0$ for $n = 1, 2, 3, \dots$. Hence, this function is not analytic at $x = 0$.

Proof. Construct a sequence of set

$$E_n = \{x \in [-1, 1] \mid f^{(n)}(x) = 0\} \implies [-1, 1] = \bigcup_{n=1}^{\infty} E_n,$$

with E_n closed (Exercise #1). Applying Baire-Category Theorem to $[-1, 1]$, at least one E_{N_1} contains a non-empty open interval, say I_1 (Exercise #2).

1. On I_1 , $f^{(N_1)} \equiv 0$, which implies f is a polynomial of degree $N_1 - 1$ (Exercise #3).
2. If $I_1 = (-1, 1)$, the proof is complete.
3. Otherwise, $[-1, 1] \setminus I_1 \neq \emptyset$. Applying Baire-Category Theorem on the set $[-1, 1] \setminus I_1 := \bigcup_{n=1}^{\infty} E_n \setminus I_1$, we conclude that at least one $E_{N_2} \setminus I_1$ contains a non-empty open interval, say I_2 , on which f is a polynomial of degree $N_2 - 1$.
4. Each time applying the same trick to construct I_1, I_2, \dots , and make sure these are the **maximal** intervals with the desired properties. Finally, we reach the stage that:

$f|_{x \in I_j}$ is a polynomial of order $N_j - 1$ for $j = 1, \dots, \infty$ and $\bigcup_{j=1}^{\infty} I_j$ is dense on $[-1, 1]$ (Exercise #4).

5. Construct and claim that

$$H = [-1, 1] \setminus \bigcup_{j=1}^{\infty} I_j = \{-1, 1\}. \text{ (Exercise #5)}$$

6. Combining (4) and (5), we derive f satisfies the condition in Proposition(4.2), and therefore is a polynomial. (Exercise #6)

■

Verification. Here we give some hints for the exercises above:

1. Since the inverse image of $\{0\}$ is closed for continuous functions, and $f^{(n)}(\cdot)$ is continuous, we derive E_n 's are closed.
2. The Baire-Category Theorem asserts that for a non-empty complete metric space X , or any subsets of X with **non-empty** interior, if it is the countably union of **closed** sets, then one of these **closed** sets has non-empty interior.
3. By integrating $f^{(N_1)}$ for N_1 times, e.g.,

$$f^{(N_1)} = 0 \implies f^{(N_1-1)} = \int f^{(N_1)} dx = a_0 \implies \dots \implies f = a_{N_1-1}x^{N_1-1} + \dots + a_0$$

4. Let $I \subseteq [-1, 1]$ be any open interval. Its closure can be expressed as:

$$\bar{I} = \bigcup_{n=1}^{\infty} \bar{I} \cap E_n$$

Applying Baire category theorem to \bar{I} , at least one $\bar{I} \cap E_{n'}$ contains an open interval I' . Thus, $I' \subseteq I$ and $I' \subseteq E_{n'}$, which implies $I' \in \bigcup_{j=1}^{\infty} I_j$ (recall that I_j 's are picked maximally). This means that $\bigcup_{j=1}^{\infty} I_j \cap I$ is non-empty for arbitrary open interval I , which implies $\bigcup_{j=1}^{\infty} I_j$ is dense.

5. We have seen that $\bigcup_{j=1}^{\infty} I_j$ is an open, dense **proper** subset of $[0, 1]$, which means H is **non-empty**, closed, and nowhere dense in $[0, 1]$. In order to show $H = \{-1, 1\}$, it suffices to show H does not contain open intervals. Otherwise applying Baire-Category Theorem to $H = \bigcup_{n=1}^{\infty} E_n \cap H$ again, for some fixed n^* , $E_{n^*} \cap H$ contains an open interval I^* . Thus, $I^* \subseteq E_{n^*}$ and $I^* \subseteq H \implies I^* \subseteq (\bigcup_{j=1}^{\infty} I_j)^c$, which leads to a contradiction as I_j 's are picked maximally.
6. Hence, $\bigcup_{j=1}^{\infty} I_j = (-1, 1)$, i.e., $f(x) = \sum_{k=0}^{\infty} a_k x^k$ in $|x| < 1$.

A simpler and more clear proof is presented in the website

<https://mathoverflow.net/questions/34059/>

if-f-is-infinitely-differentiable-then-f-coincides-with-a-polynomial

We have seen some examples of nowhere differentiable functions. Now we show that almost functions are nowhere differentiable.

Notations. We denote $\mathcal{C}[0,1]$ as the set of all continuous functions on $[0,1]$. One corresponding metric is defined as:

$$d(f,g) = \sup_{x \in [0,1]} |f(x) - g(x)|, \quad \forall f, g \in \mathcal{C}[0,1].$$

Remember that $(\mathcal{C}[0,1], d)$ is complete.

Theorem 4.1 The set of all nowhere differentiable functions in $(\mathcal{C}[0,1], d)$ is dense, i.e., forms a 2nd Category.

The trick is to show the complement of the set of nowhere differentiable functions, i.e., the set of functions that have a **finite** derivative at some point, forms a 1st Category.

Proof. Construct

$$E_n = \left\{ f \in \mathcal{C}[0,1] \left| \begin{array}{l} \forall 0 < h < 1 - x, \left| \frac{f(x+h) - f(x)}{h} \right| \leq n, \\ \text{for some } 0 \leq x \leq 1 - \frac{1}{n} \end{array} \right. \right\}$$

Thus the union of all E_n will contain all functions having a finite **right hand derivative** at some point in $[0,1)$.

Proposition 4.3 E_n is closed, i.e., for a sequence of function $\{f_m\} \subseteq E_n$ such that $f_m \rightarrow f$, we have $f \in E_n$.

Proposition 4.4 E_n is nowhere dense, i.e., $(\mathcal{C} \setminus E_n)$ is dense):

After showing these two propositions, we conclude that the set of functions, with a right derivatives at some point, is a set of the first category. Similarly, we can repeat these steps for left derivatives. In summary, the set of functions with a well-defined derivatives forms a 1st Category. The proof is complete. ■

Proof of Proposition(4.3). Since $\{f_m\} \subseteq E_n$, there exists a sequence of $\{x_m\}$ such that for

each m ,

$$0 \leq x_m \leq 1 - \frac{1}{n}$$

$$|f_m(x_m + h) - f_m(x_m)| \leq hn,$$

for $\forall 0 < h < 1 - x_m$. As $\{x_m\}$ is bounded, there exists a subsequence $\{x_{m,k}\}$ of $\{x_m\}$ with limit $x \in [0, 1 - \frac{1}{n}]$.

For $\forall 0 < h < 1 - x$, we have that $0 < h < 1 - x_{m,k}$ for large k . Applying triangle inequality, we obtain:

$$\begin{aligned} |f(x+h) - f(x)| &\leq |f(x+h) - f(x_{m,k}+h)| + |f(x_{m,k}+h) - f_m(x_{m,k}+h)| \\ &\quad + |f_m(x_{m,k}+h) - f_m(x_{m,k})| + |f_m(x_{m,k}) - f(x_{m,k})| + |f(x_{m,k}) - f(x)| \\ &\leq |f(x+h) - f(x_{m,k}+h)| + d(f, f_k) + nh + d(f_k, f) + |f(x_k) - f(x)|. \end{aligned}$$

Taking $k \rightarrow \infty$, we find all terms in RHS goes to zero except nh :

$$|f(x+h) - f(x)| \leq nh \implies f \in E_n.$$

■

Proof of Proposition(4.4). In order to show E_n is nowhere dense, by using the fact that E_n is closed, it suffices to show that an arbitrary open neighborhood $B(f, \varepsilon)$ will contain elements from the set $\mathcal{C}[0,1] \setminus E_n$, i.e., it suffices to create a function in $B(f, \varepsilon)$ that cannot be in E_n for fixed ε .

- Construct a piecewise linear function $\phi_N(x)$ on $[0,1]$ first:

$$\phi_N(x) = \begin{cases} N(x - \frac{k}{N}), & \frac{k}{N} \leq x \leq \frac{k+1}{N}, k = 0, 2, \dots, N \\ -N(x + \frac{k}{N}), & \frac{k}{N} \leq x \leq \frac{k+1}{N}, k = 1, 3, \dots, N-1 \end{cases}$$




Figure 4.1: Plot of function $\phi_N(x)$

As we can see, N is the maximum slope of the piecewise linear function ϕ_N .

- Let M be the maximum slope of the piecewise linear function f , and pick a positive even integer m such that

$$\frac{1}{2}mN\varepsilon > M + n.$$

Then we construct function

$$g(x) = f(x) + \frac{1}{2}\varepsilon\phi_{mN}(x)$$

As we can see, $d(f, g) = \frac{1}{2}\varepsilon < \varepsilon$, thus $g \in B(f, \varepsilon)$. Also note that

$$\begin{aligned} \left| \frac{g(x+h) - g(x)}{h} \right| &\geq \left| \frac{1/2\varepsilon(\phi_{mN}(x+h) - \phi_{mN}(x))}{h} \right| - \left| \frac{f(x+h) - f(x)}{h} \right| \\ &\geq \frac{1}{2}\varepsilon \left| \frac{(\phi_{mN}(x+h) - \phi_{mN}(x))}{h} \right| - M \\ &= \frac{1}{2}mN\varepsilon - M > n \end{aligned}$$

for x in $(0, 1 - \frac{1}{mN})$ and some $h \in (0, 1 - x)$. Hence, $g \notin E_n$. The proof is complete. ■

4.1.2. Continuity Analysis

Recall the definition for continuity:

- A function f is said to be continuous at $x_0 \in I$ if $\forall \varepsilon > 0$, there exists $\delta > 0$ (δ depends on x_0 and ε) such that

$$|f(x) - f(x_0)| < \varepsilon, \quad \forall |x - x_0| < \delta$$

- A function f is continuous on I if it is continuous at every point in I .

Definition 4.1 [Uniform] We say f is **uniformly continuous** on I if $\forall \varepsilon > 0$, there exists δ (depend only on ε , but independent of $x \in I$) such that

$$|f(y) - f(x)| < \varepsilon, \text{ if } |x - y| < \delta$$

- R** It is useful to note that the uniform continuity places a upper bound on the growth of the function at every point, i.e., the function cannot grow too fast.

■ **Example 4.1** Given a function $f(x) = x^2$,

1. Is it uniformly continuous on $[0, 1]$?

Yes, intuitively the growth of x^2 is limited within bounded interval.

2. Is it uniformly continuous on \mathbb{R} ?

No, intuitively the growth of x^2 tends to infinite as $x \rightarrow \infty$.

Proof: For fixed x , if $|y - x| < \delta$, if we choose $|x| \geq \frac{\varepsilon}{2\delta} + \frac{\delta}{2}$, then

$$\begin{aligned} \underbrace{|f(y) - f(x)|}_{\varepsilon} &= |y^2 - x^2| = |y + x| \underbrace{|y - x|}_{\delta} \\ &\geq (|2x| - |x - y|)|y - x| \geq \left(\frac{\varepsilon}{\delta} + \delta - \delta\right)\delta = \varepsilon \end{aligned}$$

which is a contradiction.




Figure 4.2: The proof and application for the Theorem(4.2) is Mandatory. If you don't know how to do it in the exam, Prof.Ni will fail you without hesitation.

Theorem 4.2 Suppose that f is continuous on a compact set D . Then f is uniformly continuous on D .

Proof. For given $\varepsilon > 0$, since f is continuous at x , there exists $\delta_x > 0$ s.t.

$$|f(y) - f(x)| < \frac{\varepsilon}{2}, \text{ if } |y - x| < \delta_x.$$

Construct an open cover $\{B_{\delta_x}(x) \mid x \in D\}$ of D with

$$B_{\delta_x}(x) = \{y \in D \mid |y - x| < \frac{1}{2}\delta_x\}.$$

The set D is compact implies there exists a finite subcover:

$$D \subseteq B_{\delta_{x_1}}(x_1) \cup B_{\delta_{x_2}}(x_2) \cup \dots \cup B_{\delta_{x_k}}(x_k). \quad (4.1)$$

Construct $\delta > 0$ such that $B_\delta(x)$ must be contained entirely in one of the ball, say $B_{\delta_{x_j}}(x_j)$ (Exercise #7)

Therefore given $|y - x| < \delta$ we imply $x, y \in B_{\delta_{x_j}}(x_j)$ for some j , which follows that

$$|f(y) - f(x)| \leq |f(y) - f(x_j)| + |f(x_j) - f(x)| < \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon$$

■

Verification of Exercise. Such a δ is constructed as

$$\delta = \frac{1}{2} \min\{\delta_{x_1}, \delta_{x_2}, \dots, \delta_{x_k}\}.$$

Thus for any x, y with $|y - x| < \delta$, by (4.1), there exists j such that $x \in B_{\delta_{x_j}}(x_j)$, and hence

$$|x - x_j| < \frac{1}{2} \delta_{x_j} \quad (4.2)$$

Also, we have

$$|y - x_j| \leq |y - x| + |x - x_j| \leq \delta + \frac{1}{2} \delta_{x_j} \leq \delta, \quad (4.3)$$

i.e., y is also in $B_{\delta_{x_j}}(x_j)$.

Definition 4.2 [Convex] A real-valued function f defined in (a, b) is said to be convex if

$$f(tx + (1 - t)y) \leq tf(x) + (1 - t)f(y)$$

whenever $a < x < b, a < y < b, 0 < t < 1$. ■

Check Rudin's book for the proof that a convex function is always continuous.

