

**A FIRST COURSE  
IN  
ANALYSIS**



---

# A FIRST COURSE IN ANALYSIS

## MAT2006 Notebook

---

### Lecturer

*Prof. Weiming Ni*

*The Chinese University of Hongkong, Shenzhen*

### Tex Written By

*Mr. Jie Wang*

*The Chinese University of Hongkong, Shenzhen*



香港中文大學(深圳)

The Chinese University of Hong Kong, Shenzhen



# Contents

Acknowledgments	vii
Notations	ix
<b>1 Week1</b>	<b>1</b>
1.1 Wednesday	1
1.1.1 Introduction to Set	1
1.2 Quiz	5
1.3 Friday	6
1.3.1 Proof of Schroder-Berstein Theorem	6
1.3.2 Connectedness of Real Numbers	10
<b>2 Week2</b>	<b>13</b>
2.1 Wednesday	13
2.1.1 Review and Announcement	13
2.1.2 Irrational Number Analysis	13
2.2 Friday	21
2.2.1 Set Analysis	21
2.2.2 Set Analysis Meets Sequence	22
2.2.3 Completeness of Real Numbers	23
<b>3 Week3</b>	<b>27</b>
3.1 Tuesday	27
3.1.1 Application of Heine-Borel Theorem	27
3.1.2 Set Structure Analysis	29
3.1.3 Reviewing	31

<b>3.2</b>	<b>Friday</b>	<b>33</b>
3.2.1	Review . . . . .	33
3.2.2	Continuity Analysis . . . . .	34
<b>4</b>	<b>Week4 . . . . .</b>	<b>41</b>
<b>4.1</b>	<b>Wednesday</b>	<b>41</b>
4.1.1	Function Analysis . . . . .	41
4.1.2	Continuity Analysis . . . . .	46

# Acknowledgments

This book is taken notes from the MAT2006 in fall semester, 2018. These lecture notes were taken and compiled in  $\text{\LaTeX}$  by Jie Wang, an undergraduate student in Fall 2018. Prof. Weiming Ni has not edited this document.





# Notations and Conventions

$\mathbb{R}^n$	$n$ -dimensional real space
$\mathbb{C}^n$	$n$ -dimensional complex space
$\mathbb{R}^{m \times n}$	set of all $m \times n$ real-valued matrices
$\mathbb{C}^{m \times n}$	set of all $m \times n$ complex-valued matrices
$x_i$	$i$ th entry of column vector $\mathbf{x}$
$a_{ij}$	$(i, j)$ th entry of matrix $\mathbf{A}$
$\mathbf{a}_i$	$i$ th column of matrix $\mathbf{A}$
$\mathbf{a}_i^T$	$i$ th row of matrix $\mathbf{A}$
$\mathbb{S}^n$	set of all $n \times n$ real symmetric matrices, i.e., $\mathbf{A} \in \mathbb{R}^{n \times n}$ and $a_{ij} = a_{ji}$ for all $i, j$
$\mathbb{H}^n$	set of all $n \times n$ complex Hermitian matrices, i.e., $\mathbf{A} \in \mathbb{C}^{n \times n}$ and $\bar{a}_{ij} = a_{ji}$ for all $i, j$
$\mathbf{A}^T$	transpose of $\mathbf{A}$ , i.e, $\mathbf{B} = \mathbf{A}^T$ means $b_{ji} = a_{ij}$ for all $i, j$
$\mathbf{A}^H$	Hermitian transpose of $\mathbf{A}$ , i.e, $\mathbf{B} = \mathbf{A}^H$ means $b_{ji} = \bar{a}_{ij}$ for all $i, j$
$\text{trace}(\mathbf{A})$	sum of diagonal entries of square matrix $\mathbf{A}$
$\mathbf{1}$	A vector with all 1 entries
$\mathbf{0}$	either a vector of all zeros, or a matrix of all zeros
$\mathbf{e}_i$	a unit vector with the nonzero element at the $i$ th entry
$\mathcal{C}(\mathbf{A})$	the column space of $\mathbf{A}$
$\mathcal{R}(\mathbf{A})$	the row space of $\mathbf{A}$
$\mathcal{N}(\mathbf{A})$	the null space of $\mathbf{A}$
$\text{Proj}_{\mathcal{M}}(\mathbf{A})$	the projection of $\mathbf{A}$ onto the set $\mathcal{M}$



# Chapter 1

## Week1

### 1.1. Wednesday

#### Recommended Reading.

1. (Springer-Lehrbuch) V. A. Zorich, J. Schüle-Analysis I-Springer (2006).
2. (The Carus mathematical monographs 13) Ralph P. Boas, Harold P. Boas, A primer of real functions-Mathematical Association of America (1996).
3. (International series in pure and applied mathematics) Walter Rudin, Principles of Mathematical Analysis-McGraw-Hill (1976).
4. Terence Tao, Analysis I,II-Hindustan Book Agency (2006)
5. (Cornerstones) Anthony W. Knap, Basic real analysis-Birkhäuser (2005)

#### 1.1.1. Introduction to Set

For a set  $\mathcal{A} = \{1,2,3\}$ , we have  $2^3 = 8$  subsets of  $\mathcal{A}$ . We are interested to study the collection of sets.

**Definition 1.1** [Collection of Subsets] Given a set  $\mathcal{A}$ , the the collection of subsets of  $\mathcal{A}$  is denoted as  $2^{\mathcal{A}}$ . ■

We use Cardinal to describe the order of number of elements in a set.

**Definition 1.2** Given two sets  $\mathcal{A}$  and  $\mathcal{B}$ ,  $\mathcal{A}$  and  $\mathcal{B}$  are said to be **equivalent** (or have the same cardinal) if there exists a 1-1 onto mapping from  $\mathcal{A}$  to  $\mathcal{B}$ . ■

**Definition 1.3** [Countability] The set  $\mathcal{A}$  is said to be **countable** if  $\mathcal{A} \sim \mathbb{N} = \{1, 2, 3, \dots\}$ ; an infinite set  $\mathcal{A}$  is **uncountable** if it is not equivalent to  $\mathbb{N}$ . ■

**R** Note that the set of integers, i.e.,  $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$  is also countable; the set of rational numbers, i.e.,  $\mathbb{Q} = \{p/q \mid p, q \in \mathbb{Z}, q \neq 0\}$  is countable.

We skip the process to define real numbers.

**Proposition 1.1** The set of real numbers  $\mathbb{R}$  is **uncountable**.

For example,  $\sqrt{2} \notin \mathbb{Q}$ . Some irrational numbers are the roots of some polynomials, such a number is called **algebraic** numbers. However, some irrational numbers are not, such a number is called **transcendental**. For example,  $\pi$  is **not** algebraic. We will show that the collection of algebraic numbers are countable in the future.

There are two steps for the proof for proposition(1.1):

*Proof.* 1.  $2^{\mathbb{N}}$  is **uncountable**:

Assume  $2^{\mathbb{N}}$  is countable, i.e.,

$$2^{\mathbb{N}} = \{A_1, A_2, \dots, A_k, \dots\}$$

Define  $B := \{k \in \mathbb{N} \mid k \notin A_k\}$ , it is a collection of subscripts such that the subscript  $k$  does not belong to the corresponding subsets  $A_k$ .

It follows that  $B \in 2^{\mathbb{N}} \implies B = A_n$  for some  $n$ . Then it follows two cases:

- If  $n \in A_n$ , then  $n \notin B = A_n$ , which is a contradiction
- Otherwise,  $n \in B = A_n$ , which is also a contradiction.

The proof for the claim  $2^{\mathbb{N}}$  is **uncountable** is complete.

2.  $\mathbb{R} \sim 2^{\mathbb{N}}$ :

**Firstly we have**  $\mathbb{R} \sim (0, 1)$ . This can be shown by constructing a one-to-one mapping:

$$f : \mathbb{R} \mapsto (0, 1) \quad f(x) = \frac{1}{\pi} \arctan x + \frac{1}{2}, \forall x \in \mathbb{R}$$

Secondly, we show that  $2^{\mathbb{N}} \sim (0,1)$ . We construct a mapping  $f$  such that

$$f : 2^{\mathbb{N}} \mapsto (0,1),$$

where for  $\forall A \in 2^{\mathbb{N}}$ ,

$$f(A) = 0.a_1a_2a_3\dots, \quad a_j = \begin{cases} 2, & \text{if } j \in A \\ 4, & \text{if } j \notin A \end{cases}$$

This function is only 1-1 mapping but not onto mapping.


Reversely, we construct a 1-1 mapping from  $(0,1)$  to  $2^{\mathbb{N}}$ . We construct a mapping  $g$  such that

$$g : (0,1) \mapsto 2^{\mathbb{N}}$$

where for any real number from  $(0,1)$ , we can write it into binary expansion:

binary form:  $0.a_1a_2\dots$  where  $a_j = 0$  or  $1$ .

Hence, we construct  $g(0.a_1a_2\dots) = \{j \in \mathbb{N} \mid a_j = 0\} \subseteq \mathbb{N}$ , which implies  $g(\cdot) \in 2^{\mathbb{N}}$ .

 Our intuition is that two 1-1 mappings in the reverse direction will lead to a 1-1 **onto** mapping. If this is true, then we complete the proof. This intuition is the **Schroder-Berstein Theorem**.

■

**Defining Binary Form.** However, during this proof, we must be careful about the binary form of a real number from  $(0,1)$ . Now we give a clear definition of Binary Form:

For a real number  $a$ , to construct its binary form, we define

$$a_1 = \begin{cases} 0, & \text{if } a \in (0, \frac{1}{2}) \\ 1, & \text{if } a \in [\frac{1}{2}, 1). \end{cases}$$

After having chosen  $a_1, a_2, \dots, a_{j-1}$ , we define  $a_j$  to be the largest integer such that

$$\frac{1}{2}a_1 + \frac{1}{2^2}a_2 + \dots + \frac{a_j}{2^j} \leq a$$

Then the binary form of  $a$  is  $a := 0.a_1a_2\dots$

**Theorem 1.1 — Schroder-Berstein Theorem.** If  $f : A \mapsto B$  and  $g : B \mapsto A$  are both 1-1 mapping, then there exists a 1-1 onto mapping from  $A$  to  $B$ , i.e.,  $\text{card } A$  equals to  $\text{card } B$ .

Exercise: Show that  $(0,1]$  and  $[0,1)$  have 1-1 onto mapping without applying Schroder-Berstein Theorem.

The next lecture we will take a deeper study into the proof of Schroder-Berstein Theorem and the real number.

