

# Contents

<b>1</b>	<b>Permutations and Combinations</b>	<b>2</b>
1.1	Basic Counting Principles . . . . .	2
1.2	Permutations . . . . .	4

# Permutations and Combinations

## 1.1 Basic Counting Principles

An important motivation to study combinatorics is to count the **number of ways** in which an event may occur. Intuitively, we have two approaches to count.

The first approach is to categorise the event into **non-overlapping cases**. This means that we break an event into mutually exclusive sub-events, after which we can count the number of ways for each sub-event to occur. The aggregate of these counts is the total number of ways for the original event to occur.

Those familiar with basic set theory may consider  $E$  to be the set containing all distinct ways for an event to occur. By breaking up the event, we essentially establish a **partition** of  $E$ , so that the sum of cardinalities of all the elements in that partition equals the cardinality of  $E$ .

This motivates us to write the following principle using set notations.

### Theorem 1.1.1 ► Addition Principle (AP)

Let  $k \in \mathbb{N}^+$  and let  $A_1, A_2, \dots, A_k$  be  $k$  finite sets which are pairwise disjoint, i.e.  $A_i \cap A_j = \emptyset$  whenever  $i \neq j$ , then

$$\left| \bigcup_{i=1}^k A_i \right| = \sum_{i=1}^k |A_i|.$$

*Proof.* The case where  $k = 1$  is trivial.

Suppose that when  $k = n$ , we have

$$\left| \bigcup_{i=1}^n A_i \right| = \sum_{i=1}^n |A_i|$$

for any  $n$  finite sets which are pairwise disjoint. Let  $A_{n+1}$  be an arbitrary finite set

which is disjoint with any of the  $A_i$ 's from the  $n$  sets. So we have:

$$\begin{aligned}
 \left| \bigcup_{i=1}^{n+1} A_i \right| &= \left| \left( \bigcup_{i=1}^n A_i \right) \cup A_{n+1} \right| \\
 &= \left| \bigcup_{i=1}^n A_i \right| + |A_{n+1}| - \left| \left( \bigcup_{i=1}^n A_i \right) \cap A_{n+1} \right| \\
 &= \left( \sum_{i=1}^n |A_i| \right) + |A_{n+1}| - |\emptyset| \\
 &= \sum_{i=1}^{n+1} |A_i|.
 \end{aligned}$$

Therefore, the original statement holds for all  $k \in \mathbb{N}^+$ . □

In more casual language, this means that if an event  $E_k$  has  $n_k$  distinct ways to occur, then there is  $\sum_{i=1}^k n_k$  ways for at least one of the events  $E_1, E_2, \dots, E_k$  to occur, provided that  $E_i$  and  $E_j$  can never occur concurrently whenever  $i \neq j$ .

Given an event  $E$ , the other approach to count the number of ways for it to occur is to break  $E$  up internally into **non-overlapping stages**.

With set notations, we can write the  $i$ -th stage for  $E$  to occur as  $e_i$ , and so a way for  $E$  to occur can be represented by an ordered tuple  $(e_1, e_2, \dots, e_k)$ , where  $k$  is the total number of stages to undergo for  $E$  to occur.

Let  $E_i$  denote the set of all distinct ways to undergo the  $i$ -th stage of  $E$ , then it is easy to see that  $E$  is just the **Cartesian product** of all the  $E_i$ 's. Hence, we derive the following principle:

### Theorem 1.1.2 ► Multiplication Principle (MP)

Let  $k \in \mathbb{N}^+$  and let  $A_1, A_2, \dots, A_k$  be  $k$  pairwise disjoint finite sets, then

$$\left| \prod_{i=1}^k A_i \right| = \prod_{i=1}^k |A_i|.$$

*Proof.* The case where  $k = 1$  is trivial.

Suppose that when  $k = n$ , we have

$$\left| \prod_{i=1}^n A_i \right| = \prod_{i=1}^n |A_i|$$

for any  $n$  finite sets which are pairwise disjoint. Let  $A_{n+1}$  be an arbitrary finite set which is disjoint with any of the  $A_i$ 's from the  $n$  sets. Take  $a_i, a_j \in A_{n+1}$ . Note that for all  $\mathbf{a} \in \prod_{i=1}^n A_i$ ,  $(\mathbf{a}, a_i) \neq (\mathbf{a}, a_j)$  whenever  $a_i \neq a_j$ . This means that

$$\begin{aligned} \left| \prod_{i=1}^{n+1} A_i \right| &= \left| \prod_{i=1}^n A_i \times A_{n+1} \right| \\ &= \left| \prod_{i=1}^n A_i \right| |A_{n+1}| \\ &= \left( \prod_{i=1}^n |A_i| \right) |A_{n+1}| \\ &= \prod_{i=1}^{n+1} |A_i| \end{aligned}$$

Therefore, the original statement holds for all  $k \in \mathbb{N}^+$ . □

In more casual language, this means that if an event  $E$  requires  $k$  stages to be undergone before it occurs and the  $i$ -th stage has  $n_i$  ways to complete, then there is  $\prod_{i=1}^k n_i$  ways for  $E$  to occur, provided that no two different stages complete concurrently.

## 1.2 Permutations

A fundamental problem in combinatorics is described as follows: given a set  $S$ , how many ways are there to arrange  $r$  elements in  $S$ , i.e. how many **distinct sequences** can be formed using the elements in  $S$  without repetition? The process of selecting elements from  $S$  and arranging them as a sequence is known as **permutation**.

Note that forming a sequence using  $r$  elements from a set  $S$  is an event consisting of  $r$  stages, as we need to select an element for each of the  $r$  terms of the sequence. Suppose  $S$  has  $n$  elements. For the first term of the sequence, we can choose any of the elements in  $S$ , so there is  $n$  ways to do it. For the second term, since we cannot repeat the elements, we are left with  $n - 1$  choices.

Continue choosing elements in this way, we realise that if we choose the terms sequentially, when we reach the  $k$ -th term we will be left with  $n - k + 1$  options as the previous  $(k - 1)$  terms have taken away  $(k - 1)$  elements. By Theorem 1.1.2, we know that the number of sequences which can be formed is given by  $\prod_{i=1}^r (n - r + i)$ .

**Definition 1.2.1 ► Permutations**

Let  $A$  be a finite set such that  $|A| = n$ , an  $r$ -permutation of  $A$  is a way to arrange  $r$  elements of  $A$ , denoted as  $P_r^n$  and given by

$$P_r^n = \prod_{i=1}^r (n - r + i) = \frac{n!}{(n - r)!}.$$