

- **Co-finite:** open $\iff$ complements of finite sets.
- **Co-countable:** open $\iff$ complements of countable sets.
- $\mathcal{B}$ gen. the coarsest containing itself.
- **Metric:** ≥ 0 , **definite**, **symmetric**, Δ -**inequality**.
- **Pseudo-metric:** $d(x, x) = 0$ but can be not definite.
- **Quasi-metric:** can be not symmetric.
- **Norm:** ≥ 0 , **definite**, Δ -**inequality**, $\|\lambda x\| = |\lambda| \|x\|$.
- **Distance between sets:** smallest pointwise distance.
- **Diameter of set:** sup of pointwise distance.
- L^p -**metric:** gen. the standard topo. on $\mathbb{R}^n$.

$$\max \|y_i - x_i\| \leq \left[\sum_{i=1}^n \|y_i - x_i\|^p \right]^{\frac{1}{p}} \leq n^{\frac{1}{p}} \max \|y_i - x_i\|.$$

- Metrics are equivalent $\iff c_1 d \leq d' \leq c_2 d$.
- $x \in \overline{A} \iff \forall$ open neighbourhood U of x , $U \cap A \neq \emptyset$.
- **Continuity:** $f^{-1}(S)$ is open $\forall S$ in sub-basis.
- TFAE:
 1. f is cont.;
 2. for all $A \subseteq X$, $f(\overline{A}) \subseteq \overline{f(A)}$;
 3. for any closed set $B \subseteq Y$, $f^{-1}(B)$ is closed in X ;
 4. $\forall x \in X$ and any open $V \subseteq Y$ with $f(x) \in V$, there is an open set $U \subseteq X$ s.t. $x \in U$ and $f(U) \subseteq V$.
- **Dense:** $\forall$ open U containing $x \in X$, $U \cap A \neq \emptyset$.
- Dense $\iff X \setminus A = \emptyset \iff \overline{A} = X$.
- **Pasting lemma:** if $X = A \cup B$ for closed A, B and $f = g$ for all $x \in A \cap B$, then $h = x \in A? f : (x \in B? g)$ is cont. if $f : A \rightarrow Y$ and $g : B \rightarrow Y$ are.
- **Pull-back** topo.: $\{f^{-1}(U) : U \text{ is open}\}$ is the coarsest topo. ensuring cont. f . (quotient)
- **Uniform continuity:** $\forall \epsilon > 0$, there exists some $\delta > 0$ s.t. $d_X(x, y) < \delta \implies d_Y(f(x), f(y)) < \epsilon$.
- f is uniformly cont. $\iff \forall \{x_i\}_i^\infty, \{y_i\}_i^\infty$ s.t. $\lim_{i \rightarrow \infty} d_X(x_i, y_i) = 0$, $\lim_{i \rightarrow \infty} d_Y(f(x_i), f(y_i)) = 0$.

- Limit of uniformly convergent sequence is cont..
- $x \mapsto x^n$ converges to $\mathbf{1}_{x=1}$ pointwise but not uniformly.
- **Prod. topo.:** $\mathcal{S} := \{\pi_{X_\alpha}^{-1}(U_\alpha) : \alpha \in \Lambda, U_\alpha \in \mathcal{T}_\alpha\}$.
- Prod. topo. is the coarsest for cont. π_α 's.
- $x \mapsto (x)^\mathbb{N}$ is not cont. in the box topo..
- f is cont. iff all components are cont..
- Subspace topo. of prod. topo. equals prod. topo. of subspace topologies.
- Standard topo. on $\mathbb{R}^n$ is the prod. topo. by standard topologies on $\mathbb{R}^{m_i}$'s.
- Prod. space basic open set: all but finitely many factors are not X_i .
- Box topo. is finer: e.g. $\prod_{i \in \mathbb{N}} U_i$.
- $d_1 = \sum_{i=1}^n d_{X_i}(x_i, y_i)$ and $d_\infty = \max d_{X_i}(x_i, y_i)$ both induce the prod. topo..
- $\rho(x, y) := \frac{d(x, y)}{1 + d(x, y)}$ is a metric with $\text{diam}(X) < 1$.
- $D(x, y) := \sup \left\{ \frac{\rho_i(x_i, y_i)}{i} : i \in \mathbb{Z}^+ \right\}$ is a metric inducing the infinite prod. topo..
- **Quotient map:** surj and U open iff $f^{-1}(U)$ open.
- Closed not open: $x \mapsto x \in [0, 1]? x : (x \in [2, 3]? x - 1)$.
- Open not closed: $x \mapsto x \in (0, 1)? x : (x \in (2, 3)? x - 1)$.
- Neither: $X := ([0, \infty) \times \mathbb{R}) \cup (\mathbb{R} \times \{0\})$, $(x, y) \mapsto x$. **But this is a quotient map!**
- Both: id_X .
- **Open map:** cont. map from open set to open set.
- Surjective cont. + open or closed = quotient map.
- Quotient, open and closed maps are preserved under $\circ$.
- **Saturated:** $A = f^{-1}(f(A))$ or $A = f^{-1}(S)$.
- Surjective cont. f is a quotient map iff $f(A)$ is open (closed) whenever A is saturated open (closed).
- Restriction to saturated sets preserves quotient map.
- Homeomorphism: bijective open map.
- Embedding: injective homeomorphism onto $f(X)$.

- Isometric: $d_Y(y_a, y_b) = d_X(a, b)$. Isometry if surj.

Countability

- 2nd countable $\implies \exists$ dense countable subset. $\iff$ is true if X is metrisable.
- $(\mathbb{R}, \mathcal{P}(\mathbb{R}))$ is 1st but not 2nd countable.
- Uncountable co-finite have no countable basis.
- $\exists$ nested countable basis $B_1 \subseteq B_2 \subseteq \dots$.
- $x_n \rightarrow x \implies x \in \overline{X}$. $\iff$ is true if X is 1st countable.
- If f is cont., then for any sequence $f(x_i) \rightarrow f(x)$. The converse is true if X is first countable.
- Lindelof: every open cover has a countable sub-cover.
- 2nd countable $\implies$ Lindelof. $\iff$ if X is metrisable.

Separation

- Co-finite topo. is T_1 and T_2 if X is finite.
- $T_1 \iff \{x\}$ is closed $\forall x \in X$.
- **Embedding thm:** $\forall x$ and $\forall$ open U s.t. $x \in U$, $\exists \alpha \in \Lambda$ s.t. $f_\alpha(x) > 0$ and $f_\alpha(X \setminus U) = \{0\}$, then $F(x) = (f_\alpha(x))_{\alpha \in \Lambda}$ is embedding.
- Metric spaces are T_2 so all finite subsets are closed.
- T_2 iff diagonal is closed.
- $T_3 \iff \forall x$ and $\forall$ open U s.t. $x \in U$, $\exists$ open V s.t. $x \in V$ and $\overline{V} \subseteq U$.
- $T_4 \iff \forall$ closed A and open U s.t. $A \subseteq U$, $\exists$ open V s.t. $A \subseteq V$ and $\overline{V} \subseteq U$.
- $T_4 \implies T_3 \implies T_2 \implies T_1$.
- Every metric space is T_4 .
- Every 2nd countable T_3 space is T_4 .
- Cont. f separates A, B : $f(A) = \{0\}$, $f(B) = \{1\}$.
- **Urysohn's lem:** Every T_4 space is completely normal.
- **Urysohn's metrisation thm:** every 2nd countable T_3 space is metrisable.

Compactness

- $[-\sqrt{2}, \sqrt{2}] \cap \mathbb{Q}$: closed and bounded, not cpt.
- Closed subset of cpt space is cpt.
- Cpt subset of T_2 space is closed.

- Set in co-finite space is cpt but closed iff it's finite.
- Closure of particular point is not cpt.
- $\text{Cpt} \implies (T_4 \iff T_3 \iff T_2)$
- Cont. f maps cpt set to cpt set.
- **Tube lemma:** If Y is cpt and $N \subseteq X \times Y$ is open and contains $\{x_0\} \times Y$, then $\exists W \supseteq \{x_0\}$ open s.t. $W \times Y \subseteq N$.
- Y is cpt $T_2 \implies \Gamma_f$ closed $\iff f$ is cont..
- X cpt iff $\forall$ set of closed sets $\mathcal{G}$ with f.i.p, $\bigcap \mathcal{G} \neq \emptyset$.
- x is **isolated** $\iff \{x\}$ is open.
- If $U \neq \emptyset$ is open in a T_2 space and x is not isolated, then $\exists$ non-empty open $V \subseteq U$ s.t. $x \notin \overline{V}$.
- Non-empty T_2 space $\approx \mathbb{R}$ if has no isolated point.
- **Tychonoff's thm:** prod. of cpt is cpt.
- **Lim pt cpt:** every infinite $Y \subseteq X$ has a lim pt in X . $\text{Cpt} \implies \text{lim pt cpt}$.
- Let Y be trivial and $\mathbb{Z}^+$ be discrete, then $Y \times \mathbb{Z}^+$ is lim pt cpt but not cpt.
- **Seq cpt:** every sequence has a convergent subsequence. $\text{Seq cpt} \implies \text{lim pt cpt}$.
- Metric space: cpt iff lim pt cpt iff seq cpt
- $\mathbb{R}$ with basis (a, ∞) 's is seq cpt but not lim pt cpt.
- **Lebesgue number:** $\forall S \subseteq X$ with $\text{diam}(S) < \delta$, $\exists U \in \mathcal{U}$ s.t. $S \subseteq U$.
- Every open cover of seq cpt metric space has a Lebesgue number.
- **Totally bounded:** $\forall \epsilon > 0$, $\exists$ finite cover by $B_\epsilon(x_i)$.
- Every seq cpt metric space is totally bounded.
- In a metric space, cpt iff lim pt cpt iff seq cpt.
- $f: X \rightarrow Y$ is cont. between metric spaces. If X is cpt, then f is uniformly cont..
- Totally bounded $\implies \text{diam}(X) < \infty$.
- $X \subseteq (\mathbb{R}^n, d_p)$ is totally bounded iff X is bounded.
- $(1 + \frac{1}{n})^n$ converges to e.

- **Complete:** every Cauchy seq is convergent.
- $(\mathbb{R}^\omega, D)$ is complete.
- Subspace of complete metric space is complete iff closed.
- Cpt iff complete and totally bounded.
- **Heine-Borel Thm:** cpt in $\mathbb{R}^n$ iff closed and bounded.
- **Locally cpt:** $\exists$ cpt K and open U s.t. $x \in U \subseteq K$.
- X is locally cpt T_2 iff $\exists$ cpt T_2 Y and $h_Y: X \rightarrow Y$ homeomorphic onto $h_Y(X)$ and $Y = h_Y(X) \cup \{p\}$.
- Open sets in Y are open sets in X and complements of cpt sets in X . h_Y is just the identity.
- $\exists$ homeomorphism f s.t. $f \upharpoonright_{h_Y(X)} = h_{Y'} \circ h_Y^{-1} \upharpoonright_{h_Y(X)}$
- $\overline{\mathbb{D}}$ and $\mathbb{S}^2$ are the one-point cptification for $\mathbb{D}$ and $\mathbb{R}^n$ because $\mathbf{v} \mapsto \frac{\mathbf{v}}{1+\|\mathbf{v}\|}$ is bijection.
- A T_2 space is locally cpt iff $\forall x$ and open U s.t. $x \in U$, $\exists$ open cpt V s.t. $x \in V \subseteq U$.
- X locally cpt, then $A \subseteq X$ locally cpt if A closed or A open and X T_2 .
- X locally cpt T_2 iff $\exists$ cpt T_2 Y s.t. X homeomorphic to open $A \subseteq Y$.
- **Uniform metric:** $\bar{\rho}(f, g) = \sup \{ \rho(\pi_\alpha(f), \pi_\alpha(g)) \}$.
- $\text{Prod.} \subseteq \text{Uniform} \subseteq \text{Box}$
- Y is complete $\implies Y^X$ is complete.
- $\mathcal{C}(X, Y)$ and $\mathcal{B}(X, Y)$ are closed, and complete if Y is.
- **Supremum metric:** $d_{\text{sup}}(f, g) = \sup d(f(x), g(x))$.
- **Metric completion:** isometric ϕ s.t. $\overline{\phi(X)} = Y$.
- Every metric space has a unique metric completion up to isometry s.t. $f \upharpoonright_{\phi(X)} = \phi' \circ \phi^{-1}$.
- **Topo. of cpt convergence:** gen'ed by $B(K, f, \epsilon) := \{g \in Y^X : d_{\text{sup}}(f \upharpoonright_K, g \upharpoonright_K) < \epsilon\}$'s.
- **Cpt-open topo.:** gen'ed by $S(K, U) := \{g \in \mathcal{C}(X, Y) : g(K) \subseteq U\}$'s.
- Both are the same on $\mathcal{C}(X, Y)$.
- **Cptly gen'ed:** A open (closed) in X iff $A \cap K$ open (closed) in K for all cpt sets in X .

- Locally cpt or 1st countable $\implies$ cptly gen'ed.
- Cptly gen'ed $\implies (f \text{ cont. iff } f \upharpoonright_K \text{ cont. } \forall \text{ cpt } K)$.
- X cptly gen'ed $\implies \mathcal{C}(X, Y)$ closed under topo. of cpt convergence.
- **Equicont.:** $\exists$ open $U \forall \epsilon > 0$ s.t. $\forall f \in \mathcal{Y} \subseteq \mathcal{C}(X, Y)$, $d(y_{x_0}, y_x) < \epsilon \forall x \in U$.
- Totally bounded in $\bar{\rho} \implies$ equicont.
- Converge in prod. topo. = ptwise convergence.
- Equicont. family has the same topo. for cpt and ptwise convergence.
- **Arzela-Ascoli Thm:** $\mathcal{Y}_a := \{f(a) : f \in \mathcal{Y}\}$. $\overline{\mathcal{Y}} \subseteq \mathcal{C}(X, Y)$ is cpt if it's equicont. and $\overline{\mathcal{Y}_a}$ is cpt $\forall a \in X$. $\iff$ is true if X is locally cpt T_2 .

Connectedness

- Connected iff only X and $\emptyset$ are clopen.
- $X = (U, V)$, $Y \subseteq X$ connected $\implies Y \subseteq U$ or $Y \subseteq V$.
- Connected sets with $\cap \neq \emptyset$ has connected $\cup$.
- A connected and $A \subseteq B \subseteq \overline{A} \implies B$ connected.
- Cont. f maps connected to connected.
- Prod. of connected is connected.
- Path connected $\implies$ connected.
- Path component $\subseteq$ cc.
- **Locally connected:** $\forall$ open U s.t. $x \in U$, $\exists$ open connected V s.t. $x \in V \subseteq U$.
- S : graph of $\sin \frac{2\pi}{x}$ at $(0, 1]$. $\overline{S} = S \cup \{0\} \times [-1, 1]$ is connected, but not path/locally connected.
- Locally connected iff every cc of every open set is open.
- CCs and path components are the same in locally path connected space.
- Quotient topo. on $\tilde{X}$: cc's of locally path connected space, is discrete.
- Locally path connected $\implies$ locally connected.