

Gradients in Logistic Regression and Perceptrons Supplement

CS114B Lab 4

Kenneth Lai

February 17, 2022

Gradients in Logistic Regression

- ▶ Cross-entropy loss $L(\hat{y}, y) = -[y \log \hat{y} + (1 - y) \log(1 - \hat{y})]$

Gradients in Logistic Regression

- ▶ Cross-entropy loss $L(\hat{y}, y) = -[y \log \hat{y} + (1 - y) \log(1 - \hat{y})]$
- ▶ We want to compute $\frac{\partial L}{\partial w_j}$


Gradients in Logistic Regression

- ▶ Cross-entropy loss $L(\hat{y}, y) = -[y \log \hat{y} + (1 - y) \log(1 - \hat{y})]$
- ▶ We want to compute $\frac{\partial L}{\partial w_j}$
- ▶ Chain Rule of calculus: $\frac{dy}{dx} = \frac{dy}{dz} \frac{dz}{dx}$

Gradients in Logistic Regression

- ▶ Cross-entropy loss $L(\hat{y}, y) = -[y \log \hat{y} + (1 - y) \log(1 - \hat{y})]$
- ▶ We want to compute $\frac{\partial L}{\partial w_j}$
- ▶ Chain Rule of calculus: $\frac{dy}{dx} = \frac{dy}{dz} \frac{dz}{dx}$
- ▶ Looking at the graph: $\frac{\partial L}{\partial w_j} = \frac{\partial L}{\partial \hat{y}} \frac{\partial \hat{y}}{\partial z} \frac{\partial z}{\partial w_j}$

Graphical Representation of a Linear Classifier (1)



Gradients in Logistic Regression

$$\blacktriangleright \frac{\partial L}{\partial w_j} = \frac{\partial L}{\partial \hat{y}} \frac{\partial \hat{y}}{\partial z} \frac{\partial z}{\partial w_j}$$

Gradients in Logistic Regression

$$\blacktriangleright \frac{\partial L}{\partial w_j} = \frac{\partial L}{\partial \hat{y}} \frac{\partial \hat{y}}{\partial z} x_j$$

$$\blacktriangleright \frac{\partial z}{\partial w_j} = x_j$$

Gradients in Logistic Regression

- ▶ $\frac{\partial L}{\partial w_j} = \frac{\partial L}{\partial \hat{y}} \hat{y}(1 - \hat{y}) x_j$
 - ▶ $\frac{\partial z}{\partial w_j} = x_j$
 - ▶ For the logistic function: $\frac{\partial \hat{y}}{\partial z} = \hat{y}(1 - \hat{y})$

Gradients in Logistic Regression

- ▶ $\frac{\partial L}{\partial w_j} = \frac{\hat{y} - y}{\hat{y}(1 - \hat{y})} \hat{y}(1 - \hat{y}) x_j$
 - ▶ $\frac{\partial z}{\partial w_j} = x_j$
 - ▶ For the logistic function: $\frac{\partial \hat{y}}{\partial z} = \hat{y}(1 - \hat{y})$
 - ▶ For the cross-entropy loss: $\frac{\partial L}{\partial \hat{y}} = \frac{\hat{y} - y}{\hat{y}(1 - \hat{y})}$

Gradients in Logistic Regression

- ▶ $\frac{\partial L}{\partial w_j} = \frac{\hat{y} - y}{\hat{y}(1 - \hat{y})} \hat{y}(1 - \hat{y}) x_j = (\hat{y} - y) x_j$
 - ▶ $\frac{\partial z}{\partial w_j} = x_j$
 - ▶ For the logistic function: $\frac{\partial \hat{y}}{\partial z} = \hat{y}(1 - \hat{y})$
 - ▶ For the cross-entropy loss: $\frac{\partial L}{\partial \hat{y}} = \frac{\hat{y} - y}{\hat{y}(1 - \hat{y})}$

Gradients in Logistic Regression

- ▶ $\frac{\partial L}{\partial w_j} = \frac{\hat{y} - y}{\hat{y}(1 - \hat{y})} \hat{y}(1 - \hat{y}) x_j = (\hat{y} - y) x_j$
 - ▶ $\frac{\partial z}{\partial w_j} = x_j$
 - ▶ For the logistic function: $\frac{\partial \hat{y}}{\partial z} = \hat{y}(1 - \hat{y})$
 - ▶ For the cross-entropy loss: $\frac{\partial L}{\partial \hat{y}} = \frac{\hat{y} - y}{\hat{y}(1 - \hat{y})}$
- ▶ $\frac{\partial L}{\partial b} = \frac{\hat{y} - y}{\hat{y}(1 - \hat{y})} \hat{y}(1 - \hat{y})(1) = \hat{y} - y$
 - ▶ $\frac{\partial z}{\partial b} = 1$
 - ▶ ...

Gradients in Perceptrons

- ▶ Perceptron loss $L(\hat{y}, y) = (\hat{y} - y)z$

Gradients in Perceptrons

- ▶ Perceptron loss $L(\hat{y}, y) = (\hat{y} - y)z$
- ▶ We want to compute $\frac{\partial L}{\partial w_j}$
- ▶ Chain Rule of calculus: $\frac{dy}{dx} = \frac{dy}{dz} \frac{dz}{dx}$
- ▶ Looking at the graph: $\frac{\partial L}{\partial w_j} = \frac{\partial L}{\partial \hat{y}} \frac{\partial \hat{y}}{\partial z} \frac{\partial z}{\partial w_j}$

Gradients in Perceptrons

- ▶ Perceptron loss $L(\hat{y}, y) = (\hat{y} - y)z$
- ▶ We want to compute $\frac{\partial L}{\partial w_j}$
- ▶ Chain Rule of calculus: $\frac{dy}{dx} = \frac{dy}{dz} \frac{dz}{dx}$
- ▶ Looking at the graph: $\frac{\partial L}{\partial w_j} = \frac{\partial L}{\partial \hat{y}} \frac{\partial \hat{y}}{\partial z} \frac{\partial z}{\partial w_j}$
 - ▶ Problem: $\frac{\partial \hat{y}}{\partial z} = 0$ almost everywhere

Gradients in Perceptrons

- ▶ Perceptron loss $L(\hat{y}, y) = (\hat{y} - y)z$
- ▶ We want to compute $\frac{\partial L}{\partial w_j}$
- ▶ Chain Rule of calculus: $\frac{dy}{dx} = \frac{dy}{dz} \frac{dz}{dx}$
- ▶ Looking at the graph: $\frac{\partial L}{\partial w_j} = \frac{\partial L}{\partial \hat{y}} \frac{\partial \hat{y}}{\partial z} \frac{\partial z}{\partial w_j}$
 - ▶ Problem: $\frac{\partial \hat{y}}{\partial z} = 0$ almost everywhere
 - ▶ A change in z does not result in a change in \hat{y} unless you are at the threshold

Gradients in Perceptrons

- ▶ Perceptron loss $L(\hat{y}, y) = (\hat{y} - y)z$
- ▶ We want to compute $\frac{\partial L}{\partial w_j}$
- ▶ Chain Rule of calculus: $\frac{dy}{dx} = \frac{dy}{dz} \frac{dz}{dx}$
- ▶ Looking at the graph: $\frac{\partial L}{\partial w_j} = \frac{\partial L}{\partial \hat{y}} \frac{\partial \hat{y}}{\partial z} \frac{\partial z}{\partial w_j}$
 - ▶ Problem: $\frac{\partial \hat{y}}{\partial z} = 0$ almost everywhere
 - ▶ A change in z does not result in a change in \hat{y} unless you are at the threshold
 - ▶ Solution: consider $\frac{\partial L}{\partial z}$ directly

Gradients in Perceptrons

- ▶ Perceptron loss $L(\hat{y}, y) = (\hat{y} - y)z$
- ▶ We want to compute $\frac{\partial L}{\partial w_j}$
- ▶ Chain Rule of calculus: $\frac{dy}{dx} = \frac{dy}{dz} \frac{dz}{dx}$
- ▶ Looking at the graph: $\frac{\partial L}{\partial w_j} = \frac{\partial L}{\partial z} \frac{\partial z}{\partial w_j}$

Gradients in Perceptrons

$$\blacktriangleright \frac{\partial L}{\partial w_j} = \frac{\partial L}{\partial z} \frac{\partial z}{\partial w_j}$$

Gradients in Perceptrons

$$\blacktriangleright \frac{\partial L}{\partial w_j} = \frac{\partial L}{\partial z} x_j$$

$$\blacktriangleright \frac{\partial z}{\partial w_j} = x_j$$

Gradients in Perceptrons

- ▶ $\frac{\partial L}{\partial w_j} = (\hat{y} - y)x_j$

- ▶ $\frac{\partial z}{\partial w_j} = x_j$

- ▶ For the perceptron loss: $\frac{\partial L}{\partial z} = \hat{y} - y$

Gradients in Perceptrons

- ▶ $\frac{\partial L}{\partial w_j} = (\hat{y} - y)x_j$
 - ▶ $\frac{\partial z}{\partial w_j} = x_j$
 - ▶ For the perceptron loss: $\frac{\partial L}{\partial z} = \hat{y} - y$
- ▶ $\frac{\partial L}{\partial b} = \hat{y} - y$
 - ▶ $\frac{\partial z}{\partial b} = 1$
 - ▶ ...