

Hidden Variables

Qi Zhao

August 5, 2018

1. Introduction

To make the density multi-modal, it introduces mixture models [4]. To make the density robust, it replaces the normal with the t-distribution [2]. To cope with parameter estimation in high dimensions, it introduces subspace models [6].

The new models have much in common with each other. In each case it introduces a hidden or latent variable h_i associated with each observed data point x_i . The hidden variable induces the more complex properties of the resulting pdf. To model a complex probability density function over the variable x , it will introduce a hidden or latent variable h , which may be discrete or continuous.

2. Descriptions

To exploit the hidden variables, it describes the final density $Pr(x)$ as the marginalization of the joint density $Pr(x,h)$ [5] between x and h in Equation 1 so that it is relatively simple to model, but produces an expressive family of marginal distributions $Pr(x)$ when it integrates over h (see Figure 1).

$$Pr(x) = \int Pr(x, h) dh \quad (1)$$

Whatever form can be choose for the joint distribution in Equation 2, there are two possible approaches to fitting the model to training data $\{x_i\}_{i=1}^I$ using the maximum likelihood [3] method. It could directly maximize the log likelihood of the distribution $Pr(x)$ from the left hand side of equation (see Equation 3). It uses the expectation maximization [1] algorithm, which works directly with the right-hand side of equation (see Equation 4).

$$Pr(x|\theta) = \int Pr(x, h|\theta) dh \quad (2)$$

$$\hat{\theta} = \operatorname{argmax}_{\theta} \left[\sum_{i=1}^I \log Pr(x_i|\theta) \right] \quad (3)$$

$$\hat{\theta} = \operatorname{argmax}_{\theta} \left[\sum_{i=1}^I \log \left[\int Pr(x_i, h_i|\theta) dh_i \right] \right] \quad (4)$$




Figure 1. Using hidden variables to help model complex densities. One way to model the density $Pr(x)$ is to consider the joint probability distribution $Pr(x,h)$ between the observed data x and a hidden variable h . The density $Pr(x)$ can be considered as the marginalization of (integral over) this distribution with respect to the hidden variable h . As we manipulate the parameters of this joint distribution, the marginal changes and the agreements with the observed data $\{x_i\}_{i=1}^I$ increases or decreases. Sometimes it is easier to fit the distribution in this indirect way than to directly manipulate $Pr(x)$.

References

- [1] C. Carson, S. Belongie, H. Greenspan, and J. Malik. Blobworld: image segmentation using expectation-maximization and its application to image querying. *IEEE Transactions on Pattern Analysis & Machine Intelligence*, 24(8):1026–1038, 2002. 1
- [2] E. A. Coornish. The multivariate t-distribution associated with a set of normal sample deviates. *Australian Journal of Physics*, 7(4):531–542, 1954. 1

- [3] A. P. Dempster, N. M. Laird, and D. B. Rubin. Maximum likelihood from incomplete data via the EM algorithm. *Journal of the Royal Statistical Society*, 39(1):1–38, 1977. [1](#)
- [4] M. A. T. Figueiredo and A. K. Jain. Unsupervised learning of finite mixture models. *IEEE Transactions on Pattern Analysis & Machine Intelligence*, 24(3):381–396, 2000. [1](#)
- [5] L. A. Shepp. The joint density of the maximum and its location for a wiener process with drift. *Journal of Applied Probability*, 16(2):423–427, 1979. [1](#)
- [6] M. Verhaegen and D. Westwick. Identifying MIMO wiener systems using subspace model identification methods. In *IEEE Conference on Decision and Control*, pages 4206–4211, 1995. [1](#)