

Week 1 • Problem Set 4 • Deviations and acyclic closures

Quick Problem:

Let (Q, \mathfrak{n}, k) be a regular local ring and let $R = Q/I$ with $I \subseteq \mathfrak{n}^2$.

- From how the acyclic closure is constructed, identify $\varepsilon_2(R)$ as a commonly known quantity involving some kind of homology.
- Explain what the correspondence between the acyclic closure of $R \rightarrow k$ and the minimal model of $Q \rightarrow R$ implies about the deviation $\varepsilon_2(R)$.

Problems:

- Consider the running example from the lectures:

Let $R = k[[x, y]]/(f)$ with $f = x^3 + y^3$, and set $S = k = R/(x, y)$.

- Determine some or all of the deviations of R .
- Determine some or all of the complex $\text{Ind}_R^\gamma R\langle X \rangle$ of Γ -indecomposables where $R\langle X \rangle$ is an acyclic closure of k over R .

- Consider the example you computed in #1 of Problem Set 2:

$$R = k[[x, y]]/(x^2, xy) \longrightarrow S = k = R/(x, y)$$

Running the Macaulay2 code from Wednesday's M2 session yields the output:

$$\{\mathbf{x}, \mathbf{y}, \mathbf{xT}_1, \mathbf{yT}_1, -\mathbf{xT}_1\mathbf{T}_2, -\mathbf{yT}_1\mathbf{T}_3, \mathbf{xT}_1\mathbf{T}_2\mathbf{T}_3, \mathbf{xT}_1\mathbf{T}_2\mathbf{T}_4\}$$

giving a list of the images of the variables under the differential of the variables denoted by T_1, T_2, \dots , numbered in the order that they were adjoined. Translating, we see that the acyclic closure looks like:

$$R\langle T \rangle = \langle T_1, T_2, \dots \rangle$$

with

$$\begin{aligned} \partial(T_1) &= x, \partial(T_2) = y, \\ \partial(T_3) &= xT_1, \partial(T_4) = yT_1, \\ \partial(T_5) &= -xT_1T_2, \partial(T_6) = -yT_1T_3, \\ \partial(T_7) &= xT_1T_2T_3, \partial(T_8) = xT_1T_2T_4 \end{aligned}$$

- Find the degree of each T_i above.
 - Determine some or all of the deviations of R .
 - Determine some or all of the complex $\text{Ind}_R^\gamma R\langle T \rangle$ of Γ -indecomposables.
- Here is an example of a homomorphism whose acyclic closure is not a minimal complex. Let k be a field of characteristic zero. Consider the canonical surjection.

$$R = k[[x, y]] \longrightarrow S = R/(x^2, xy)$$

Compute a few steps of the acyclic closure until you see the non-minimality appear.

- Show that a surjective map of finite projective dimension $\varphi: R \rightarrow S$ is complete intersection if and only if the acyclic closure is minimal.

Lastly, here are two proofs that we skipped in the lecture for you to try.

Let $\varphi: R \rightarrow S$ be a surjective homomorphism of local rings. (Much does not actually require surjectivity, but we assume this for simplicity here.)

5. Let $R \rightarrow R\langle X \rangle$ be a semi-free Γ -extension resolving S with $X_0 = \emptyset$.

For any $x \in X_n$ such that the class of x generates a free summand of the cokernel of $\partial_{n+1}^I: SX_{n+1} \rightarrow SX_n$, there exists an R -linear chain Γ -derivation $\theta_x: R\langle X \rangle \rightarrow R\langle X \rangle$ of degree $-n$ such that $\theta_x(x) = 1$ and $\theta_x(X_n \setminus \{x\}) = 0$.

Hint: Note that $Sx \cap \text{im } \partial_{n+1}^I = 0$. Define a map from I to $R\langle X \rangle$. Why is it a chain map? Why can you extend it to a derivation from $R\langle X \rangle$?

6. Let $R \rightarrow R\langle X \rangle$ be a semi-free Γ -extension resolving S with $X_0 = \emptyset$.

Prove that $R\langle X \rangle$ is an acyclic closure of S over R if and only if the complex $I = \text{Ind}_R^\gamma R\langle X \rangle$ of Γ -indecomposables is minimal ($\partial(I) \subseteq \mathfrak{m}I$).