
a
Longitudinal transfusion target trial
18,742 triggers - 2,432 prior RBC - 742 not at landmark
Baseline lookback
RBC start grace period
Post-landmark Hb response
90-day mortality
Hb 7--9 g/dL trigger
Treatment landmark
Final n = 15,568: 3,017 early RBC; 3,312 deaths
-24 h
0
6 h
24 h
90 d
Timeline shown schematically
b
Retrospective temporal holdout
Discover graph 2008--2016
n = 10,179
RBC 22.9%
Select learner 2017--2019
n = 3,127
RBC 17.1%
Temporal test 2020--2022
n = 2,262
RBC 6.5%
Non-overlapping eras separate discovery, selection, and evaluation.
c
Role safety and overlap
ESS/n=0.905
Assignment set C (7)
Retained S (1)
Before
After
0.0
0.1
0.2
0.3
0.4
Mean |SMD|
0.90
d
R-loss gain (×10−4)
20 repeated splits
1.25
Temporal holdout
−1
0
1
2
3
Positive values favor CCIA C+S
0.005
e
DR-Qini gain
0.032
0.00
0.02
0.04
0.06
Positive values favor CCIA C+S
0.46
f
Policy-risk gain (pp)
2.38
0
2
4
6
Positive values favor CCIA C+S