
a
Target trial and analysis cohort
Baseline lookback
RBC initiation 0-6 h
Hb response
90-day mortality
-24 h
0 h
6 h
24 h
90 d
Trigger
Landmark
18,742 triggers | 15,568 landmark-eligible
2,432 prior RBC | 742 unavailable
1,578 excluded: no treated temporal support
13,990 analyzed | 2,819 RBC | 2,649 deaths
b
Treatment and outcome shift across eras
Early RBC
90-day mortality
Cohort rate (%)
23.9%
21.1%
24.3%
17.0%
18.1%
7.0%
n=9,018
n=2,865
n=2,107
2008--2016 Discovery
2017--2019 Selection
2020--2022 Temporal test
c
Temporal balance across 11 adjustment variables
Mean 0.386 to 0.111 | ESS/n=0.773
0.10
Before
Weighted
Temperature
Cardiac ICU
Urine recorded
Medical service
Trigger time
RASS
Glucose unrecorded
Trigger Hb
Minimum Hb
Lactate
GCS
Absolute standardized mean difference
d
Effect risk: R-loss
All variables
RITE
Gain [95% interval]
Repeated
3.48 [0.68, 6.25]
12/20 wins
Temporal
1.27 [-6.61, 9.22]
P(gain > 0)=62.0%
Gain (×10⁻⁴)
e
Ranking: DR-Qini
All variables
RITE
Gain [95% interval]
Repeated
0.010 [0.003, 0.017]
15/20 wins
Temporal
0.014 [-0.022, 0.051]
P(gain > 0)=78.2%
Gain
f
Policy value
All variables
RITE
Gain [95% interval]
Repeated
1.59 [0.62, 2.54]
15/20 wins
Temporal
1.78 [-5.55, 7.37]
P(gain > 0)=51.9%
Gain (pp)