
RITE versus all-variable baseline · Positive gain favours RITE
a
R-loss
Gain (×10⁻⁴) · 12/20 positive
b
DR-Qini
Gain · 15/20 positive
c
Policy risk reduction
Gain (pp) · 15/20 positive
M
T
Bars: paired splits · M: mean · T: temporal test · Whiskers: 95% CI · pp: percentage points
d
Temporal covariate balance
Before
Weighted
Reduction
Minimum Hb 0.366 0.074
GCS 0.376 0.040
RASS 0.260 0.124
Temperature 0.109 0.185
Glucose unrecorded 0.461 0.091
Lactate 0.284 0.050
Medical service 0.615 0.143
Trigger time 0.434 0.126
Trigger Hb 0.345 0.075
Cardiac ICU 0.422 0.163
Urine recorded 0.574 0.147
Mean |SMD| 0.386 → 0.111 ESS/n = 0.773
Variables with |SMD| ≤ 0.10: 0/11 → 5/11
e
Clinical eras
Early RBC
90-day mortality
23.9%
21.1%
24.3%
17.0%
18.1%
7.0%
2008–16 Discovery
2017–19 Selection
2020–22 Temporal
n 9,018 2,865 2,107
RBC 2,154 518 147
Deaths 1,534 604 511