
a
BP medication: SBP
n = 7,711; S-learner
R-loss
mmHg²
DR-Qini
mmHg
Budget outcome
mmHg
Cumulative splits (%)
100
50
0
−1
0
2
−0.5
0.5
1.5
−0.5
0.5
1.5
15/20 positive
9/20 positive
12/20 positive
+0.516
+0.017
+0.073
b
Cholesterol medication: TC
n = 4,521; Class transformation
R-loss
(mg/dL)²
DR-Qini
mg/dL
Budget outcome
mg/dL
Cumulative splits (%)
100
50
0
0
60
120
0
1
2
−2
0
4
19/20 positive
16/20 positive
10/20 positive
+78.739
+0.747
+0.558
Gain > 0 favors RITE; diamonds: mean and 95% CI
c
BP task: variable roles
10 variables in C/S at either stage
Statistical
Reviewed
C 9
S 1
6 retained in C
C 6
Height
S 2
Insulin use; vigorous work
Weight
O 2
C: proximal
S: precision
O: outside C/S
Review decisions: 37 pairs
12
4
4
15
2
Oriented/confirmed
Semantic removal
Task-order removal
Retained
Query/DAG rules
d
Cholesterol task: variable roles
12 variables in C/S at either stage
Statistical
Reviewed
C 6
S 5
O 1
6 retained in C
4 retained in S
Height
CHD history
C 7
S 4
O 1
C: proximal
S: precision
O: outside C/S
Review decisions: 33 pairs
10
2
2
18
1