B2 Lunn–McNeil test of whether pattern effects differ across intrahepatic / extrahepatic / systemic HCC death

Overall likelihood-ratio tests (reference-independent; non-HCC and undetermined deaths censored; strata = cause x treatment group; all covariates interact with cause)

Test	LR chi-square	df	P
Pattern x cause (9 patterns)	60.17	16	<0.001
Site x cause (IH / EH / Sys)	48.80	4	<0.001
Form x cause (TG / NL / TG+NL)	7.69	4	0.104

Follow-up (weighted effect coding): csHR of each pattern vs the pattern-weighted average, per cause; Wald test that the effect is equal across the three causes

Pattern	IH death	EH death	Systemic death	Wald chi2 (df=2)	P	P (Holm)
IH-TG	0.99 (0.88-1.12)	0.78 (0.55-1.10)	0.70 (0.56-0.89)	7.19	0.028	0.138
IH-NL	0.90 (0.77-1.05)	0.46 (0.30-0.72)	0.61 (0.45-0.81)	11.50	0.003	0.019
IH-TG+NL	1.49 (1.16-1.91)	1.17 (0.57-2.42)	0.85 (0.50-1.45)	3.62	0.164	0.499
EH-TG	1.41 (1.05-1.90)	2.07 (1.22-3.50)	1.22 (0.77-1.94)	2.28	0.319	0.639
EH-NL	0.75 (0.62-0.91)	1.40 (0.98-1.99)	1.18 (0.91-1.53)	13.21	0.001	0.011
EH-TG+NL	1.02 (0.55-1.91)	1.90 (0.59-6.16)	1.41 (0.65-3.06)	0.98	0.613	0.639
Sys-TG	1.39 (0.89-2.19)	0.88 (0.28-2.78)	2.50 (1.50-4.16)	4.16	0.125	0.499
Sys-NL	0.93 (0.64-1.33)	2.72 (1.55-4.76)	2.00 (1.32-3.04)	12.89	0.002	0.011
Sys-TG+NL	1.14 (0.94-1.40)	1.49 (0.93-2.39)	2.20 (1.68-2.89)	14.52	<0.001	0.006