A Adjusted 12-month cumulative incidence of death after first progression

Cause-specific Cox + G-computation (standardised to all 1786 progressors); % (95% bootstrap CI, B = 1000); n = events within 12 months

	All-cause	Intrahepatic HCC death	Extrahepatic HCC death	Systemic HCC death	Non-HCC death
Sys-TG	77.5 (65.1–88.6) n=36	43.5 (29.7–57.6) n=18	4.1 (0.0–11.5) n=3	26.4 (14.7–40.5) n=14	3.2 (0.0–11.2) n=1
EH-TG+NL	75.4 (63.1–87.5) n=21	36.1 (20.3–51.3) n=10	9.4 (0.0–20.5) n=2	17.1 (6.5–31.3) n=6	12.5 (0.0–29.7) n=3
Sys-TG+NL	75.2 (69.6–81.3) n=154	39.0 (33.3–45.6) n=88	7.3 (3.9–11.7) n=13	25.3 (19.5–31.0) n=47	3.3 (0.9–6.4) n=6
EH-TG	74.9 (67.7–81.8) n=72	47.0 (37.6–55.9) n=42	10.1 (5.2–15.8) n=11	14.2 (8.2–21.1) n=15	3.3 (0.6–8.0) n=3
Sys-NL	74.0 (67.3–81.3) n=63	33.7 (24.9–42.1) n=26	13.4 (6.9–22.6) n=13	24.3 (15.6–33.3) n=22	2.3 (0.0–6.2) n=2
IH-TG+NL	71.7 (64.7–78.8) n=74	51.5 (43.8–59.6) n=55	6.2 (2.1–11.1) n=6	10.5 (5.3–16.0) n=10	3.2 (0.7–6.5) n=3
EH-NL	64.9 (60.5–69.5) n=159	32.2 (27.4–37.2) n=76	8.4 (5.5–11.4) n=18	17.7 (13.4–22.5) n=48	6.2 (3.9–9.2) n=15
IH-TG	62.1 (58.6–66.2) n=269	42.1 (38.0–46.5) n=193	4.9 (3.0–7.0) n=19	10.8 (8.1–13.9) n=38	3.8 (2.2–5.6) n=16
IH-NL	58.0 (53.5–62.6) n=161	40.5 (35.3–45.8) n=109	3.1 (1.6–4.7) n=8	10.0 (7.1–13.4) n=29	3.9 (2.3–6.0) n=15

12-month CIF (%)

0 25 50 75

Cause composition at 12 months



Adjusted for TTP, age, sex, HBV, ECOG PS, ALBI grade, AFP ≥400, PIVKA-II ≥1000, tumour number >3, tumour size, Vp class, HVTT, EHS, cirrhosis; stratified by treatment group. Rows ordered by adjusted all-cause CIF. Cause-specific CIFs plus undetermined deaths sum to all-cause. Composition labels show % of deaths by 12 months (segments <8% unlabelled).