B1 Subdistribution hazard ratios (Fine–Gray) by progression pattern (reference: Sys-TG)

Fine–Gray via survival::finegray with stratum-specific censoring weights; robust SE

	Intrahepatic HCC death	Extrahepatic HCC death	Systemic HCC death	Non-HCC death
IH-TG	0.91 (0.61–1.36)	1.07 (0.35–3.24)	0.41 (0.24–0.72)**	NE (sparse reference)
IH-NL	0.84 (0.55–1.27)	0.71 (0.22–2.29)	0.36 (0.20–0.64)***	NE (sparse reference)
IH-TG+NL	1.25 (0.78–2.00)	1.29 (0.36–4.60)	0.40 (0.19–0.82)*	NE (sparse reference)
EH-TG	0.92 (0.56–1.51)	2.37 (0.73–7.65)	0.49 (0.25–0.96)*	NE (sparse reference)
EH-NL	0.59 (0.38–0.92)*	1.98 (0.67–5.87)	0.68 (0.40–1.17)	NE (sparse reference)
EH-TG+NL	0.68 (0.32–1.44)	1.84 (0.41–8.35)	0.69 (0.28–1.69)	NE (sparse reference)
Sys-TG	Reference	Reference	Reference	Reference (1 event)
Sys-NL	0.62 (0.37–1.04)	3.10 (0.94–10.25)	0.96 (0.51–1.78)	NE (sparse reference)
Sys-TG+NL	0.78 (0.49–1.23)	1.53 (0.49–4.80)	1.00 (0.57–1.75)	NE (sparse reference)
Overall Wald P (df=8)	**P = 0.003**	**P = 0.003**	**P < 0.001**	**P = 0.256**

$\log_2(\text{sHR})$

-2 -1 0 1 2

Adjusted for TTP, age, sex, HBV, ECOG PS, ALBI grade, AFP ≥400, PIVKA-II ≥1000, tumour number >3, tumour size, Vp class, HVTT, EHS, cirrhosis; stratified by treatment group. NE: <3 events in the cell or in the reference cell. Overall Wald test (df = 8, robust variance) is reference-independent.