B2 Lunn–McNeil test of whether pattern effects differ across intrahepatic / extrahepatic / systemic HCC death

Overall likelihood-ratio tests (reference-independent; non-HCC and undetermined deaths censored; strata = cause x treatment group; all covariates interact with cause)

Test	LR chi-square	df	P
Pattern x cause (9 patterns)	60.54	16	<0.001
Site x cause (IH / EH / Sys)	48.63	4	<0.001
Form x cause (TG / NL / TG+NL)	7.36	4	0.118

Follow-up (weighted effect coding): csHR of each pattern vs the pattern-weighted average, per cause; Wald test that the effect is equal across the three causes

Pattern	IH death	EH death	Systemic death	Wald chi2 (df=2)	P	P (Holm)
IH-TG	0.99 (0.88-1.12)	0.78 (0.55-1.10)	0.70 (0.56-0.89)	7.18	0.028	0.138
IH-NL	0.90 (0.77-1.04)	0.46 (0.30-0.72)	0.61 (0.45-0.81)	11.42	0.003	0.020
IH-TG+NL	1.49 (1.16-1.91)	1.17 (0.57-2.42)	0.85 (0.50-1.45)	3.61	0.164	0.534
EH-TG	1.41 (1.05-1.90)	2.07 (1.22-3.52)	1.22 (0.77-1.93)	2.33	0.312	0.624
EH-NL	0.75 (0.62-0.91)	1.40 (0.98-1.99)	1.18 (0.91-1.53)	13.25	0.001	0.011
EH-TG+NL	1.02 (0.55-1.91)	1.90 (0.58-6.15)	1.41 (0.65-3.07)	0.97	0.614	0.624
Sys-TG	1.58 (1.11-2.24)	1.03 (0.39-2.76)	2.54 (1.62-3.97)	4.03	0.134	0.534
Sys-NL	0.93 (0.64-1.33)	2.72 (1.55-4.76)	2.00 (1.32-3.05)	12.86	0.002	0.011
Sys-TG+NL	1.07 (0.86-1.33)	1.47 (0.90-2.40)	2.16 (1.63-2.88)	15.05	<0.001	0.005