BLS	ОТ	AM	FR	EMR	AFA	MAFA
			✓	✓	✓	✓

Steps

Ensure you are always wearing proper PPE before performing the skill.

USING AN INHALER

- 1. Check the Six Rights of Medication before proceeding.
- 2. Tell the patient to rapidly shake the inhaler 3 or 4 times.
- 3. Have the patient remove the cap from the inhaler.
- 4. Tell the patient to breathe out, away from the inhaler.
- 5. Have the patient bring the inhaler to the mouth. Help to put the mouthpiece between the patient's teeth and then tell patient to close his or her lips around it.
- 6. Tell the patient to press the top of the inhaler once while taking one slow, full breath in.
- 7. Have the patient hold the breath for as long as comfortable (up to 10 seconds) and then exhale.

USING AN INHALER WITH A SPACER

- 1. Check the Six Rights of Medication before proceeding.
- 2. Tell the patient to rapidly shake the inhaler 3 or 4 times.
- 3. Have the patient remove the cap from the inhaler (and spacer, if applicable).
- 4. Have the patient put the inhaler into the spacer.
- 5. Tell the patient to breathe out, away from the spacer.
- 6. Have the patient bring the spacer to the mouth.
- 7. Tell the patient to press the top of the inhaler once while taking one slow, full breath in.
- 8. Have the patient hold the breath for as long as comfortable (up to 10 seconds) and then exhale.

Organizational/local protocol for :							

Skill Notes

• If the patient is unable to press the top of the inhaler, you may do it instead if the patient asks you to.