RGH: CLAS12 WITH A TRANSVERSELY POLARIZED TARGET

ZACH MEADOR CHRISTOPHER NEWPORT UNIVERSITY

1. Luminosity of CLAS12

The luminosity of the CLAS12 detector can be calculated by:

$$L = N_e \times N_p = \left(\frac{1}{q_e}\right) INl\rho$$

 N_e and N_p is the the number of electrons and protons, respectively. I is the current of the CLAS12 detector. The density and length of the target is represented by ρ and l, respectively. The fraction $1/q_e$ is the inverse of the charge of an electron. $N = \frac{n_p N_A}{M}$ where n_p is the number of protons, N_A is Avogadro's constant, and M is the molar mass of the target. This equation can be rearranged to find the desired current for running the CLAS12 at the correct luminosity:

$$I = \frac{L \times q_e}{Nl\rho}$$

The current is set by inside the gcard used to set the geometry of the CLAS12 detector:: <option name="LUMI_EVENT" value=" n_e , time window, bunch time" /> n_e is the number of electrons per event. The time window to observe occupancies inside the Drift Chamber subsystem of the CLAS12 detector is 250 ns. Once the desired current is known we can find the number of electrons per event:

$$I = \frac{n_e \times q_e}{timewindow}$$

$$n_e = \frac{I \times time \text{ window}}{q_e}$$

The target cell is not a single chunk of atoms. The atoms do not fill the entirety of the target cell. Therefore, atomic packing fraction, APF, must be calculated:

$$APF = \frac{N_{\text{particles}} \times V_{\text{particle}}}{V_{\text{target cell}}}$$
$$= \frac{2 \times \frac{4}{3} \pi r^{3}}{\left(\frac{4r}{\sqrt{3}}\right)^{3}}$$
$$APF = 0.68$$

Since we are interested in the empty space around the atom, the shape in question is a cube with an atom in the center and an eighth of an atom in each corner. The sum of all volumes

Date: March 2022.

in the cube leads to N=2: Therefore, the actual number of electrons in LUMI_EVENT




FIGURE 1. Cell Structure

needed to run the CLAS12 detector at the correct luminosity is:

$$n_e = n_e + 0.68n_e.$$