

IP/OP NO



Patient Name : Ms.Shubneet Kaur
Age/Gender : 24 Y 0 F 0 D /M
UHID/MR No : DPWA.0000005652
Visit ID : DPWAOPV8439

Ref Doctor : Dr.SELF

 Collected
 : 16/August/2021 09:55AM

 Received
 : 16/August/2021 02:52PM

 Reported
 : 17/August/2021 05:12PM

Status : Final Report
Client Name : PCC WAKAD
Patient location : Pune,Pune

| DEPARTMENT OF MOLECULAR BIOLOGY. |        |      |                 |        |  |  |  |
|----------------------------------|--------|------|-----------------|--------|--|--|--|
| COVID 19                         |        |      |                 |        |  |  |  |
| Test Name                        | Result | Unit | Bio. Ref. Range | Method |  |  |  |

| COVID-19(SARS-CoV 2)-REAL TIME PCR(RT-PCR) |   |                            |  |  |  |
|--|---|----------------------------|--|--|--|
| SAMPLE TYPE                                | NASOPHARYNGEAL<br>SWAB AND<br>OROPHARYNGEAL<br>SWAB |                            |  |  |  |
| SARS-CoV-2                                 | NOT DETECTED  | REAL TIME PCR (RT-<br>PCR) |  |  |  |
| RdRp Gene CT VALUE                         | -   |                            |  |  |  |
| N-Gene CT VALUE                            | - CT E 40   |                            |  |  |  |
| ORF1ab Gene CT VALUE                       |   |                            |  |  |  |

SRF ID: 1

## **Comment:**

## **Please Note:**

- \*Not Detected (Negative) Result indicate CT Value of >35 as per the testing kit used.
- \*The cut-off of the reported CT value is as per the manufacturer.
- \*CT values vary with the type of kit used, the types of samples collected and the various pre-analytical factors.

## Kit used: CoviPath TM COVID-19 RTPCR KIT

| Result                         | Interpretation  |  |  |
|--------------------------------|---|--|--|
| Detected (Positive) Result     | Indicates presence of detectable levels SARS-CoV-2 specific RNA (ORF1ab gene and/ or N gene) in patient's samp  |  |  |
| Not Detected (Negative) Result | Indicates absence of detectable levels SARS-CoV-2 specific RNA (ORF1ab gene and/ or N gene) in patient's sample |  |  |
| Indeterminate Result           | Indicates the target concentration is too low, or presence of interfering substances leading to PCR inhibition. |  |  |

**NOTE:** 

Page 1 of 2

<sup>\*</sup>The published studies are conflicting to definitively prove a direct correlation between disease severity/infectiousness and CT values therefore; relying on CT values for determining infectiousness of COVID-19 patients and deciding management protocols is left to clinical discretion.





Patient Name : Ms.Shubneet Kaur Age/Gender : 24 Y 0 F 0 D /M UHID/MR No : DPWA.0000005652

Visit ID : DPWAOPV8439
Ref Doctor : Dr.SELF

IP/OP NO :

| Collected : 16/August/2021 09:55AM | Received : 16/August/2021 02:52PM | Reported : 17/August/2021 05:12PM

Status : Final Report
Client Name : PCC WAKAD
Patient location : Pune,Pune

| DEPARTMENT OF MOLECULAR BIOLOGY. |        |      |                 |        |  |  |  |
|----------------------------------|--------|------|-----------------|--------|--|--|--|
| COVID 19                         |        |      |                 |        |  |  |  |
| Test Name                        | Result | Unit | Bio. Ref. Range | Method |  |  |  |

- If either of the genes cannot be determined or will give indeterminate results, further testing maybe required using another kit with a different primer and probe.
- A **Not Detected** (**Negative**) **Result**do not preclude SARS-CoV-2 and should not be used as the sole basis for patient management decisions. Kindly repeat test after 48 to 72 hrs if clinically suspected.
- Lower respiratory tract specimens are more representative and are preferred.
- If at a later date, suspicion of COVID-19 is strong, a fresh sample for Real Time PCR after a gap of 2-4 days after collection
  of first sample maybe considered.
- Please contact your Physician for necessary action to be taken and to limit the spread of infection.
- Clinical Correlation and Correlation with the history of the patient is required before arriving at any conclusion. Presence of Non-Specific interfering substances during this assay to be kept in mind. Please correlate clinically before arriving at any conclusion.
- False positive and false negative results can be due to multiple factors including sampling technique, transport & interference in the assay can affect the result.
- Presence PCR inhibitors in sample may lead to false negative or invalid results.
- Mutation in the target sequence of SARS-CoV-2 or change in the sequence due to virus evolution may lead to false negative results.
- Invalid Result: There is no typical S-shape amplification curve or Ct >35 or No Ct detected for target genes and internal control, indicating the specimen concentration is too low, or there are interfering substances that inhibit the reaction. If upon retest, the result is invalid again, another fresh sample should be collected and tested.

## **Disclaimer:**

This Test is based on real-time reverse transcriptase PCR technology for the qualitative detection of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) specific RNA

\*ICMR ID-AHLS001- HYDERABAD

\*ICMR ID-AHLVDLA- VIJAYAWADA

\*ICMR ID- DLUAHTMH- MUMBAI

\*ICMR ID-AHLLKDLKAP- KURNOOL

\*ICMR ID-ADNTKLKWB - KOLKATA

\*ICMR ID-AHLLDLDD-DELHI

\*ICMR ID-AHLLKDIALBK- BENGALURU

\*ICMR ID-APHLILKDLCTN- CHENNAI

\*ICMR ID-AHHHSPPMH- PUNE

(For kits determining RdRp gene or ORF1b, sensitivity / specificity of such kits shall be mentioned).

\*\*\* End Of Report \*\*\*

DR. PALAK KANSRA M.B.B.S. M.D(Microbiology) Consultant Microbiologist.

Page 2 of 2