

ID: W2678323552

TITLE: Severe impacts of brown tides caused by *Sargassum* spp. on near-shore Caribbean seagrass communities

AUTHOR: ['Brigitta I. van Tussenbroek', 'Héctor A. Hernández Arana', 'Rosa E. Rodríguez-Martínez', 'Julio Espinoza-Ávalos', 'Hazel M. Canizales-Flores', 'Carlos E. González-Godoy', 'M. Guadalupe Barba-Santos', 'Alejandro Vega-Zepeda', 'Ligia Collado-Vides']

ABSTRACT:

From mid-2014 until the end of 2015, the Mexican Caribbean coast experienced a massive influx of drifting *Sargassum* spp. that accumulated on the shores, resulting in build-up of decaying beach-cast material and near-shore murky brown waters (*Sargassum*-brown-tides, Sbt). The effects of Sbt on four near-shore waters included reduction in light, oxygen (hypoxia or anoxia) and pH. The monthly influx of nitrogen, and phosphorus by drifting *Sargassum* spp. was estimated at 6150 and 61 kg km⁻² respectively, resulting in eutrophication. Near-shore seagrass meadows dominated by *Thalassia testudinum* were replaced by a community dominated by calcareous rhizophytic algae and drifting algae and/or epiphytes, resulting in 61.6–99.5% loss of below-ground biomass. Near-shore corals suffered total or partial mortality. Recovery of affected seagrass meadows may take years or even decades, or changes could be permanent if massive influxes of *Sargassum* spp. recur.

SOURCE: Marine pollution bulletin

PDF URL: None

CITED BY COUNT: 264

PUBLICATION YEAR: 2017

TYPE: article

CONCEPTS: ['*Sargassum*', 'Seagrass', '*Thalassia testudinum*', 'Epiphyte', 'Algae', 'Eutrophication', 'Oceanography', 'Fucales', 'Rocky shore', 'Shore', 'Bay', 'Environmental science', 'Ecology', 'Fishery', 'Biology', 'Ecosystem', 'Geology', 'Nutrient']