

TITLE: Conservation and management of northeast Atlantic and Mediterranean maerl beds

AUTHOR: ['Carmen Barberá', 'César Bordehore', 'Joseph Borg', 'M. Glémarec', 'Jacques Grall', 'Jason M. Hall-Spencer', 'R. de la Huz', 'Edwin Lanfranco', 'Mariano Lastra', 'P. G. Moore', 'José Carlos Romero Mora', 'Marta Pita', 'Alfonso A. Ramos-Esplá', 'Miraine Rizzo', 'A. Sánchez-Mata', 'A.M. Seva', 'Patrick J. Schembri', 'C. Valle']

ABSTRACT:

Aquatic Conservation: Marine and Freshwater Ecosystems Volume 13, Issue S1 p. S65-S76 Research Article
 Conservation and management of northeast Atlantic and Mediterranean maerl beds C. Barbera, C. Barbera Ciencias Ambientales (Biología marina), Universidad de Alicante, 03080 Alicante, Spain Authorship alphabetical: cite as BIOMAERL teamSearch for more papers by this authorC. Bordehore, C. Bordehore Ciencias Ambientales (Biología marina), Universidad de Alicante, 03080 Alicante, Spain Authorship alphabetical: cite as BIOMAERL teamSearch for more papers by this authorJ.A. Borg, J.A. Borg Department of Biology, University of Malta, Msida, Malta Authorship alphabetical: cite as BIOMAERL teamSearch for more papers by this authorM. Glémarec, M. Glémarec UMR CNRS 6539 Ecologie benthique, Institut Universitaire Européen de la Mer, Place Copernic, 29280 Plouzane, France Authorship alphabetical: cite as BIOMAERL teamSearch for more papers by this authorJ. Grall, J. Grall UMR CNRS 6539 Ecologie benthique, Institut Universitaire Européen de la Mer, Place Copernic, 29280 Plouzane, France Authorship alphabetical: cite as BIOMAERL teamSearch for more papers by this authorJ. M. Hall-Spencer, J. M. Hall-Spencer School of Biological Sciences, University of Plymouth, Drake Circus, Plymouth, PL4 8AA, UK Authorship alphabetical: cite as BIOMAERL teamSearch for more papers by this authorCh. de la Huz, Ch. de la Huz Departamento de Ecología y Biología Animal, Facultad de Ciencias, Universidad de Vigo, Apdo.874, 36200 Vigo, Spain Authorship alphabetical: cite as BIOMAERL teamSearch for more papers by this authorE. Lanfranco, E. Lanfranco Department of Biology, University of Malta, Msida, Malta Authorship alphabetical: cite as BIOMAERL teamSearch for more papers by this authorM. Lastra, M. Lastra Departamento de Ecología y Biología Animal, Facultad de Ciencias, Universidad de Vigo, Apdo.874, 36200 Vigo, Spain Authorship alphabetical: cite as BIOMAERL teamSearch for more papers by this authorP.G. Moore, Corresponding Author P.G. Moore University Marine Biological Station, Millport, Isle of Cumbrae, UK, KA28 0EG Authorship alphabetical: cite as BIOMAERL teamCoordinatorUniversity Marine Biological Station Millport, Isle of Cumbrae, KA28 0EG, UKSearch for more papers by this authorJ. Mora, J. Mora Departamento de Biología Animal, Facultad de Biología, Universidad de Santiago de Compostela, 15706 Santiago de Compostela, Galicia, Spain Authorship alphabetical: cite as BIOMAERL teamSearch for more papers by this authorM.E. Pita, M.E. Pita Departamento de Ecología y Biología Animal, Facultad de Ciencias, Universidad de Vigo, Apdo.874, 36200 Vigo, Spain Authorship alphabetical: cite as BIOMAERL teamSearch for more papers by this authorA.A. Ramos-Esplá, A.A. Ramos-Esplá UMR CNRS 6539 Ecologie benthique, Institut Universitaire Européen de la Mer, Place Copernic, 29280 Plouzane, France Authorship alphabetical: cite as BIOMAERL teamSearch for more papers by this authorM. Rizzo, M. Rizzo Department of Biology, University of Malta, Msida, Malta Authorship alphabetical: cite as BIOMAERL teamSearch for more papers by this authorA. Sánchez-Mata, A. Sánchez-Mata Departamento de Ecología y Biología Animal, Facultad de Ciencias, Universidad de Vigo, Apdo.874, 36200 Vigo, Spain Departamento de Biología Animal, Facultad de Biología, Universidad de Santiago de Compostela, 15706 Santiago de Compostela, Galicia, Spain Authorship alphabetical: cite as BIOMAERL teamSearch for more papers by this authorA. Seva, A. Seva Ciencias Ambientales (Biología marina), Universidad de Alicante, 03080 Alicante, Spain Authorship alphabetical: cite as BIOMAERL teamSearch for more papers by this authorP.J. Schembri, P.J. Schembri Department of Biology, University of Malta, Msida, Malta Authorship alphabetical: cite as BIOMAERL teamSearch for more papers by this authorC. Valle, C. Valle Ciencias Ambientales (Biología marina), Universidad de Alicante, 03080 Alicante, Spain Authorship alphabetical: cite as BIOMAERL teamSearch for more papers by this author C. Barbera, C. Barbera Ciencias Ambientales (Biología marina), Universidad de Alicante, 03080 Alicante, Spain Authorship alphabetical: cite as BIOMAERL teamSearch for more papers by this authorC. Bordehore, C. Bordehore Ciencias Ambientales (Biología marina), Universidad de Alicante, 03080 Alicante, Spain Authorship alphabetical: cite as BIOMAERL teamSearch for more papers by this authorJ.A. Borg, J.A. Borg Department of Biology, University of Malta, Msida, Malta Authorship alphabetical: cite as BIOMAERL teamSearch for more papers by this authorM. Glémarec, M. Glémarec UMR CNRS 6539 Ecologie benthique, Institut Universitaire Européen de la Mer, Place Copernic, 29280 Plouzane, France Authorship alphabetical: cite as BIOMAERL teamSearch for more papers by this authorJ. Grall, J. Grall UMR CNRS 6539 Ecologie benthique, Institut Universitaire

Européen de la Mer, Place Copernic, 29280 Plouzane, France Authorship alphabetical: cite as BIOMAERL teamSearch for more papers by this authorJ. M. Hall-Spencer, J. M. Hall-Spencer School of Biological Sciences, University of Plymouth, Drake Circus, Plymouth, PL4 8AA, UK Authorship alphabetical: cite as BIOMAERL teamSearch for more papers by this authorCh. de la Huz, Ch. de la Huz Departamento de Ecología y Biología Animal, Facultad de Ciencias, Universidad de Vigo, Apdo.874, 36200 Vigo, Spain Authorship alphabetical: cite as BIOMAERL teamSearch for more papers by this authorE. Lanfranco, E. Lanfranco Department of Biology, University of Malta, Msida, Malta Authorship alphabetical: cite as BIOMAERL teamSearch for more papers by this authorM. Lastra, M. Lastra Departamento de Ecología y Biología Animal, Facultad de Ciencias, Universidad de Vigo, Apdo.874, 36200 Vigo, Spain Authorship alphabetical: cite as BIOMAERL teamSearch for more papers by this authorP.G. Moore, Corresponding Author P.G. Moore University Marine Biological Station, Millport, Isle of Cumbrae, UK, KA28 0EG Authorship alphabetical: cite as BIOMAERL team CoordinatorUniversity Marine Biological Station Millport, Isle of Cumbrae, KA28 0EG, UKSearch for more papers by this authorJ. Mora, J. Mora Departamento de Biología Animal, Facultad de Biología, Universidad de Santiago de Compostela, 15706 Santiago de Compostela, Galicia, Spain Authorship alphabetical: cite as BIOMAERL teamSearch for more papers by this authorM.E. Pita, M.E. Pita Departamento de Ecología y Biología Animal, Facultad de Ciencias, Universidad de Vigo, Apdo.874, 36200 Vigo, Spain Authorship alphabetical: cite as BIOMAERL teamSearch for more papers by this authorA.A. Ramos-Esplá, A.A. Ramos-Esplá UMR CNRS 6539 Ecologie benthique, Institut Universitaire Européen de la Mer, Place Copernic, 29280 Plouzane, France Authorship alphabetical: cite as BIOMAERL teamSearch for more papers by this authorM. Rizzo, M. Rizzo Department of Biology, University of Malta, Msida, Malta Authorship alphabetical: cite as BIOMAERL teamSearch for more papers by this authorA. Sánchez-Mata, A. Sánchez-Mata Departamento de Ecología y Biología Animal, Facultad de Ciencias, Universidad de Vigo, Apdo.874, 36200 Vigo, Spain Departamento de Biología Animal, Facultad de Biología, Universidad de Santiago de Compostela, 15706 Santiago de Compostela, Galicia, Spain Authorship alphabetical: cite as BIOMAERL teamSearch for more papers by this authorA. Seva, A. Seva Ciencias Ambientales (Biología marina), Universidad de Alicante, 03080 Alicante, Spain Authorship alphabetical: cite as BIOMAERL teamSearch for more papers by this authorP.J. Schembri, P.J. Schembri Department of Biology, University of Malta, Msida, Malta Authorship alphabetical: cite as BIOMAERL teamSearch for more papers by this authorC. Valle, C. Valle Ciencias Ambientales (Biología marina), Universidad de Alicante, 03080 Alicante, Spain Authorship alphabetical: cite as BIOMAERL teamSearch for more papers by this author First published: 12 February 2003 <https://doi.org/10.1002/aqc.569>Citations: 142 Authorship alphabetical: cite as BIOMAERL team Coordinator AboutPDF ToolsRequest permissionExport citationAdd to favoritesTrack citation ShareShare Give accessShare full text accessShare full-text accessPlease review our Terms and Conditions of Use and check box below to share full-text version of article.I have read and accept the Wiley Online Library Terms and Conditions of UseShareable LinkUse the link below to share a full-text version of this article with your friends and colleagues. Learn more.Copy URL Share a linkShare onEmailFacebookTwitterLinkedInRedditWechat

Abstract 1. Maerl beds occur worldwide and are formed by an accumulation of unattached calcareous red algae (Rhodophyta). 2. Maerl-forming algae grow in a superficial living layer on sediments within the photic zone. 3. Maerl beds are spatially complex habitats with a high degree of species and trophic group diversity. 4. The European Commission's 'Habitats Directive' mandates the conservation management of two of the main European maerl-forming species, *Phymatolithon calcareum* and *Lithothamnion corallioides*. 5. Mediterranean maerl beds are to be considered for inclusion in national inventories of sites of conservation interest, as required by the SPABIM Protocol of the Barcelona Convention. 6. In spite of their importance, and the requirement for their conservation management, European maerl grounds suffer a variety of anthropogenic perturbations including direct exploitation through extraction, fishing impacts and chemical pollution by organic matter and excess nutrients. 7. The ecology of northeast Atlantic and Mediterranean maerl beds has received little attention, in contrast to other marine communities (e.g. kelp forests, sea-grass meadows). 8. Key conservation and management measures proposed include: the recognition that maerl beds are non-renewable resources and cannot sustain direct exploitation; prohibitions on the use of towed gear on maerl grounds; moratoria on the issue of further permits for the siting of aquaculture units above maerl grounds; monitoring of existing exploited or impacted maerl beds; the designation of 'no-take' reserves; measures to limit the impacts that might affect water quality above maerl beds; a programme of monitoring of the 'health' of European maerl beds; an awareness campaign on the biological importance of maerl beds; a higher conservation status for maerl habitats and maerl-forming species in European legislation; and further research on maerl ecosystems. Copyright © 2003 John Wiley & Sons, Ltd.

REFERENCES Ballesteros E. 1989. Composición y estructura de los fondos de maerl de Tossa de Mar (Gerona, España) *Collectanea Botanica* (Barcelona) 17 (2): 161?182. Basso D. 1996. Adaptive strategies and convergent morphologies in some Mediterranean coralline algae. In Cherchi, A. (ed.); *Autecology of selected fossil organisms: achievements and problems*. *Bollettino della Societa Paleontologica Italiana* 3, 1?8. Bellan-Santini D, Lacaze JC, Poizat

C. 1994. Les biocénoses marines et littorales de Méditerranée: synthèse, menaces et perspectives. Museum National d'Histoire Naturelle: Paris: 246 pp.

BIOMAERL Team. 1998. Maerl grounds: habitats of high biodiversity in European waters, In Proceedings of the third European Marine Science and Technology Conference, Project Synopses, vol. I, Marine Ecosystems: Lisbon; 169?178.

BIOMAERL Team. 1999. Final Report, BIOMAERL project (Coordinator: P.G.Moore, University Marine Biological Station Millport, Scotland), EC Contract No.MAS3-CT95-0020, 1: 1?541, 2: 542-973pp. + Appendix.

Birkett DA, Maggs CA, Dring MJ. 1998. An Overview of Dynamic and Sensitivity Characteristics for Conservation Management of Marine SACs, vol. 5. Maerl, [UK Marine SACs Project], Scottish Association for Marine Science, Scotland, 116 pp.

Blunden G, Farnham WF, Jephson N, Barwell CJ, Fenn RH, Plunkett BA. 1981. The composition of maerl beds of economic interest in northern Brittany, Cornwall and Ireland. In Proceedings of the Xth International Seaweed Symposium. Walter de Gruyter & Co.: Berlin; 651?656.

Blunden G, Farham WF, Jephson N, Fenn RH, Plunkett BA. 1977. The composition of maerl from the Glenan Islands of Southern Brittany *Botanica Marina* 20: 121?125.

Bohsanck JA, Ault JS. 1996. Management strategies to conserve marine biodiversity. *Oceanography* 9 (1): 73?82.

Bordehore C, Barberá C, Mallol J, Martínez MC, Mena C, Patiño JL, Valle C, Vivas M, Ramos-Esplá AA. 1999. Algal composition and habitat structure in two maerl bottoms: marine reserve of Tabarca and the Island of Benidorm. Proceedings of the First International Workshop on Marine Reserves, Murcia.

Bordehore C, Borg JA, Lanfranco E, Ramos-Esplá AA, Rizzo M, Schembri P. 2000. Trawling as a major threat to Mediterranean maerl beds, presented at the First Mediterranean Symposium on Marine Vegetation, Regional Activity Centre for Specially Protected Areas [UNEP Mediterranean Action Plan], Ajaccio, Corsica, France 2?3 October 2000.

Borg JA, Lanfranco E, Mifsud JR, Rizzo M, Schembri PJ. 1998. Does fishing have an impact on Maltese maerl grounds? ICES Conference on the Ecosystem Effects of Fishing, Iraklion, Crete.

Bosence DWJ. 1979. Live and dead faunas from coralline algal gravels, Co. Galway. *Palaeontology* 19: 365?395.

Bosence DWJ. 1983. The occurrence and ecology of recent rhodoliths ? a review. In Coated Grains, TM Peryt (Ed.) Springer Verlag: Berlin; 225?242.

Brenchley GA. 1981. Disturbance and community structure: an experimental study of bioturbation in marine soft-bottom environments. *Journal of Marine Research* 39: 767?790.

Briand X. 1989. Le Lithothamne: tradition d'hier et agrochimie de demain. *Oceanis* 15: 693?739.

Cabioch L. 1968. Contribution à la connaissance des peuplements benthiques de la Manche occidentale. *Cahiers de Biologie Marine* 9 (suppl 5.): 493?720.

Cabioch J. 1969. Les fonds de maerl de la baie de Morlaix et leur peuplement végétal. *Cahiers de Biologie Marine* 10, 139?161.

Canals M, Ballesteros E. 1997. Production of carbonate particles by phytobenthic communities on the Mallorca-Menorca shelf, northwestern Mediterranean Sea. *Deep Sea Research* 44: 611?629.

Council of Europe. 1996. Convention relative à la conservation de la vie sauvage et du milieu naturel de l'Europe [Rapport T-PVS (96) 102], Council of Europe, Strasbourg, 99 pp.

Davies CE, Moss D. 1998. EUNIS habitat classification: final draft with further revisions to marine habitats. [Work Programme Task 7.5.1], Unpublished report to the European Environment Agency, European Topic Centre on Nature Conservation.

De Grave S, Fazakerley H, Kelly L, Guiry MD, Ryan M, Walshe J. 2000. A study of selected maerl beds in Irish waters and their potential for sustainable extraction [Marine Resource Series 10]. Marine Institute, Dublin, 44 pp.

De Grave S, Whitaker A. 1999. Benthic community re-adjustment following dredging of a muddy-maerl matrix. *Marine Pollution Bulletin* 38 (2): 102?108.

Engle VD, Summers JK. 1999. Refinement, validation, and application of a Benthic Condition Index for Northern Gulf of Mexico Estuaries. *Estuaries* 22 (3A): 624?635.

European Commission. 1999. Interpretation manual of European Union habitats EUR15/2, European Commission DG Environment, Brussels, 119 pp.

Falconetti C. 1970. Etude faunistique d'un faciès: 'La gravelette' ou maerl de Castiglione (Algérie). *Tethys* 1(4): 1057?1096.

Freiwald A, Henrich R. 1994. Reefal coralline algal build-ups within the Arctic Cycle: morphology and sedimentary dynamics under extreme environment seasonality. *Sedimentology* 41: 963?984.

Grall J, Glémarec M. 1997a. Using biotic indices to estimate macrobenthic community perturbations in the Bay of Brest. *Estuarine, Coastal and Shelf Science* 44 (Suppl. A): 43?53.

Grall J, Glémarec M. 1997b. Biodiversité des fonds de maerl en Bretagne: approche fonctionnelle et impacts anthropogéniques. *Vie et Milieu* 47(4): 339?349.

Grall J, Hall-Spencer JM. 2003. Problems facing maerl conservation in Brittany. *Aquatic Conservation: Marine and Freshwater Ecosystems*. 13: S55?S64.

Hall-Spencer JM. 1998. Conservation issues relating to maerl beds as habitats for mollusks. *Journal of Conchology, Special Publication* 2: 271?286.

Hall-Spencer JM, Grall J, Moore PG, Atkinson RJA. 2003. Bivalve fishing and maerl bed conservation in France and the UK?retrospect and prospect. *Aquatic Conservation: Marine and Freshwater Ecosystem* 13: S33?S41.

Hall-Spencer JM, Moore PG. 2000a. Scallop dredging has profound, long-term impacts on maerl habitats. *ICES Journal of Marine Science* 57: 1407?1415.

Hall-Spencer JM, Moore PG. 2000b. Impacts of scallop dredging on maerl grounds. In *Effects of Fishing on Non-Target Species and Habitats: Biological, Conservation and Socio-Economic Issues*. MJ Kaiser, SJ de Groot. (eds). Blackwell Science: Oxford; 105?117.

Irvine LM, Chamberlain YMC. 1994. Seaweeds of the British Isles, vol. 1, Part 2B. British Museum (Natural History): London, 276 pp.

Jacquotte R. 1962. Etude des fonds de maerl de Méditerranée. *Recueils des travaux de la Station marine d'Endoume* 26 (41): 141?235.

Keegan BF. 1974. The macrofauna of maerl substrates

on the west coast of Ireland. *Cahiers de Biologie Marine* 15: 513-530. Lanfranco E, Rizzo M, Hall-Spencer J, Borg JA, Schembri PJ. 1999. Maerl-forming coralline algae and associated phytobenthos from the Maltese Islands. *The Central Mediterranean Naturalist* 3 (1): 1-6. Littler MM, Littler DS, Hanisack MD. 1991. Deep-water rhodolith distribution, productivity, and growth history at sites of formation and subsequent degradation. *Journal of Experimental Marine Biology and Ecology* 150: 163-182. Massutí E, Reñones O, Carbonell A, Oliver P. 1996. Demersal fish communities exploited on the continental shelf and slope off Majorca (Balearic Islands, NW Mediterranean). *Vie et Milieu* 46 (1): 45-55. Mora J. 1980. Poblaciones bentónicas de la Ría de Arosa, Tesis Doctoral. Universidad de Santiago, 335 pp. Norse EA (ed.), 1993. *Global Marine Biological Diversity: a Strategy for Building Conservation into Decision Making*. Island Press: Washington; 384 pp. Oliver P. 1983. Les ressources halieutiques de la Méditerranée, Première Partie: Méditerranée Occidentale. *Studies and Reviews of the General Fisheries Council of the Mediterranean* 59: 135. OSPAR. 1998. OSPAR strategy on the protection and conservation of the ecosystems and biological diversity of the maritime area [Reference number: 1998-19], OSPAR Convention for the Protection of the Marine Environment of the North-East Atlantic, Ministerial Meeting of the OSPAR Commission, Sintra: 22-23 July 1998; 3pp. Parsons TR. 1992. Biological coastal communities: productivity and impacts, in *Coastal systems studies and sustainable development*. UNESCO Technical Papers in Marine Science 64: 27-37. Picard J. 1965. Recherches qualitatives sur les biocoenoses marines des substrats meubles dragables de la région marseillaise. *Recueils des travaux de la Station marine d'Endoume* 31 (47): 5-137. Potin P, Floc'h JY, Augris C, Cabioch J. 1990. Annual growth rate of the calcareous red alga *Lithothamnion corallioides* (Corallinales, Rhodophyta) in the Bay of Brest, France. *Hydrobiologia* 204/205: 263-267. Ramos-Esplá AA. 1992. *Ascidias litorales del Mediterráneo ibérico: faunística, ecología y biogeografía*. Universidad de Alicante, Alicante, 410 pp. Ramos-Esplá AA, Guillén JE, Bayle JT, Sánchez-Jérez P. 2000. Artificial antitrawling reefs off Alicante, South-Eastern Iberian Peninsula: evolution of reefs block and set design. In *Artificial Reefs in European Waters* A Jensen, K Collins (eds). Kluwer Academic Publishers: London; 195-217. Ramos-Esplá AA. 1990. The marine reserve of Nueva Tabarca (Alicante, Spain). Management aspects. In *Parchi marini del Mediterraneo. Aspetti naturalistici e gestionali*. A Cossu, V Gazale, I Milella (eds). ICM/Università di Sassari/CO.RI.SA: San Teodoro; 107-117. Rijnsdorp AD, Buijs AM, Storbeck F, Visser EG. 1998. Micro-scale distribution of beam trawl effort in the southern North Sea between 1993 and 1996 in relation to the trawling frequency of the seabed and the impact on benthic organisms. *ICES Journal of Marine Science* 55: 403-419. Rowe GA, Shearer M, Jensen AC. 1990. The infauna of Handfast Point maerl bed, [Report No. SUDO/TEC/90/16C] Nature Conservancy Council, Peterborough, 57pp. Sánchez-Mata A, Abella FE, Mora J. 1998. Biodiversidad específica de los fondos de maerl *Lithothamnion corallioides* (P. & H. Crouan) P. & H. Crouan y *Phymatholithon calcareum* (Pallas) Adey & McKibbin de la Ría de Vigo. In *Proceedings of the 10th Simposio Ibérico de Estudios del Bentos Marino*, Universidade do Algarve, Praia de Carvoeiro; 168-169. Sánchez-Mata A, Lastra M, de la Huz Ch, Pita ME, Mora J. 1999. The maerl grounds as islands of high biodiversity in temperate latitudes: the Galician case. In *Proceedings of the 34th European Marine Biology Symposium*, Universidade dos Açores. Ponta Delgada; 77-78. Sánchez-Lizaso JL, Guillén JE, Ramos-Esplá AA. 1990. The regression of *Posidonia oceanica* meadows in El Campello. *Rapports C.I.E.S.M.* 32(1): 7. Seaman W, Sprague LM. 1991. Artificial habitat practices in aquatic systems. In *Artificial Habitats for Marine and Freshwater Fisheries*, W Seaman, LM Sprague (eds). Academic Press: London; 1-29. Smith VH, Tilman GD, Nekola JC. 1999. Eutrophication: impacts of excess nutrient inputs on freshwater, marine, and terrestrial ecosystems. *Environmental Pollution* 100: 179-196. Sousa WP. 1984. The role of disturbance in natural communities. *Annual Review of Ecology and Systematics* 15: 353-391. Steller DL, Foster MS. 1995. Environmental factors influencing distribution and morphology of rhodoliths in Bahía Concepción, B.C.S., México. *Journal of Experimental Marine Biology and Ecology* 194: 201-212. Steller DL, Riosmena-Rodríguez R, Foster MS, Roberts CA. 2003. Rhodolith bed diversity in the Gulf of California: the importance of rhodolith structure and consequences of anthropogenic disturbances. *Aquatic Conservation: Marine and Freshwater Ecosystems* 13(Suppl 1): S5-S20. Thouzeau G. 1991. Experimental collection of postlarvae of *Pecten maximus* (L.) and other benthic macrofaunal species in the Bay of Saint-Brieuc, France. II. Reproduction patterns and postlarval growth of five mollusc species. *Journal of Experimental Marine Biology and Ecology* 148: 181-200. UNEP/IUCN/GIS Posidonie. 1990. Livre rouge 'Gérard Vuignier' des végétaux, peuplements et paysages marins menacés de Méditerranée. MAP Technical Report Series 43; United Nations Environment Programme, Athens, 250 pp. Citing Literature Volume13, IssueS1Supplement: International Workshop on 'The Conservation and Management of Maerl', 23-27 February, 2001. University Marine Biological Station, Millport, Isle of Cumbrae, Scotland January/February 2003 Pages S65-S76

ReferencesRelatedInformation

SOURCE: Aquatic conservation

PDF URL: None

CITED BY COUNT: 182

PUBLICATION YEAR: 2003

TYPE: article

CONCEPTS: ['Ecology', 'Habitat', 'Fishery', 'Environmental science', 'Biology']