

ID: W1967849499

TITLE: Antibiotics in typical marine aquaculture farms surrounding Hailing Island, South China: Occurrence, bioaccumulation and human dietary exposure

AUTHOR: ['Hui Chen', 'Shan Liu', 'Xiangrong Xu', 'Shuangshuang Liu', 'Guang?Jie Zhou', 'Kun Sun', 'Jian?Liang Zhao', 'Guang?Guo Ying']

ABSTRACT:

The occurrence, bioaccumulation, and human dietary exposure via seafood consumption of 37 antibiotics in six typical marine aquaculture farms surrounding Hailing Island, South China were investigated in this study. Sulfamethoxazole, salinomycin and trimethoprim were widely detected in the water samples (0.4?36.9 ng/L), while oxytetracycline was the predominant antibiotic in the water samples of shrimp larvae pond. Enrofloxacin was widely detected in the feed samples (16.6?31.8 ng/g) and erythromycin?H₂O was the most frequently detected antibiotic in the sediment samples (0.8?4.8 ng/g). Erythromycin?H₂O was the dominant antibiotic in the adult *Fenneropenaeus penicillatus* with concentrations ranging from 2498 to 15,090 ng/g. In addition, trimethoprim was found to be bioaccumulative in young *Lutjanus russelli* with a median bioaccumulation factor of 6488 L/kg. Based on daily intake estimation, the erythromycin?H₂O in adult *F. penicillatus* presented a potential risk to human safety.

SOURCE: Marine pollution bulletin

PDF URL: None

CITED BY COUNT: 254

PUBLICATION YEAR: 2015

TYPE: article

CONCEPTS: ['Bioaccumulation', 'Enrofloxacin', 'Erythromycin', 'Aquaculture', 'Oxytetracycline', 'Biology', 'Trimethoprim', 'Florfenicol', 'Antibiotics', 'Fishery', 'Veterinary medicine', 'Ecology', 'Microbiology', 'Fish <Actinopterygii>', 'Ciprofloxacin', 'Medicine']