

ID: W2055667516

TITLE: Primary production of prochlorophytes, cyanobacteria, and eucaryotic ultraphytoplankton: Measurements from flow cytometric sorting

AUTHOR: ['William K. W. Li']

ABSTRACT:

A partitioning of ultraphytoplankton primary production among prochlorophytes, cyanobacteria, and eucaryotic algae was made by shipboard flow cytometric sorting of ^{14}C -labeled cells. Aggregate primary production was derived from the sum, over all three ultraplankton groups, of the product of cell abundance and cell-specific rate of ^{14}C uptake which ranged from 0.03 to 4 fg C cell $^{-1}$ h $^{-1}$ for prochlorophytes and 0.2 to 10 fg C cell $^{-1}$ h $^{-1}$ for cyanobacteria. Results indicated that the dominant primary producer was not necessarily the numerical dominant nor necessarily the group with the highest cell-specific rate of ^{14}C uptake. Generally, eucaryotic ultraphytoplankton are dominant because of their high cell-specific rate of ^{14}C uptake and in spite of their relatively low abundance. Less often, it seems, procaryotic picoplankton may dominate in spite of their low cell-specific rate of ^{14}C uptake because of their high abundance.

SOURCE: Limnology and oceanography/The oel & o on cd-rom

PDF URL: <https://onlinelibrary.wiley.com/doi/pdfdirect/10.4319/lo.1994.39.1.0169>

CITED BY COUNT: 475

PUBLICATION YEAR: 1994

TYPE: article

CONCEPTS: ['Cyanobacteria', 'Picoplankton', 'Abundance (ecology)', 'Algae', 'Biology', 'Cell sorting', 'Flow cytometry', 'Cell', 'Botany', 'Chemistry', 'Biochemistry', 'Ecology', 'Molecular biology', 'Bacteria', 'Genetics']