

ID: W2046794716

TITLE: An interlaboratory study of dissolved oxygen in water

AUTHOR: ['Robert J. Wilcock', 'C. Stevenson', 'Cindy A. Roberts']

ABSTRACT:

Abstract Dissolved oxygen concentrations in water samples free from chemical reductants may be stabilised by the addition of mercuric chloride ( $40 \text{ g m}^{-3}$ ) and storage in gas-tight bottles. This preservation technique has been used in an interlaboratory study of dissolved oxygen analysis in New Zealand laboratories. At reference concentrations of 1.20 and  $5.86 \text{ g m}^{-3}$ , there was a significant positive bias in results reported for both the Winkler method (0.24 and  $0.22 \text{ g m}^{-3}$  respectively) and the membrane electrode method (0.59 and  $0.62 \text{ g m}^{-3}$  respectively). Inadequate precautions to avoid sample aeration during handling and analysis probably caused the bias.

SOURCE: Water research

PDF URL: None

CITED BY COUNT: 22

PUBLICATION YEAR: 1981

TYPE: article

CONCEPTS: ['Environmental science', 'Environmental chemistry', 'Oxygen', 'Water quality', 'Environmental engineering', 'Hydrology (agriculture)', 'Chemistry', 'Engineering', 'Ecology', 'Biology', 'Organic chemistry', 'Geotechnical engineering']