

ID: W2618878949

TITLE: Organophosphate Esters in Air, Snow, and Seawater in the North Atlantic and the Arctic

AUTHOR: ['Jing Li', 'Zhiyong Xie', 'Wenying Mi', 'Shengli Lai', 'Chongguo Tian', 'Kay?Christian Emeis', 'Ralf Ebinghaus']

ABSTRACT:

The concentrations of eight organophosphate esters (OPEs) have been investigated in air, snow and seawater samples collected during the cruise of ARK-XXVIII/2 from sixth June to third July 2014 across the North Atlantic and the Arctic. The sum of gaseous and particle concentrations (Σ OPE) ranged from 35 to 343 pg/m³. The three chlorinated OPEs accounted for $88 \pm 5\%$ of the Σ OPE. The most abundant OPE was tris(2-chloroethyl) phosphate (TCEP), with concentrations ranging from 30 to 227 pg/m³, followed by three major OPEs, such as tris(1-chloro-2-propyl) phosphate (TCPP, 0.8 to 82 pg/m³), tri-n-butyl phosphate (TnBP, 2 to 19 pg/m³), and tri-iso-butyl phosphate (TiBP, 0.3 to 14 pg/m³). The Σ OPE concentrations in snow and seawater ranged from 4356 to 10561 pg/L and from 348 to 8396 pg/L, respectively. The atmospheric particle-bound dry depositions of TCEP ranged from 2 to 12 ng/m²/day. The air-seawater gas exchange fluxes were dominated by net volatilization from seawater to air for TCEP (mean, 146 ± 239 ng/m²/day), TCPP (mean, 1670 ± 3031 ng/m²/day), TiBP (mean, 537 ± 581 ng/m²/day) and TnBP (mean, 230 ± 254 ng/m²/day). This study highlighted that OPEs are subject to long-range transport via both air and seawater from the European continent and seas to the North Atlantic and the Arctic.

SOURCE: Environmental science & technology

PDF URL: None

CITED BY COUNT: 146

PUBLICATION YEAR: 2017

TYPE: article

CONCEPTS: ['Seawater', 'TCEP', 'Snow', 'Organophosphate', 'Environmental chemistry', 'Arctic', 'Phosphate', 'Chemistry', 'Oceanography', 'Pesticide', 'Meteorology', 'Geology', 'Ecology', 'Organic chemistry', 'Biology', 'Physics', 'Phosphine', 'Catalysis']