

ID: W2083956291

TITLE: Seamount benthos in a cobalt-rich crust region of the central Pacific: conservation challenges for future seabed mining

AUTHOR: ['Thomas A. Schlacher', 'Amy R. Baco', 'Ashley A. Rowden', 'Timothy D. O'Hara', 'Malcolm R. Clark', 'Chris Kelley', 'John F. Dower']

ABSTRACT:

**Abstract Aim** The benthic fauna of seamounts typically includes organisms that are slow-growing, long-lived and sensitive to mechanical disturbance, making seamounts susceptible to anthropogenic impacts. Such impacts may arise from mining cobalt-rich crusts, envisaged for seamounts in the central North Pacific; this scenario requires that environmental guidelines for mining operations on seamounts be developed. Here, we provide the biological information essential for effective conservation planning of deep-sea features targeted for such mining. **Location** Central North Pacific, Hawaiian Seamount Chain. **Methods** Spatial analysis of seamount benthos using a large biological dataset (> 600 taxa) obtained from 144 submersible dives (depth range: 113–1985 m) on 27 seamounts covering a distance of over 2200 km of ocean. **Results** Benthic assemblages of invertebrates are structurally different between seamounts located inside and outside a region with cobalt-rich crusts. This spatial contrast results from variations in species composition and relative abundance of species, rather than differences in species richness, challenging historical notions of an impoverished cobalt-rich crust fauna in the region. Seamount assemblages also have high species turnover with depth and distance at the scale of individual seamounts, but geographic separation was a poor predictor of ecological separation for the region at large. **Main conclusion** Several implications for the design of spatial management and conservation tools with respect to mining emerge: (1) conservation of seamounts outside the cobalt-rich crust region is unlikely to capture the full range of ecological features found inside the region; (2) conservation areas need to encompass a broad bathymetric gradient; (3) ideally, mining blocks on individual seamounts should not exceed 2 km in length. Overall, the life history characteristics and morphological traits of the deep-water invertebrate fauna typical of seamounts in the region imply that any recovery from mechanical impacts is likely to be very slow.

SOURCE: Diversity and distributions

PDF URL: None

CITED BY COUNT: 86

PUBLICATION YEAR: 2013

TYPE: article

CONCEPTS: ['Seamount', 'Benthos', 'Benthic zone', 'Fauna', 'Geology', 'Crust', 'Range (aeronautics)', 'Species richness', 'Oceanography', 'Habitat', 'Ecology', 'Geography', 'Paleontology', 'Biology', 'Materials science', 'Composite material']