

ID: W2132644839

TITLE: Cholera and Other Types of Vibriosis: A Story of Human Pandemics and Oysters on the Half Shell

AUTHOR: ['J. Glenn Morris', 'David W. K. Acheson']

ABSTRACT:

Vibrios are ubiquitous in the aquatic environment and are commonly present in or on shellfish and other seafood. A small subset of strains/species are able to cause human disease, including the cholera toxin-producing strains of *Vibrio cholerae* that are responsible for epidemic/pandemic cholera; thermostable direct hemolysin-producing strains of *Vibrio parahaemolyticus*; and *Vibrio vulnificus*, which can cause fulminant sepsis. Cholera outbreaks can be initiated by transmission of "epidemic" *V. cholerae* strains from their environmental reservoir to humans through seafood or other environmentally related food or water sources. "Nonepidemic" strains of *V. cholerae* and strains of other *Vibrio* species, including *V. parahaemolyticus* and *V. vulnificus*, are generally acquired by eating seafood (particularly raw oysters/oysters on the half shell). Although the primary clinical manifestation of infection with these strains is gastroenteritis, they can also cause wound infections and (particularly for *V. vulnificus*) septicemia in persons who have liver disease or are immunocompromised.

SOURCE: Clinical infectious diseases/Clinical infectious diseases (Online. University of Chicago. Press)

PDF URL: <https://academic.oup.com/cid/article-pdf/37/2/272/6104314/37-2-272.pdf>

CITED BY COUNT: 271

PUBLICATION YEAR: 2003

TYPE: article

CONCEPTS: ['*Vibrio vulnificus*', '*Vibrio parahaemolyticus*', '*Vibrio cholerae*', 'Cholera', 'Microbiology', 'Vibrio Infections', 'Outbreak', 'Vibrionaceae', 'Pandemic', 'Vibrio', 'Biology', 'Food poisoning', 'Virology', 'Medicine', 'Bacteria', 'Disease', 'Infectious disease (medical specialty)', 'Pathology', 'Genetics', 'Coronavirus disease 2019 (COVID-19)']