

ID: W2142113544

TITLE: Successful high-level accumulation of fish oil omega-3 long-chain polyunsaturated fatty acids in a transgenic oilseed crop

AUTHOR: ['Noemí Ruiz-López', 'Richard P. Haslam', 'Johnathan A. Napier', 'Olga Sayanova']

ABSTRACT:

Omega-3 (also called n-3) long-chain polyunsaturated fatty acids ( $C_{20}$ ; LC-PUFAs) are of considerable interest, based on clear evidence of dietary health benefits and the concurrent decline of global sources (fish oils). Generating alternative transgenic plant sources of omega-3 LC-PUFAs, i.e. eicosapentaenoic acid (20:5 n-3, EPA) and docosahexaenoic acid (22:6 n-3, DHA) has previously proved problematic. Here we describe a set of heterologous genes capable of efficiently directing synthesis of these fatty acids in the seed oil of the crop *Camelina sativa*, while simultaneously avoiding accumulation of undesirable intermediate fatty acids. We describe two iterations: RRes\_EPA in which seeds contain EPA levels of up to 31% (mean 24%), and RRes\_DHA, in which seeds accumulate up to 12% EPA and 14% DHA (mean 11% EPA and 8% DHA). These omega-3 LC-PUFA levels are equivalent to those in fish oils, and represent a sustainable, terrestrial source of these fatty acids. We also describe the distribution of these non-native fatty acids within *C. sativa* seed lipids, and consider these data in the context of our current understanding of acyl exchange during seed oil synthesis.

SOURCE: Plant journal

PDF URL: <https://onlinelibrary.wiley.com/doi/pdfdirect/10.1111/tpj.12378>

CITED BY COUNT: 284

PUBLICATION YEAR: 2013

TYPE: article

CONCEPTS: ['Polyunsaturated fatty acid', 'Docosahexaenoic acid', 'Camelina sativa', 'Eicosapentaenoic acid', 'Fish oil', 'Food science', 'Biology', 'Context (archaeology)', 'Crop', 'Fatty acid', 'Chemistry', 'Biochemistry', 'Fish <Actinopterygii>', 'Agronomy', 'Fishery', 'Paleontology']