

ID: W2142731244

TITLE: Effects of seismic energy releases on the survival and development of zoeal larvae of dungeness crab (*Cancer magister*)

AUTHOR: ['Walter H. Pearson', 'John R. Skalski', 'Stephen D. Sulkin', 'Charles I. Malme']

ABSTRACT:

In blind, controlled field experiments, early Stage II zoeae of Dungeness crab (*Cancer magister* DANA) were exposed to sounds from single discharges of a 13.8-litre array of seven air guns. Their survival and development were followed during subsequent laboratory culture. Immediate mortality was low (0 to 2%) and showed no significant difference between control and exposed larvae ( $P > 0.05$ ). Across all treatments and blocks of the experiment, survival to the molt to Stage III averaged 88.8%. The conditional Stage IV survival rate averaged 69.8%. The times to the molts to Stage III and Stage IV averaged 14.4 and 34.9 days, respectively. For immediate and long-term survival and time to molt, the field experiment revealed no statistically significant ( $P > 0.05$ ) effects on zoeae for exposures as close as 1 m from the array, nor for mean sound pressure as high as 231 dB re 1  $\mu$ Pa and cumulative energy density up to 251 J/M<sup>2</sup>. Post hoc power calculations showed that any reduction in zoeal survival as a result of sound exposure was less than 7% for survival to Stage III and less than 12% for Stage IV conditional survival ( $1 - P = 0.90$ ,  $P = 0.05$  one-tailed). The sound exposures in our study were at the maximum levels likely to be experienced by a zoea during an actual survey.

SOURCE: Marine environmental research

PDF URL: None

CITED BY COUNT: 28

PUBLICATION YEAR: 1994

TYPE: article

CONCEPTS: ['Larva', 'Stage (stratigraphy)', 'Biology', 'Energy density', 'Animal science', 'Ecology', 'Physics', 'Paleontology', 'Theoretical physics']