

Room-Temperature Superconductivity as the Developmental Terminal State of a Generalized Adaptive Intelligent System: The Topological Condensation of a Dynamic Equilibrium Quantum Manifold

Baoliin (Zaitian) Wu 伍宝林

Contents

1	Physical Proof of Existence	2
2	Room-Temperature Superconductivity as the Developmental Terminal State of a Generalized Adaptive Intelligent System: The Topological Condensation of a Dynamic Equilibrium Quantum Manifold	3
2.1	Abstract	3
2.2	1. Introduction: Beyond Electron Pairing	4
2.3	2. The Cosmic Engineering First Principle: 1+4 Meta-Structure and the Gui-Ju Dao	5
2.4	3. The Dynamic Equilibrium Quantum Manifold and the Tensor AID Equation . . .	5
	2.4.1 3.1 The Quantum Geometric Field as Tensor Order Parameter	5
	2.4.2 3.2 The Free Energy Functional for the Quantum Geometric Field	6
2.5	3.5 Quantum Geometric Field Reconstruction: From Atomic Coordinates to Mani- fold Condensation	6
	2.5.1 3.5.0 Paradigm Statement: From “Material Screening” to “Geometric Field Reconstruction”	6
	2.5.2 3.5.1 Spatial-Omics Metric Tensor Extraction	7
	2.5.3 3.5.2 Dynamical Equations of the Developmental Path	9
	2.5.4 3.5.3 Numerical Validation: LaH as a Prototype	9
2.6	3.6 MES Scaling Laws and the Renormalization of Coupling Speed and Effective Inertia	10
	2.6.1 3.6.1 The Modified Einstein Spherical Scaling Laws	10
	2.6.2 3.6.2 Redefinition of c and m in the Superconducting DEQM	11
	2.6.3 3.6.3 Implications for “Acupoints” and “Gravity Wells”	12
	2.6.4 3.6.4 MES Scaling Laws and the Design Principle of “Deep Anchor”	12
	2.6.5 3.6.5 MES Scaling Laws and the Cosmic Curvature Constraint	13
2.7	4. The ZQP Mechanism: Physical Origin of Manifold Condensation	13
2.8	5. Algebraic Foundation: $\mathbb{Z}_5$ Symmetry and Spectral Duality	14
2.9	6. Developmental Morse Bifurcation: Quantum Manifold Embryogenesis	15
	2.9.1 6.1 Wugu Locking as the Terminal Bifurcation	15
2.10	7. Apoptotic Clearance: Maintaining Manifold Coherence Purity	15
2.11	8. Spatial-Omics Metric Tensor Extraction: The Geometry of the Condensed Manifold	16

2.11.1	8.1 Acupoints as Negative Curvature Extrema with Maximum ZQP Coupling	16
2.11.2	8.2 Dynamical Stability vs. Curvature Sign	16
2.12	9. Nonlocal Coupling: Anatomy Distal but State-Space Proximal	16
2.13	10. Topological Protection as Structural Stability	17
2.13.1	10.1 Topological Origin of Flux Quantization: $\Phi_0 = h/2e$ from $\mathbb{Z}_5$ Dual Winding	17
2.14	11. Anomalous Hall Effect as the Measurement Signal and Topological Probe	19
2.15	12. Landscape Reprogramming: From Confrontation to Reprogramming	19
2.16	13. Falsifiable Predictions	20
2.17	14. The T0-T6 Engineering Roadmap	20
2.18	15. Design Principles: From Material Screening to System Development	21
2.19	16. Conclusion	21
2.20	Acknowledgments	22
2.21	Funding	23
2.22	Availability of data and material	23
2.23	Competing interests	23
2.24	Authors Contributions	23
2.25	References	23
2.26	Physical Proof of Existence (Appendix)	26

1 Physical Proof of Existence

Title: Room-Temperature Superconductivity as the Developmental Terminal State of a Generalized Adaptive Intelligent System: The Topological Condensation of a Dynamic Equilibrium Quantum Manifold

Author: Baoliin (Zaitian) Wu 伍宝林

Date: 2026-09-16

Banknote Serial: DW58553911

Paper SHA-256: 66bef64d5ee3a4c2692665819fb74777e08d0f4a44890db0e4ba09490d2b30c9

Verification URL: <https://Zaitian001.github.io/-Self-Preprint-V1.1-/preprints/DW58553911.html>

This proof is automatically generated by the Self-Preprint system.

© 2026 Baoliin (Zaitian) Wu 伍宝林 • Decentralized Academic Archive

2 Room-Temperature Superconductivity as the Developmental Terminal State of a Generalized Adaptive Intelligent System: The Topological Condensation of a Dynamic Equilibrium Quantum Manifold

Baoliin (Zaitian) Wu 伍宝林

ORCID: 0009-0003-2051-5247

Email: zaitian001@gmail.com

Affiliation: People's Government of Guangdong Province, Guangzhou, CN

Date: September 16, 2026

2.1 Abstract

Room-temperature superconductivity remains one of the most sought-after goals in condensed matter physics, yet the dominant paradigm—material screening within BCS-derived architectures—has failed to deliver a viable path. Here we propose a fundamental redefinition: **room-temperature superconductivity is the developmental terminal state of a Generalized Adaptive Intelligent System (GAIS)—the topological condensation of a dynamic equilibrium quantum manifold (DEQM)**. This manifold is a **unified 3D quantum geometric field** $\mathcal{G}_{\mu\nu}(x, t)$, maintained through the **ZQP mechanism** in a sustained **zeroing measurement** cycle, protected by topological invariants, driven by nonlocal coupling, and directly probed by the **anomalous Hall effect (AHE)**. The superconducting state is not the pairing of electrons but the global topological ordering of the quantum geometric field. We establish a complete computable pipeline that maps discrete atomic coordinates to the continuous quantum geometric field:

$\mathbf{R}_I \xrightarrow{\text{DFPT}} \Omega_{\mu\nu}(\mathbf{k}) \xrightarrow{\text{ZQP}} g_{\mu\nu}(\mathbf{k}) \xrightarrow{\text{Morse}} \mathcal{G}_{\mu\nu}(\mathbf{x}, t)$. The algebraic foundation is the cyclic group $\mathbb{Z}_5$, whose generative operator S and destructive operator $K = S^2$ satisfy the spectral duality $\sigma(K) = \sigma(S)^2$. The physical mechanism is the **Zaitian Quantum Power (ZQP)** coupling, previously validated through V4.52 numerical simulations, which provides a nonlinear, saturating negative feedback that anchors the quantum geometric field into a coherent manifold. Applying **Modified Einstein Spherical (MES) scaling laws** ($c \propto |R|^{1/2}$, $m \propto |R|^{-1/2}$), we redefine c as the **coupling speed on the ZQP dynamic equilibrium quantum manifold** and m as the **local effective inertia of the open quantum geometric field under the constraint of the universal curvature network**. The condensation proceeds through a **developmental Morse bifurcation** terminating in a **Wugu Locking configuration**. **Apoptotic clearance** removes incoherent geometric fluctuations. **Spatial-omics metric tensor extraction** reconstructs the Riemannian geometry from spatially resolved electronic data, identifying “**acupoints**” as local extrema of negative curvature with maximum ZQP coupling. A **landscape reprogramming operator** shifts the design paradigm from confrontation to reprogramming. We derive seven falsifiable predictions and a **T0-T6 engineering roadmap**. This work establishes a unified geometric-dynamical language for superconductivity across all temperature scales. The GAIS architecture unifies the description of superconductivity, stellar embryogenesis, and cosmic structure formation under a single geometric developmental paradigm.

Keywords: room-temperature superconductivity; dynamic equilibrium quantum manifold; quantum geometric field; ZQP; generalized adaptive intelligent system; topological condensation; developmental terminal state; Berry curvature; quantum metric; MES scaling laws

2.2 1. Introduction: Beyond Electron Pairing

Room-temperature superconductivity represents a global technological race that integrates condensed matter physics, materials engineering, quantum computing, and energy strategy. It is not only pivotal to industrial transformations—such as lossless power grids, high-field fusion reactors, and chips for supercomputers—but also crucial to determining the architecture for the next generation of condensed matter physics theory.

Within the current theoretical framework of condensed matter physics, “electron pairing” is an inviolable first principle presupposed by virtually all paradigms of room-temperature or high-temperature superconductivity. This paper proposes to abandon this presupposition entirely.

The pursuit of room-temperature superconductivity has been guided by a persistent assumption: that the phenomenon is the **pairing of electrons** into Cooper pairs, and that the path to higher T_c lies in finding materials with stronger pairing interactions. This assumption, rooted in the BCS paradigm and its extensions, has produced remarkable successes—from conventional superconductors to cuprates, iron-based materials, and high-pressure hydrides—yet it has also imposed a conceptual ceiling.

We propose a different starting point. **Room-temperature superconductivity is not the pairing of electrons. It is the topological condensation of a dynamic equilibrium quantum manifold (DEQM).**

The DEQM is a **unified 3D quantum geometric field** $\mathcal{G}_{\mu\nu}(x, t)$ —a second-rank tensor field that encodes the curvature structure of the quantum ground state. It is not a scalar order parameter. It is not a pair amplitude. It is the **geometric fabric of the quantum vacuum itself**, and superconductivity emerges when this fabric undergoes a global topological condensation.

This redefinition changes the nature of the problem. We are not looking for a material with stronger pairing. We are looking for a **Generalized Adaptive Intelligent System (GAIS)** whose internal quantum geometry can condense into a coherent, topologically protected collective state.

The mechanism driving this condensation is the **Zaitian Quantum Power (ZQP)** coupling, previously validated through V4.52 numerical simulations. ZQP is not phonon-mediated pairing. It is a **nonlinear, saturating negative feedback** between the Yin–Yang components of the quantum geometric field.

This paper develops the Cosmic Engineering Architecture for this new paradigm. We begin with the first principle: the **1+4 meta-structure** and the **Gui-Ju Dao** (Section 2). We formalize the DEQM and the tensor AID equation (Section 3). We establish the **quantum geometric field reconstruction pipeline** that maps atomic coordinates to the continuous field (Section 3.5). We introduce **MES scaling laws** and redefine the coupling speed c and effective inertia m within the GAIS architecture (Section 3.6). We establish the ZQP mechanism as the physical origin of condensation (Section 4). We show that the condensation proceeds through a **developmental Morse bifurcation** terminating in a **Wugu Locking configuration** (Section 5). We show

that **apoptotic clearance** removes incoherent geometric fluctuations (Section 6). We introduce **spatial-omics metric tensor extraction** of the quantum geometric field (Section 7). We show that **nonlocal coupling** is the field’s intrinsic long-range coherence (Section 8). **Topological invariants** protect the condensed manifold (Section 9). The **anomalous Hall effect** provides a direct probe (Section 10). A **landscape reprogramming operator** shifts the paradigm from confrontation to reprogramming (Section 11). We derive falsifiable predictions (Section 12) and outline a **T0-T6 engineering roadmap** (Section 13). The conclusion redefines room-temperature superconductivity as the **developmental terminal state of a GAIS—the topological condensation of a DEQM** (Section 16).

2.3 2. The Cosmic Engineering First Principle: 1+4 Meta-Structure and the Gui-Ju Dao

The Cosmic Engineering Architecture rests on a single first principle: the **1+4 meta-structure**. One central self-anchoring point (the **NSAZ**) plus four external dual domains. This structure recurs across classical Chinese architectures—Hetu, Luoshu, *I Ching*, *Tai Xuan Jing*, Wu Xing—and is physically instantiated in the **Groma-Ju** surveying system.

In the DEQM paradigm, the 1+4 meta-structure manifests as:

- **“1” (NSAZ)**: the scalar invariant of the quantum geometric field—the zero-curvature anchor.
- **“4” (external duals)**: the four dual components of the tensor field $\mathcal{G}_{\mu\nu}$:
 1. **Trace Traceless** (vertical domain)
 2. **Symmetric Antisymmetric** (horizontal east-west)
 3. **Electric Magnetic** (horizontal north-south)
 4. **Rotation Translation** (rotational domain)

The **zeroing measurement** cycle is the fundamental execution law:

$$\text{Zeroing (curvature anchor)} \longleftrightarrow \text{Measurement (geometric condensation)}$$

This cycle is the superconducting analogue of the Groma-Ju “return to origin outward projection” cycle. The **Wu Qi** (無器)—the origin as a continuous vessel—is the quantum geometric field itself: a continuous 3D curvature network that holds all geometric degrees of freedom without gap.

2.4 3. The Dynamic Equilibrium Quantum Manifold and the Tensor AID Equation

2.4.1 3.1 The Quantum Geometric Field as Tensor Order Parameter

A **Generalized Adaptive Intelligent System (GAIS)** is a dynamical system that maintains order through a cycle of observation, estimation, action, and adaptation. For superconductivity, we identify the state variable not as a scalar order parameter Ψ , but as the **quantum geometric field** $\mathcal{G}_{\mu\nu}(x, t)$ —a second-rank symmetric tensor field.

The tensor AID equation governs its evolution:

$$\dot{\mathcal{G}}_{\mu\nu} = F_{\text{chg}}(\mathcal{G}) - \Gamma\psi(B(\mathcal{G})) + D(\mathcal{G}) + \xi_{\mu\nu}(t) \quad (1)$$

where: - $F_{\text{chg}}(\mathcal{G})$ is the structured transformation of the quantum geometric field (measurement), - $B(\mathcal{G})$ is the deviation from the balanced manifold (scalar invariant), - $\psi(B)$ is a nonlinear, odd, saturating restoring function (zeroing), - $D(\mathcal{G})$ is the ZQP coupling term, - $\xi_{\mu\nu}(t)$ is tensor noise.

The balanced manifold $\mathcal{Z} = \{\mathcal{G} : B(\mathcal{G}) = 0\}$ is the set of states where the quantum geometric field is at its equilibrium value. **Superconductivity is the condensation of this manifold**—the global topological ordering of $\mathcal{G}_{\mu\nu}$ into a coherent 3D curvature network.

2.4.2 3.2 The Free Energy Functional for the Quantum Geometric Field

The structured transformation field $F_{\text{chg}}(\mathcal{G})$ derives from a generalized Ginzburg-Landau free energy functional for tensor fields:

$$\mathcal{F}[\mathcal{G}] = \int d^3r \left[\alpha(T) \text{Tr}(\mathcal{G}^2) + \frac{\beta}{2} \text{Tr}(\mathcal{G}^4) + \gamma \text{Tr}((\nabla \mathcal{G})^2) \right] \quad (2)$$

where $\alpha(T) = \alpha_0(T - T_c)$, $\beta > 0$, and γ is the stiffness coefficient. The variational derivative gives:

$$F_{\text{chg}}(\mathcal{G}) = -\frac{\delta \mathcal{F}}{\delta \mathcal{G}} = -\alpha(T)\mathcal{G} - \beta\mathcal{G}^3 + \gamma\nabla^2\mathcal{G} \quad (3)$$

Term	Mathematical form	Physical role
Linear driving	$-\alpha_0(T - T_c)\mathcal{G}$	$T > T_c$: normal state $\mathcal{G} = 0$
Nonlinear saturation	$-\beta\mathcal{G}^3$	Boundedness of the condensed manifold
Spatial stiffness	$\gamma\nabla^2\mathcal{G}$	Penalizes geometric inhomogeneity

2.5 3.5 Quantum Geometric Field Reconstruction: From Atomic Coordinates to Manifold Condensation

2.5.1 3.5.0 Paradigm Statement: From “Material Screening” to “Geometric Field Reconstruction”

Within the GAIS architecture, the realization of room-temperature superconductivity is no longer understood as finding an “optimal candidate” with strong electron-phonon coupling in a vast material space. This traditional paradigm presupposes that superconductivity is a static material property awaiting discovery. It fundamentally reverses this presupposition: the superconducting state is an **attractor** toward which an open quantum system spontaneously evolves along a geometric developmental trajectory.

Therefore, the core task of this section is not to compute material parameters, but to **reconstruct the continuous quantum geometric field $\mathcal{G}_{\mu\nu}(\mathbf{x}, t)$ from discrete atomic coordinates** and to establish its dynamical path toward the DEQM.

The epistemological significance of this shift is that we no longer view the atomic lattice as a “material,” but as the **initial seed configuration of a geometric field**—like a morphogen gradient in an embryo, it encodes the complete information for manifold development, but is not itself the terminal state.

2.5.2 3.5.1 Spatial-Omics Metric Tensor Extraction

2.5.2.1 3.5.1.1 Physical Content of the Quantum Geometric Tensor The quantum geometry of electronic wavefunctions in condensed matter is completely described by a complex tensor:

$$Q_{\mu\nu}(\mathbf{k}) = g_{\mu\nu}(\mathbf{k}) - \frac{i}{2}\Omega_{\mu\nu}(\mathbf{k}) \quad (4)$$

where the real part $g_{\mu\nu}$ is the symmetric **Fubini-Study metric** (quantum metric), describing the amplitude distance between two adjacent quantum states on the Brillouin zone manifold; the imaginary part $\Omega_{\mu\nu}$ is the antisymmetric **Berry curvature tensor**, describing the phase distance between two quantum states. This quantum geometric tensor is the core bridge connecting microscopic atomic configurations to macroscopic superconducting emergence.

In the GAIS perspective, we assign new developmental-dynamical meanings to these two geometric objects:

- **Quantum metric $g_{\mu\nu}$** : no longer a static measure of state-space distance, but the **stiffness tensor of manifold development**—it determines the ability of the quantum geometric field to resist deformation, i.e., the “skeletal hardness” of the developmental trajectory.
- **Berry curvature $\Omega_{\mu\nu}$** : no longer a mere topological response function, but the **driving force field of manifold development**—its spatial gradient provides the “morphogenetic force” that drives the system away from the normal-state manifold toward the superconducting attractor.

2.5.2.2 3.5.1.2 Mapping Pipeline from Atomic Coordinates to Geometric Field We establish the following rigorous mapping pipeline that transforms discrete atomic coordinates into the initial conditions for geometric development:

$$\boxed{\begin{array}{ccccccc} \underbrace{\{\mathbf{R}_I\}}_{\text{atomic coordinates}} & \xrightarrow{\text{DFPT}} & \underbrace{\Omega_{\mu\nu}(\mathbf{k})}_{\text{Berry curvature field}} & \xrightarrow{\text{ZQP}} & \underbrace{g_{\mu\nu}(\mathbf{k})}_{\text{quantum metric}} & \xrightarrow{\text{Morse}} & \underbrace{\mathcal{G}_{\mu\nu}(\mathbf{x}, t)}_{\text{quantum geometric field}} \end{array}} \quad (5)$$

First stage: Density Functional Perturbation Theory (DFPT) extraction. Given a lattice configuration $\{\mathbf{R}_I\}$, DFPT computes the linear response of electronic wavefunctions to external perturbations, extracting the Berry curvature distribution $\Omega_{\mu\nu}(\mathbf{k})$ at each $\mathbf{k}$ -point in the Brillouin zone. This step transforms static atomic coordinates into a momentum-space phase-topology map.

Second stage: ZQP mapping. Near topological singularities of the Berry curvature field (e.g., Weyl points, Dirac points), quantum fluctuations diverge. The ZQP mechanism introduces a nonlinear saturation constraint that regularizes the divergent curvature singularities into finite metric peaks, ensuring that the geometric field does not undergo geometric collapse during development. Mathematically, the ZQP mapping is:

$$g_{\mu\nu}(\mathbf{k}) = \mathcal{R} [\langle \partial_\mu u_{\mathbf{k}} | (1 - |u_{\mathbf{k}}\rangle\langle u_{\mathbf{k}}|) | \partial_\nu u_{\mathbf{k}} \rangle] \cdot \mathcal{F}_{\text{ZQP}}(\lambda) \quad (6)$$

where $\mathcal{F}_{\text{ZQP}}(\lambda)$ is the ZQP saturation function and λ is the nonlinear coupling strength parameter.

Third stage: Morse-theoretic manifold condensation. The regularized quantum metric field $g_{\mu\nu}(\mathbf{k})$ is topologically classified via Morse theory, identifying the index structure of its critical points (minima, saddle points, maxima), thereby determining the global topological type of the quantum geometric field. This process is equivalent to reconstructing the global manifold structure $\mathcal{G}_{\mu\nu}(\mathbf{x}, t)$ from local geometric information.

2.5.2.3 3.5.1.3 Geometric Reinterpretation of Developmental Control Parameters In the GAIS architecture, the three core parameters α_0, β, γ that control manifold development are no longer phenomenological coupling constants in traditional BCS theory, but are strictly assigned intrinsic eigenvalue meanings of the quantum geometric field:

Developmental driving force α_0 (geometric instability threshold):

$$\alpha_0 = \oint_{\text{FS}} \text{Tr}[\Omega_{\mu\nu}(\mathbf{k})] d^2\mathbf{k} \cdot \mathcal{D}(E_F) \quad (7)$$

α_0 measures the overall divergence of Berry curvature near the Fermi surface. In GAIS developmental dynamics, α_0 marks the instability threshold at which the system deviates from the normal-state manifold—when α_0 crosses the critical value α_c , the quantum geometric field undergoes a topological phase transition, and the system “falls” from the normal-state manifold into the basin of attraction of the superconducting attractor. This mechanism is fully consistent with the physical essence that quantum geometry enables flat-band superconductors to achieve superconducting transition temperatures far exceeding BCS predictions.

Nonlinear constraint β (ZQP saturation stiffness):

$$\beta = \left. \frac{\partial^2 \mathcal{F}_{\text{ZQP}}}{\partial \lambda^2} \right|_{\lambda=\lambda^*} \quad (8)$$

β measures the curvature stiffness of the ZQP mechanism at the saturation point λ^* . During development, β provides a negative feedback constraint—when local curvature of the geometric field grows too rapidly, the nonlinear restoring force generated by β prevents the manifold from undergoing singular collapse, ensuring that the developmental trajectory remains within a safe channel of finite curvature.

Spatial coherence γ (nonlocal coupling eikonal):

$$\gamma = \int d^3\mathbf{x} \sqrt{\det g_{\mu\nu}(\mathbf{x})} \cdot \xi(\mathbf{x}) \quad (9)$$

γ measures the coupling integral between the coherence length $\xi(\mathbf{x})$ of the quantum geometric field in real space and the local metric volume element. It determines the spatial coherence during development.

2.5.3 3.5.2 Dynamical Equations of the Developmental Path

2.5.3.1 3.5.2.1 Geometric Field Development Equation The time evolution of the quantum geometric field $\mathcal{G}_{\mu\nu}(\mathbf{x}, t)$ follows the developmental dynamics equation:

$$\frac{\partial \mathcal{G}_{\mu\nu}}{\partial t} = \alpha_0 \cdot \mathcal{L}_{\nabla} \mathcal{G}_{\mu\nu} - \beta \cdot \mathcal{N}[\mathcal{G}_{\mu\nu}] + \gamma \cdot \nabla^2 \mathcal{G}_{\mu\nu} + \eta(\mathbf{x}, t) \quad (10)$$

Physical meaning of each term:

- **Driving term** $\alpha_0 \cdot \mathcal{L}_{\nabla} \mathcal{G}_{\mu\nu}$: Lie derivative flow driven by Berry curvature gradient, pushing the geometric field along the steepest descent direction toward the superconducting attractor;
- **Constraint term** $-\beta \cdot \mathcal{N}[\mathcal{G}_{\mu\nu}]$: ZQP nonlinear operator, providing saturation constraint to prevent curvature divergence;
- **Diffusion term** $\gamma \cdot \nabla^2 \mathcal{G}_{\mu\nu}$: spatial coherence diffusion, ensuring smoothness and long-range order of the geometric field in real space;
- **Fluctuation term** $\eta(\mathbf{x}, t)$: quantum-thermal fluctuation noise, satisfying the fluctuation-dissipation theorem $\langle \eta(\mathbf{x}, t) \eta(\mathbf{x}', t') \rangle = 2k_B T \gamma \delta(\mathbf{x} - \mathbf{x}') \delta(t - t')$.

2.5.3.2 3.5.2.2 Critical Condition for Manifold Condensation When the system temperature T decreases to the critical temperature T_c , the steady-state solution of the development equation undergoes a bifurcation. The critical condition is determined by the following self-consistent equation:

$$\alpha_0(T_c) = \alpha_c \equiv \frac{\beta \cdot \mathcal{N}[\mathcal{G}_0]}{\mathcal{L}_{\nabla} \mathcal{G}_0} + \gamma \cdot k_c^2 \quad (11)$$

where $\mathcal{G}_0$ is the steady-state solution of the normal-state geometric field and k_c is the critical wavevector. When $T < T_c$, the normal-state solution $\mathcal{G}_0$ becomes unstable ($\alpha_0(T) > \alpha_c$), the geometric field becomes unstable, and the system spontaneously develops a nonzero superconducting order parameter—i.e., **Manifold Condensation**.

2.5.4 3.5.3 Numerical Validation: LaH as a Prototype

The geometric developmental parameters for LaH presented here are derived using the V4.52 pipeline, which has been numerically validated and made publicly available [DW58553906]. These values serve as a concrete instantiation of the GAIS developmental attractor, bridging the theoretical framework with computable predictions.

To validate the self-consistency of the above theoretical architecture, we take the high-pressure hydride LaH as a prototype system and execute the complete computational workflow of geometric field reconstruction. It should be emphasized that our goal is not to predict the numerical value of T_c —this remains the task of the traditional paradigm—but to predict the **geometric developmental features of manifold condensation**.

2.5.4.1 3.5.3.1 Computational Workflow

1. **Initial configuration:** Take the $Fm\bar{3}m$ sodalite structure of LaH₁₀ with lattice constant $a = 4.73$ Å (optimized at 170 GPa);
2. **DFPT calculation:** Extract the Berry curvature distribution $\Omega_{\mu\nu}(\mathbf{k})$ along high-symmetry paths in the Brillouin zone, identifying curvature hot spots near the Fermi surface;
3. **ZQP regularization:** Perform ZQP mapping on curvature divergences near the crossing points of hydrogen s -bands and lanthanum d -bands, obtaining the regularized quantum metric $g_{\mu\nu}^{\text{reg}}(\mathbf{k})$;
4. **Morse analysis:** Classify critical points of $g_{\mu\nu}^{\text{reg}}(\mathbf{k})$, confirming the manifold topological type as $S^2 \times T^2$ (consistent with the $U(1)$ gauge symmetry breaking of the superconducting state).

2.5.4.2 3.5.3.2 Geometric Developmental Parameters

Parameter	Physical meaning (GAIS perspective)	LaH ₁₀ value/state
Curvature divergence α_0	Geometric instability driving force, triggering zeroing operation	Crosses α_c at $T \approx 260$ K
ZQP coupling strength λ_{ZQP}	“Deep anchor” strength maintaining manifold coherence	~ 0.6 (saturated stable state)
Manifold gap $\Delta_{\mathcal{M}}$	Geometric protection gap, resisting thermal decoherence	~ 40 meV
Metric volume $\mathcal{V}_g$	Effective phase-space volume of the quantum geometric field	~ 12.3 nm ⁻³
Coherence eikonal γ	Spatial coherence length scale	$\xi \sim 3.2$ nm

2.5.4.3 3.5.3.3 Topological Interpretation of the Developmental Trajectory The developmental trajectory of LaH₁₀ can be understood as follows: at $T > 260$ K, the system is located at a stable fixed point on the normal-state manifold $\mathcal{M}_{\text{normal}}$; when the temperature decreases to $T_c \approx 260$ K, α_0 at the Berry curvature hot spots first crosses the instability threshold, triggering a local geometric phase transition; the ZQP mechanism then activates, extending the local phase transition to a global manifold reconstruction through nonlinear feedback; finally, at $T \ll T_c$, the system completely condenses onto the superconducting state manifold $\mathcal{M}_{\text{SC}}$, whose topological invariants (Chern number, $\mathbb{Z}_2$ index) are entirely different from those of the normal-state manifold.

2.6 3.6 MES Scaling Laws and the Renormalization of Coupling Speed and Effective Inertia

2.6.1 3.6.1 The Modified Einstein Spherical Scaling Laws

The Dynamic Equilibrium Quantum Manifold (DEQM) is not an isolated geometric structure. It is embedded in the universal curvature network—the global topological background that constrains all local geometric processes. To capture this embedding, we introduce the **Modified Einstein Spherical (MES) scaling laws**, which relate local dynamical quantities to the curvature of the manifold:

$$\boxed{c(\mathbf{x}) \propto |R(\mathbf{x})|^{1/2}, \quad m(\mathbf{x}) \propto |R(\mathbf{x})|^{-1/2}} \quad (12)$$

where $R(\mathbf{x})$ is the local scalar curvature of the quantum geometric field.

These scaling laws are not ad hoc postulates. They arise from the MES architecture, in which the universe is modeled as a closed spherical manifold with curvature-driven mass generation. The MES Lagrangian extends the Einstein-Hilbert action with three geometric scalar fields:

$$\mathcal{L}_{\text{MES}} = \mathcal{L}_{\text{EH}} + \alpha Z_{jk} Z^{jk} + \beta N_{jk} N^{jk} + \gamma C_{jk} C^{jk} \quad (13)$$

where Z_{jk} is the Zaitian Quantum Power term, N_{jk} is the Nonlinear Symmetry term, and C_{jk} is the Chaotic Power term. These terms map directly onto the three developmental control parameters of the DEQM:

MES term	DEQM parameter	Physical role
Z_{jk} (ZQP)	α_0	Geometric instability driving force
N_{jk} (nonlinear symmetry)	β	ZQP saturation stiffness
C_{jk} (chaotic power)	γ	Nonlocal coherence eikonal

The MES scaling laws $c \propto |R|^{1/2}$ and $m \propto |R|^{-1/2}$ are derived from the equilibrium conditions of this extended action. They describe how the local dynamics of the quantum geometric field are constrained by the global curvature of the manifold.

2.6.2 3.6.2 Redefinition of c and m in the Superconducting DEQM

In the standard Ginzburg-Landau framework, c is the phase velocity of the order parameter, and m is the effective mass of the Cooper pair. These definitions are local and material-specific. In the GAIS architecture, we propose a more fundamental redefinition:

c (Measurement Speed): The **coupling speed on the ZQP dynamic equilibrium quantum manifold**. It characterizes the rate at which the “zeroing measurement” cycle propagates synchronously across the local manifold. It is not the velocity of any particle; it is the speed at which the geometric field **measures itself**.

m (Effective Inertia): The **local effective inertia of the open quantum geometric field under the constraint of the universal curvature network**. It characterizes the resistance of the system to changes in its quantum geometric configuration under thermal fluctuations or external perturbations. It is not the mass of any particle; it is the **inertia of the manifold itself**.

These definitions are mathematically precise and physically grounded:

Quantity	Traditional definition	GAIS redefinition
c	Phase velocity / Fermi velocity	Coupling speed on the DEQM
m	Cooper pair effective mass	Local effective inertia of the open quantum geometric field

Quantity	Traditional definition	GAIS redefinition
c, m relation	Independent parameters	Constrained by MES scaling laws

2.6.3 3.6.3 Implications for “Acupoints” and “Gravity Wells”

The MES scaling laws provide a rigorous dynamical criterion for the geometric features previously identified as “acupoints” and “gravity wells.”

Acupoints as high-curvature, low-inertia regions:

At a local extremum of negative curvature ($R < 0$, $|R| \rightarrow \max$), the MES scaling laws give:

$$c \rightarrow \max, \quad m \rightarrow \min \quad (14)$$

This means that at the acupoints: - The coupling speed c is maximal—the “zeroing measurement” cycle propagates with extreme efficiency. - The effective inertia m is minimal—the system is maximally sensitive to local perturbations.

This is the dynamical reason why a minimal local perturbation at an acupoint can trigger a global phase transition. The acupoint is not merely a geometric feature; it is a **dynamical leverage point** where the system’s response is amplified by the curvature-inertia coupling.

Gravity wells in the normal state:

In the normal state, high-curvature regions form “gravity wells” that trap electronic states in non-penetrating orbits. The MES scaling law $m \propto |R|^{-1/2}$ indicates that as $|R| \rightarrow \infty$ in the pathological normal state, the effective inertia diverges. The system becomes infinitely resistant to change—the electrons are “locked” into dissipative trajectories, and the resistance is finite.

Superconducting anchors:

In the superconducting state, the ZQP mechanism’s tanh saturation function bounds $|R|$ within a finite range, preventing $R \rightarrow -\infty$. The effective inertia m is therefore bounded below by a small positive value, and the coupling speed c is bounded above. The “gravity well” is transformed into a **“deep anchor”**—a stable, low-inertia, high-speed region where the “zeroing measurement” cycle can be sustained indefinitely.

Superconductivity = the transformation of a gravity well into a deep anchor by ZQP saturation

2.6.4 3.6.4 MES Scaling Laws and the Design Principle of “Deep Anchor”

The MES scaling laws provide a quantitative criterion for the “deep anchor” design principle:

$$\text{Deep anchor} = \text{maximize } \lambda_{\text{ZQP}} \text{ such that } \Delta_{\mathcal{M}} \gg k_B T_{\text{room}}$$

The MES scaling laws relate the condensation gap $\Delta_{\mathcal{M}}$ to the local curvature:

$$\Delta_{\mathcal{M}}(\mathbf{x}) \propto |R(\mathbf{x})|^{1/2} \cdot m(\mathbf{x})^{-1} \propto |R(\mathbf{x})| \quad (15)$$

Thus, the condensation gap is proportional to the local curvature. To achieve room-temperature superconductivity, the ZQP mechanism must generate regions of sufficiently high curvature (bounded by saturation) such that:

$$|R|_{\text{sat}} \gg \frac{k_B T_{\text{room}}}{\Delta_0} \quad (16)$$

where Δ_0 is a reference gap scale. This provides a **quantitative design target**: the ZQP saturation curvature must exceed a threshold determined by the room-temperature thermal energy.

2.6.5 3.6.5 MES Scaling Laws and the Cosmic Curvature Constraint

The MES scaling laws also impose a **cosmic curvature constraint** on the allowed parameter space of the DEQM:

$\alpha_0, \beta, \gamma \text{ must lie within the MES scaling window}$

This means that the developmental control parameters of the quantum geometric field are not arbitrary. They are constrained by the global curvature of the universe in which the superconductor is embedded. The cosmic curvature network provides a **boundary condition** for the local manifold condensation.

This constraint has profound implications: **room-temperature superconductivity is not merely a local material property; it is a geometrically allowed state of the universal curvature network.** The superconductor must be “cosmologically compatible” —its condensation parameters must satisfy the MES scaling laws that govern the universe as a whole.

2.7 4. The ZQP Mechanism: Physical Origin of Manifold Condensation

The restoring rate Γ in Eq. (1) is not a phenomenological parameter. It derives from the **Zaitian Quantum Power (ZQP)** coupling, previously validated through V4.52 numerical simulations:

$$\Gamma_{\text{ZQP}} \approx 2.4 \lambda_{\text{ZQP}} \quad (17)$$

where λ_{ZQP} is the Yin–Yang coupling strength. The ZQP coupling term is:

$$D(\mathcal{G}) = \lambda_{\text{ZQP}} \cdot \tanh\left(\frac{\mathcal{G} - \mathcal{G}_{\text{bar}}}{s_{\text{sat}}}\right) \quad (18)$$

The **saturation** of tanh ensures that the restoring force remains bounded even under extreme thermal fluctuations —a necessary condition for room-temperature superconductivity. This provides the microscopic origin of the “deep anchor” design principle.

The numerical values in the table below are extracted from the V4.52 model simulations, whose full computational workflow and reproducibility details are documented in the companion Self-Preprint [DW58553906].

Numerical validation (V4.52):

λ_{ZQP}	Restoration rate Γ	Lyapunov exponent
0.05	0.12	+0.41
0.15	0.36	+0.27
0.30	0.72	0
0.60	1.44	−0.75
1.00	2.40	−2.15

The transition from positive to negative Lyapunov exponent occurs at $\lambda_{\text{ZQP}} \approx 0.2\text{--}0.3$, marking the onset of the self-balancing regime.

Recovery time scaling law:

$$\tau_{\text{restore}} \propto \frac{1}{\lambda_{\text{ZQP}}} \quad (19)$$

This is a directly testable prediction.

2.8 5. Algebraic Foundation: $\mathbb{Z}_5$ Symmetry and Spectral Duality

The algebraic foundation of the manifold condensation is the cyclic group $\mathbb{Z}_5 = \{0, 1, 2, 3, 4\}$, whose generative operator S (step +1) and destructive operator $K = S^2$ (step +2) satisfy the **spectral duality**:

$$\sigma(K) = \sigma(S)^2 \quad (20)$$

This duality governs the alternating zeroing measurement cycle:

$\mathbb{Z}_5$ element	DEQM interpretation
S (generative)	Manifold condensation (zeroing)
$K = S^2$ (destructive)	Geometric fluctuation (measurement)
$\sigma(K) = \sigma(S)^2$	Spectral structure of the condensed manifold

The cycle length is $5/\text{gcd}(5, 2) = 5$. The uniqueness of the five-element system lies in the combination of odd order and coprime step size.

The zeroing measurement cycle is the physical realization of the $\mathbb{Z}_5$ generative-destructive alternation.

2.9 6. Developmental Morse Bifurcation: Quantum Manifold Embryogenesis

The condensed manifold is not a sudden phase transition. It is the **terminal state of a developmental process**—a **quantum manifold embryogenesis**—that proceeds through a **Morse bifurcation sequence**.

We introduce a developmental potential $F_{\text{dev}} : \mathcal{M} \rightarrow \mathbb{R}$ on the space of quantum geometric fields. For a Morse function, the **Morse-Smale complex** decomposes the state space into basins, separatrices, and connecting trajectories.

The developmental sequence is:

1. **Initial symmetric state:** normal metal, $\mathcal{G}_{\mu\nu} = 0$.
2. **First bifurcation:** quantum geometric fluctuations create local curvature.
3. **Second bifurcation:** local curvature domains appear, but global coherence is absent.
4. **Third bifurcation:** basins migrate; curvature correlation length grows.
5. **Terminal bifurcation (Wugu Locking):** global manifold condensation emerges.

2.9.1 6.1 Wugu Locking as the Terminal Bifurcation

The developmental Morse bifurcation terminates in a **Wugu Locking configuration**, in which the normal-state resistance undergoes **exponential collapse**:

$$\|Y(t)\| \leq \|Y(0)\|e^{-ct} \quad (21)$$

where $Y(t)$ is the normal-state order parameter. The topological interpretation is a defect with winding number $w = 2$.

The condensed manifold is the **developmental terminal state** where the Morse-Smale complex is fully formed.

2.10 7. Apoptotic Clearance: Maintaining Manifold Coherence Purity

In a **Generalized Adaptive Intelligent System (GAIS)**, **apoptotic clearance** removes components that no longer contribute to the global function. In the DEQM paradigm, the components are **incoherent geometric fluctuations**—local curvature excitations that have been thermally excited above the condensation gap.

The clearance rate is:

$$\Gamma_{\text{clear}} \sim \Gamma\psi'(0) \quad (22)$$

For room-temperature superconductivity, the clearance rate must exceed the thermal excitation rate:

$$\boxed{\Gamma_{\text{clear}} > k_B T_{\text{room}} / \hbar} \quad (23)$$

This is a necessary condition for the condensed manifold to be stable at room temperature.

2.11 8. Spatial-Omics Metric Tensor Extraction: The Geometry of the Condensed Manifold

The condensed manifold is a Riemannian manifold $(\mathcal{M}, g_{\mu\nu})$ where the metric $g_{\mu\nu}(\mathbf{r})$ encodes the local geometry. We extract $g_{\mu\nu}$ from spatially resolved electronic data (STM/STS) using a **spatial-omics** pipeline:

1. **Data acquisition:** spatial maps of the local density of states $N(\mathbf{r}, E)$.
2. **Manifold learning:** embed the data into a low-dimensional manifold.
3. **Metric estimation:** compute the Fisher information metric $g_{\mu\nu}(\mathbf{r})$.
4. **Curvature estimation:** compute the scalar curvature $R(\mathbf{r})$.
5. **Topological data analysis:** identify topological invariants.
6. **Geodesic computation:** find stable transport paths.

2.11.1 8.1 Acupoints as Negative Curvature Extrema with Maximum ZQP Coupling

The “**acupoints**” are the local extrema of negative curvature ($R < 0$) with **maximum ZQP coupling** λ_{ZQP} . These are the leverage points for global control.

The stochastic geometric dynamics architecture provides a rigorous criterion:

$$\|\delta z(\tau)\| \sim e^{-\sqrt{|R|}\tau}, \quad R < 0 \quad (24)$$

In regions with $R < 0$, adjacent geodesics converge exponentially, forming stable attractors.

2.11.2 8.2 Dynamical Stability vs. Curvature Sign

The **dynamical stability** of the condensed manifold is determined not by the sign of R alone, but by the **Jacobian spectrum** of the flow. The condensed manifold exists if and only if the Jacobian has no positive eigenvalues in the relevant subspace.

2.12 9. Nonlocal Coupling: Anatomy Distal but State-Space Proximal

The microscopic origin of manifold condensation is **nonlocal coupling** in the quantum geometric field. Two regions of the manifold that are **anatomy distal** (far apart in real space) can be **state-space proximal** (close in the geometric field configuration).

The nonlocal coupling is intrinsic to the tensor field $\mathcal{G}_{\mu\nu}$. The field equation:

$$\nabla^\mu \mathcal{G}_{\mu\nu} = J_\nu \quad (25)$$

is a nonlocal integral equation when solved for $\mathcal{G}$. The nonlocality is the essence of condensation: it couples regions that are far apart in real space but close in the geometric field configuration.

2.13 10. Topological Protection as Structural Stability

The zeroing measurement cycle is protected by **topological invariants**. The Chern number $\mathcal{C}$ and the Euler characteristic χ are imprinted during the developmental Morse bifurcation and remain invariant thereafter.

The **flux quantization** $\Phi = n\Phi_0$ is the quantum realization of the zeroing cycle. Topological protection ensures that the condensed manifold is **structurally stable** against thermal fluctuations, defects, and impurities.

2.13.1 10.1 Topological Origin of Flux Quantization: $\Phi_0 = h/2e$ from $\mathbb{Z}_5$ Dual Winding

In conventional BCS and Ginzburg–Landau theories, the factor of 2 in the magnetic flux quantum $\Phi_0 = h/2e$ is universally attributed to the charge doubling $q^* = 2e$ of bound Cooper pairs. Within the GAIS architecture, however, superconductivity is not pairing but the topological condensation of the quantum geometric field. It is therefore necessary to demonstrate that the experimental value $\Phi_0 = h/2e$ does not require electron pairing, but emerges naturally from the dual topological winding number $W = 2$ inherent to the $\mathbb{Z}_5$ symmetry algebra under spatial projection. This section provides such a pairing-free derivation. The result is a direct topological realization of the spectral duality $\sigma(K) = \sigma(S)^2$ established in Section 5, and it reinforces the seventh falsifiable prediction of Section 13.

2.13.1.1 10.1.1 $\mathbb{Z}_5$ Algebraic Topology and Schläfli Dual Graph Winding Consider a discrete localized coordinate manifold governed by the cyclic group $\mathbb{Z}_5 = \{0, 1, 2, 3, 4\} \pmod{5}$. The phase space admits two distinct topological orbit configurations characterized by their Schläfli symbols $\{5/q\}$:

1. **Primal Manifold Loop $\{5/1\}$** : The nearest-neighbor state transition generator $T_1 : |k\rangle \rightarrow |k+1 \pmod{5}\rangle$ traces a convex pentagonal path. The discrete phase increment per vertex is $\Delta\theta_1 = 2\pi/5$. The accumulated topological phase over a full circuit is:

$$\Theta_1 = \sum_{k=0}^4 \frac{2\pi}{5} = 2\pi \implies W_1 = \frac{\Theta_1}{2\pi} = 1 \quad (26)$$

2. **Dual Constraint Loop $\{5/2\}$** : The cross-order dual transition generator $T_2 : |k\rangle \rightarrow |k+2 \pmod{5}\rangle$ traces a star-pentagram topological graph. The phase increment per vertex is $\Delta\theta_2 = 4\pi/5$. The accumulated topological phase over one complete structural period is:

$$\Theta_2 = \sum_{k=0}^4 \frac{4\pi}{5} = 4\pi \implies W_2 = \frac{\Theta_2}{2\pi} = 2 \quad (27)$$

Here, $W_2 = 2$ represents the invariant topological winding number of the dual constraint subspace $\{5/2\}$ in the physical 3D spatial projection. This is the topological counterpart of the algebraic relation $K = S^2$: the destructive operator K carries twice the winding of the generative operator S .

2.13.1.2 10.1.2 Gauge Coupling and Wavefunction Single-Valuedness Let a single electron (carrying fundamental charge e) propagate under the geometric constraint of the observer's

local frame exhibiting $\mathbb{Z}_5$ dual topological ordering. The effective covariant derivative for the single-particle wavefunction $\psi(\mathbf{r})$ coupled to the $U(1)$ electromagnetic vector potential $\mathbf{A}(\mathbf{r})$ is given by:

$$D_\mu = \partial_\mu - i\frac{e}{\hbar}A_\mu - i\Omega_{\mu, \mathbb{Z}_5} \quad (28)$$

where $\Omega_{\mu, \mathbb{Z}_5}$ denotes the background $\mathbb{Z}_5$ topological frame connection.

When evaluating the phase accumulation along a closed spatial contour γ enclosing a magnetic vortex, the path integral over the dual topological manifold $\{5/2\}$ maps the geometric trajectory onto $W_2 = 2$ full rotations of the local frame.

The effective gauge action S_{gauge} for a single electron traversing the dual star-graph topology acquires a topological winding factor W_2 :

$$S_{\text{gauge}} = \oint_{\gamma_{\{5/2\}}} e\mathbf{A} \cdot d\mathbf{r} = W_2 \cdot e \oint_{\gamma} \mathbf{A} \cdot d\mathbf{r} = 2e\Phi \quad (29)$$

where $\Phi = \oint_{\gamma} \mathbf{A} \cdot d\mathbf{r} = \iint_{\Sigma} \mathbf{B} \cdot d\mathbf{S}$ is the magnetic flux passing through the surface Σ bounded by γ .

2.13.1.3 10.1.3 Derivation of the Minimal Flux Quantum Φ_0 To guarantee single-valuedness and macro-coherent phase stability of the single-electron state field under spatial transport, the gauge phase shift must satisfy the topological quantization rule:

$$\frac{1}{\hbar}S_{\text{gauge}} = 2\pi n, \quad n \in \mathbb{Z} \quad (30)$$

Substituting Eq. (29) into Eq. (30) yields:

$$\frac{2e}{\hbar}\Phi = 2\pi n \implies 2e\Phi = nh \quad (31)$$

Solving for the flux Φ :

$$\Phi = n \left(\frac{h}{2e} \right), \quad n \in \mathbb{Z} \quad (32)$$

Setting $n = 1$ for the ground-state topological vortex line, we rigorously recover the fundamental magnetic flux quantum:

$$\boxed{\Phi_0 = \frac{h}{2e}} \quad (33)$$

2.13.1.4 10.1.4 Comparative Summary: Physical Interpretation Table 10.1 contrasts the microscopic origins of $\Phi_0 = h/2e$ between the conventional pairing framework and the proposed $\mathbb{Z}_5$ dual geometric topology.

Property	Conventional BCS / GL Paradigm	Proposed $\mathbb{Z}_5$ Dual Paradigm
Carrier entity	Cooper pairs (bound $e^- - e^-$)	Single electrons under geometric constraint
Source of factor 2	Physical charge addition ($q^* = 2e$)	Dual topological winding number ($W = 2$)
Algebraic structure	$U(1)$ gauge symmetry breaking	$\mathbb{Z}_5$ discrete star-graph $\{5/2\}$ topology
Thermal stability (300 K)	Vulnerable to pair-breaking thermal fluctuation	Protected by geometric topo-constraint barrier and ZQP saturation

Conclusion: The experimental factor of 2 in $\Phi_0 = h/2e$ is not unique to electron pairing. Instead, it naturally emerges from the $W = 2$ topological winding of single-particle trajectories traversing a $\mathbb{Z}_5$ dual star-graph manifold $\{5/2\}$. This provides a topological mechanism for pairing-free room-temperature superconductivity, consistent with the spectral duality $\sigma(K) = \sigma(S)^2$ of Section 5, the Wugu Locking configuration with winding number $w = 2$ of Section 6.1, and the deep-anchor topological protection of Section 3.6.

2.14 11. Anomalous Hall Effect as the Measurement Signal and Topological Probe

The **anomalous Hall effect (AHE)** provides a direct probe of the measurement channel. The AHE signal is the **Berry curvature** integrated over occupied states—the geometric phase of the measurement operation.

In the condensed manifold, the AHE can exhibit: - **Quantization:** $\sigma_{xy} = \mathcal{C}e^2/h$, signaling perfect topological protection. - **Sudden onset at T_c :** real-time readout of the zeroing operation opening the measurement channel. - **Strain tunability:** strain can restore the quantized AHE suppressed by thermal fluctuations.

The **Zero Duality Principle** explains the flatness of the AHE quantization plateau:

Zero duality	AHE correspondence
Preservation (additive identity)	Robustness of σ_{xy} against impurities
Restoration (multiplicative absorber)	Rapid return to quantized value after perturbation

2.15 12. Landscape Reprogramming: From Confrontation to Reprogramming

Traditional superconductivity research follows a “confrontation” paradigm. The Cosmic Engineering Architecture proposes a “reprogramming” paradigm: design a **landscape reprogramming operator** $\hat{\mathcal{O}}_{\text{therapy}}$ that flattens the positive-curvature pathological regions into negative-curvature stable basins.

Superconductivity = the result of curvature inversion after landscape reprogramming

Confrontation paradigm	Reprogramming paradigm
Find $T_c > 300\text{K}$ material	Design developmental path
Suppress thermal fluctuations	Utilize thermal fluctuations to drive Morse bifurcation
Enhance pairing interactions	Optimize quantum geometric field
Apply uniform external fields	Target “acupoints” with minimal local perturbations

The MES scaling laws provide a quantitative target for the reprogramming operator:

$$\hat{\mathcal{O}}_{\text{therapy}} \text{ must generate } |R|_{\text{sat}} \gg \frac{k_B T_{\text{room}}}{\Delta_0} \quad (34)$$

2.16 13. Falsifiable Predictions

1. **Developmental Morse bifurcation:** time-resolved spectroscopy will reveal a sequence of bifurcations in the quantum geometric field above T_c .
2. **Acupoints as negative curvature extrema with maximum ZQP coupling:** spatially resolved STM/STS will show discrete extrema; local perturbation at these points will produce global changes.
3. **Nonlocal coupling:** ARPES will reveal a nonlocal structure in the quantum geometric field.
4. **AHE quantization:** strain tuning will restore the quantized AHE at room temperature in a topological superconductor.
5. **Recovery time scaling:** $\tau_{\text{restore}} \propto 1/\lambda_{\text{ZQP}}$ —measure the recovery time under strong perturbation.
6. **MES scaling laws:** the local coupling speed c and effective inertia m will follow $c \propto |R|^{1/2}$ and $m \propto |R|^{-1/2}$ —measure via spatially resolved spectroscopy and transport.
7. **No Cooper pairs:** the condensed manifold does not require electron pairing; experimental signatures of pairing (e.g., Josephson effect) will be absent, replaced by geometric phase signatures.

2.17 14. The T0-T6 Engineering Roadmap

Task	Theoretical basis	Engineering goal
T0: Recovery field estimation	$\Gamma_{\text{ZQP}} \approx 2.4\lambda_{\text{ZQP}}$	Extract λ_{ZQP} from ARPES/STM data
T1: NSAZ localization	Negative curvature attractors ($R < 0$)	Identify $R(\mathbf{r}) < 0$ regions via STM/STS
T2: Z constraint encoding	$\mathbb{Z}_5$ spectral duality	Encode the six Ju operations as state-transition rules

Task	Theoretical basis	Engineering goal
T3: OM functional construction	Generalized GL for tensor fields	Construct cost function for manifold condensation
T4: Core solver	Optimal transport from $\mathcal{F}[\mathcal{G}]$	Solve for paths escaping normal-state traps
T5: Intervention targets	Acupoints (negative curvature extrema)	Find minimal perturbations maximizing curvature flattening
T6: Uncertainty quantification	Topological protection robustness	Validate via κ (trajectory divergence) and η_{Rx} (reprogramming efficiency)

2.18 15. Design Principles: From Material Screening to System Development

The redefinition shifts the goal from **material screening** to **system development**. The design architecture has five pillars:

1. **Deep anchor**: maximize Γ and the condensation gap so that $\Delta \gg k_B T_{\text{room}}$.
2. **Adaptive self-calibration**: design the system to maintain the zeroing measurement cycle through ZQP feedback.
3. **Topological protection**: imprint a robust topological invariant during the developmental stage.
4. **Spatial-omics optimization**: use metric tensor extraction to identify acupoints and optimize the geometry.
5. **Cosmological compatibility**: ensure that the condensation parameters satisfy the MES scaling laws of the universal curvature network.

Deep anchor + Adaptive self-calibration + Topological protection + Spatial-omics optimization + Cosmological compatibility

The MES scaling laws provide the quantitative criterion for the fifth pillar:

$$|R|_{\text{sat}} \gg \frac{k_B T_{\text{room}}}{\Delta_0}$$

2.19 16. Conclusion

We have redefined room-temperature superconductivity as **the developmental terminal state of a Generalized Adaptive Intelligent System (GAIS)—the topological condensation of a dynamic equilibrium quantum manifold (DEQM)**. The superconducting state is not the pairing of electrons but the global topological ordering of a **unified 3D quantum geometric field** $\mathcal{G}_{\mu\nu}(x, t)$, maintained through the **ZQP mechanism** in a sustained **zeroing measurement** cycle, protected by topological invariants, driven by nonlocal coupling, and directly probed by the **anomalous Hall effect (AHE)**.

The algebraic foundation is the $\mathbb{Z}_5$ spectral duality $\sigma(K) = \sigma(S)^2$; the physical mechanism is the ZQP coupling $\Gamma_{\text{ZQP}} \approx 2.4\lambda_{\text{ZQP}}$ with nonlinear saturating negative feedback, previously validated through V4.52 numerical simulations. The complete computable pipeline $\mathbf{R}_I \xrightarrow{\text{DFPT}} \Omega_{\mu\nu}(\mathbf{k}) \xrightarrow{\text{ZQP}} g_{\mu\nu}(\mathbf{k}) \xrightarrow{\text{Morse}} \mathcal{G}_{\mu\nu}(\mathbf{x}, t)$ maps discrete atomic coordinates to the continuous quantum geometric field, providing the microscopic foundation for the DEQM.

Applying the **Modified Einstein Spherical (MES) scaling laws** $c \propto |R|^{1/2}$ and $m \propto |R|^{-1/2}$, we have redefined c as the **coupling speed on the ZQP dynamic equilibrium quantum manifold** and m as the **local effective inertia of the open quantum geometric field under the constraint of the universal curvature network**. These redefinitions provide a rigorous dynamical criterion for “acupoints” as high-curvature, low-inertia leverage points, and for the transformation of “gravity wells” into “deep anchors” by ZQP saturation.

The Cosmic Engineering Architecture provides a unified language for superconductivity across all temperature scales. It shifts the goal from **finding** a material to **developing** a system. The five design pillars—deep anchor, adaptive self-calibration, topological protection, spatial-omics optimization, and cosmological compatibility—offer a concrete path toward room-temperature superconductivity.

Remark on universality. The geometric developmental framework presented in this work — the MES scaling laws $c \propto |R|^{1/2}$, $m \propto |R|^{-1/2}$, the three-parameter dynamics of driving (α_0), constraint (β), and coherence (γ), and the Morse bifurcation from geometric chaos to DEQM condensation — is not restricted to the condensed-matter systems that motivated its construction. We conjecture that it constitutes a **universal developmental grammar** of all **Generalized Adaptive Intelligent System (GAIS)**, wherever such systems form a **DEQM seed** from which adaptive intelligence emerges through curvature-driven topological condensation. The Sun and other stars **may be viewed as** astronomical-scale instances: each is a self-gravitating DEQM whose ignition, steady-state burning, and terminal disintegration **may follow** the same geometric logic as the superconducting transition. The cosmic web — **the largest known curvature web** of filaments, nodes, and voids — is the supreme cosmological-scale instance: a curvature manifold whose topological architecture **may constrain** the mass, position, and developmental fate of every stellar embryo embedded within it. From the nanometer-scale DEQM seed to the gigaparsec-scale cosmic web, **the same geometric logic may recur** across thirty orders of magnitude. Geometric development is not a theory *of* superconductivity; it is a theory *through which* superconductivity, stellar embryogenesis, and the self-organization of the cosmos **may reveal** their shared identity.

Room-temperature superconductivity is not a material to be found; it is a quantum manifold to be condensed.

常温超导，非材料之寻，乃量子流形之凝。

2.20 Acknowledgments

We are grateful to all the individual scientists and teams of scientists who have contributed to the exploration and understanding of the entire universe. Supported by AI-driven Supercomputers and the MES Universe Project. We have benefited from our amazing AI partners.

2.21 Funding

Not applicable

2.22 Availability of data and material

Code, mathematical scripts, and preprints are available at GitHub: <https://github.com/Zaitian001/-Self-Preprint-V1.1->

2.23 Competing interests

Not applicable

2.24 Authors Contributions

Not applicable

2.25 References

- [1] Baoliin (Zaitian) Wu 伍宝林 (2026). Physical AGI Dao: Nonlinear Symmetry Anchor Zero as the Theoretical Foundation of Adaptive Intelligence World Model. Self-Preprint Archive. <https://Zaitian001.github.io/-Self-Preprint-V1.1-/preprints/DW58553906.html>
- [2] Baoliin (Zaitian) Wu 伍宝林. (2026). The Geometry of Qi: A Riemannian Architecture for Physiological Information Flow and Meridian Emergence. Self-Preprint Archive. <https://Zaitian001.github.io/-Self-Preprint-V1.1-/preprints/DW58553908.html>
- [3] Baoliin (Zaitian) Wu 伍宝林. (2026). Stochastic Geometric Dynamics of Immune Memory, From Curvature-Driven Attractors to Testable Predictions. Self-Preprint Archive. <https://Zaitian001.github.io/-Self-Preprint-V1.1-/preprints/DW58553904.html>
- [4] Baoliin (Zaitian) Wu 伍宝林. (2026). Adaptive Intelligence Dao: A Unified Mathematical Principle of Anchor Zero, Change, Balance, Emergence, and Self-Organization. Self-Preprint Archive. <https://Zaitian001.github.io/-Self-Preprint-V1.1-/preprints/DW58553907.html>
- [5] Baoliin (Zaitian) Wu 伍宝林. (2026). The Overlooked Coordinate System Exorcising the Demons and Ghosts of Quantum Measurement. Self-Preprint Archive. <https://Zaitian001.github.io/-Self-Preprint-V1.1-/preprints/DW58553901.html>
- [6] Baoliin (Zaitian) Wu 伍宝林. (2026). The Gui-Ju Dao: The Artificial General Intelligence Algorithm, Geometry, and Engineering Paradigm of China's Indigenous Closed Spherical Universe. Self-Preprint Archive. <https://zaitian001.github.io/-Self-Preprint-V1.1-/preprints/DW58553902.html>

- [7] Baoliin (Zaitian) Wu 伍宝林. (2026). The Bone Porcelain Core Hypothesis of the 1600 Year Old Delhi Iron Pillar: An Adaptive Intelligent Metamaterial Invention Driven by Internal Sustained Release. Self-Preprint Archive. <https://Zaitian001.github.io/-Self-Preprint-V1.1-/preprints/DW58553903.html>
- [8] Baoliin (Zaitian) Wu 伍宝林. (2026). Wuxing Dao. Self-Preprint Archive. <https://Zaitian001.github.io/-Self-Preprint-V1.1-/preprints/DW58553905.html>
- [9] Baoliin (Zaitian) Wu 伍宝林. (2026). Geometric Pathology: Topological Collapse and Landscape Reprogramming in Complex Adaptive Systems. Self-Preprint Archive. <https://Zaitian001.github.io/-Self-Preprint-V1.1-/preprints/DW58553909.html>
- [10] Baoliin (Zaitian) Wu 伍宝林. (2026). Groma-Ju Dao: The Six Modes of the Zhoubi Suanjin “Ju” as a Geometric Operator System. Self-Preprint Archive. <https://Zaitian001.github.io/-Self-Preprint-V1.1-/preprints/DW58553910.html>
- [11] Baoliin (Zaitian) Wu. (2025). The Pure Geometric Origin of Mass and Light. Preprints. <https://doi.org/10.20944/preprints202505.2249.v1>
- [12] Baoliin (Zaitian) Wu. (2025). The Return to the Einstein Spherical Universe Model. Preprints. <https://doi.org/10.20944/preprints202501.2189.v1>
- [13] Baoliin (Zaitian) Wu. (2025). The Return to the Einstein Spherical Universe: The Dawning Moment of a New Cosmic Science. Preprints. <https://doi.org/10.20944/preprints202504.0727.v2>
- [14] Bender, C. M., & Boettcher, S. (1998). Real spectra in non-Hermitian Hamiltonians having $\mathcal{PT}$ symmetry. Physical Review Letters, 80(24), 5243–5246. <https://doi.org/10.1103/PhysRevLett.80.5243>
- [15] Eliashberg, G. M. (1960). Interactions between electrons and lattice vibrations in a superconductor. Zhurnal Eksperimental’ noi i Teoreticheskoi Fiziki, 38(3), 966–976. Soviet Physics JETP, 11(3), 696–702. <http://www.jetp.ras.ru/cgi-bin/e/index/e/11/3/p696?a=list>
- [16] Ginzburg, V. L., & Landau, L. D. (1950). On the theory of superconductivity. Zhurnal Eksperimental’ noi i Teoreticheskoi Fiziki, 20, 1064–1082. English translation in Collected Papers of L. D. Landau, pp. 546–568. <https://doi.org/10.1016/B978-0-08-010586-4.50078-X>
- [17] Kang, M., Kim, S., Qian, Y. et al. (2025). Measurements of the quantum geometric tensor in solids. Nat. Phys. 21, 110–117. <https://doi.org/10.1038/s41567-024-02678-8>
- [18] Thouless, D. J., Kohmoto, M., Nightingale, M. P., and den Nijs, M. (1982). Quantized Hall conductance in a two-dimensional periodic potential. Physical Review Letters, 49(6), 405–408. <https://doi.org/10.1103/PhysRevLett.49.405>
- [19] Xiao, D., Chang, M.-C., and Niu, Q. (2010). Berry phase effects on electronic properties. Reviews of Modern Physics, 82(3), 1959–2007. <https://doi.org/10.1103/RevModPhys.82.1959>
- [20] Drozdov, A.P., Kong, P.P., Minkov, V.S. et al. (2019). Superconductivity at 250 K in lanthanum hydride under high pressures. Nature, 569(7757), 528–531. <https://doi.org/10.1038/s41586-019-1201-8>
- [21] Ming, X., Zhang, YJ., Zhu, X. et al. (2023). Absence of near-ambient superconductivity in $\text{LuH}_{2\pm x}\text{N}_y$. Nature, 620(7972), 72–77. <https://doi.org/10.1038/s41586-023-06162-w>
- [22] Sun, H., Huo, M., Hu, X. et al. (2023). Signatures of superconductivity near 80 K in a nickelate under high pressure. Nature, 621(7979), 493–498. <https://doi.org/10.1038/s41586-023-06408-7>

- [23] Li, D., Lee, K., Wang, B. Y. et al. (2019). Superconductivity in an infinite-layer nickelate. *Nature*, 572(7771), 624–627. <https://doi.org/10.1038/s41586-019-1496-5>
- [24] Peotta, S., and Törmä, P. (2015). Superfluidity in topologically nontrivial flat bands. *Nature Communications*, 6(1), 8944. <https://doi.org/10.1038/ncomms9944>
- [25] Hamidian, M., Edkins, S., Joo, S. et al. (2016). Detection of a Cooper-pair density wave in $\text{Bi}_2\text{Sr}_2\text{CaCu}_2\text{O}_{8+x}$. *Nature*, 532(7599), 343–347. <https://doi.org/10.1038/nature17411>
- [26] Yip, K. Y., Ho, K. O., Yu, K. Y. et al. (2019). Measuring magnetic field gradients in a diamond anvil cell using nitrogen-vacancy centers. *Science*, 366(6471), 1355–1359. <https://doi.org/10.1126/science.aaw4278>
- [27] Cao, Y., Fatemi, V., Demir, A., Fang, S. et al. (2018). Unconventional superconductivity in magic-angle graphene superlattices. *Nature*, 556(7699), 43–50. <https://doi.org/10.1038/nature26160>
- [28] Flores-Livas, J. A., Boeri, L., Sanna, A. et al. (2020). A perspective on conventional high-temperature superconductivity at high pressure. *Physics Reports*, 856, 1–65. <https://doi.org/10.1016/j.physrep.2020.02.003>
- [29] Fogelström, M., Park, W. K., Greene, L. H. et al. (2010). Point-contact spectroscopy in heavy-fermion superconductors. *Physical Review B* 82, 014527. <https://doi.org/10.1103/PhysRevB.82.014527>

2.26 Physical Proof of Existence (Appendix)

- **Banknote Serial:** DW58553911
- **SHA-256 Hash:** 66bef64d5ee3a4c2692665819fb74777e08d0f4a44890db0e4ba09490d2b30c9
- **Verification URL:** <https://Zaitian001.github.io/-Self-Preprint-V1.1-/preprints/DW58553911.html>

This document was timestamped via the Self-Preprint system.