

Chapter 1

Universality of multi-particle scattering

Hard, but worthwhile

1.1 Multi-particle quantum walk

Note that this is exactly what I wanted to talk about.

Very difficult in general.

1.1.1 Two-particle scattering on an infinite path

The one thing we can actually compute It might be interesting to talk about what happens with spins.

1.2 Applying an encoded $C\theta$ -gate

1.2.1 Finite truncation

Theorem 1. *Let $H^{(2)}$ be a two-particle Hamiltonian of the form (??) with interaction range at most C , i.e., $\mathcal{V}(|r|) = 0$ for all $|r| > C$. Let $\theta_{\pm}(p_1, p_2)$ be given by equation (??). Define $\theta = \theta_{\pm}(\pi/4, 3\pi/8)$. Let $L \in \mathbb{N}$, let $M \in \{C+1, C+2, \dots\}$, and define*

$$\begin{aligned} |\chi_{z,k}\rangle &= \frac{1}{\sqrt{L}} \sum_{x=z-L}^{z-1} e^{ikx} |x\rangle \\ |\psi(0)\rangle &= \frac{1}{\sqrt{2}} \left(|\chi_{-M, -\frac{\pi}{2}}\rangle |\chi_{M+L+1, \frac{\pi}{4}}\rangle \pm |\chi_{M+L+1, \frac{\pi}{4}}\rangle |\chi_{-M, -\frac{\pi}{2}}\rangle \right). \end{aligned}$$

Let c_0 be a constant independent of L . Then, for all $0 \leq t \leq c_0 L$, we have

$$\left\| e^{-iH^{(2)}t} |\psi(0)\rangle - |\alpha(t)\rangle \right\| = \mathcal{O}(L^{-1/8}),$$

where

$$|\alpha(t)\rangle = \sum_{x,y} a_{xy}(t)|x,y\rangle, \quad (1.1)$$

$a_{xy}(t) = \pm a_{yx}(t)$, and, for $x \leq y$,

$$a_{xy}(t) = \frac{1}{\sqrt{2}L} e^{-\sqrt{2}it} \left[e^{-i\pi x/2} e^{i\pi y/4} F(x,y,t) \pm e^{i\theta} e^{i\pi x/4} e^{-i\pi y/2} F(y,x,t) \right] \quad (1.2)$$

where

$$F(u,v,t) = \begin{cases} 1 & \text{if } u - 2[t] \in \{-M-L, \dots, -M-1\} \text{ and } v + 2 \left\lfloor \frac{t}{\sqrt{2}} \right\rfloor \in \{M+1, \dots, M+L\} \\ 0 & \text{otherwise.} \end{cases}$$

In this section we prove [Theorem 1](#). The main proof appears in [Section 1.2.1.1](#), relying on several technical lemmas proved in [Section 1.2.1.2](#). The proof follows the method used in the single-particle case, which is based on the calculation from the appendix of reference [\[?\]](#).

Recall from [\(??\)](#) that for each $p_1 \in (-\pi, \pi)$ and $p_2 \in (0, \pi)$ there is an eigenstate $|\text{sc}(p_1; p_2)\rangle_{\pm}$ of $H^{(2)}$ of the form

$$\langle x, y | \text{sc}(p_1; p_2) \rangle_{\pm} = \frac{e^{-ip_1(\frac{x+y}{2})}}{\sqrt{2}} \begin{cases} e^{-ip_2(x-y)} \pm e^{i\theta_{\pm}(p_1, p_2)} e^{ip_2(x-y)} & \text{if } x - y \leq -C \\ e^{-ip_2(x-y)} e^{i\theta_{\pm}(p_1, p_2)} \pm e^{ip_2(x-y)} & \text{if } x - y \geq C \\ f(p_1, p_2, x - y) \pm f(p_1, p_2, y - x) & \text{if } |x - y| < C \end{cases} \quad (1.3)$$

where

$$e^{i\theta_{\pm}(p_1, p_2)} = T(p_1, p_2) \pm R(p_1, p_2),$$

C is the range of the interaction, T and R are the transmission and reflection coefficients of the interaction at the chosen momentum, f describes the amplitudes of the scattering state within the interaction range, and the \pm depends on the type of particle ($+$ for bosons, $-$ for fermions). The state $|\text{sc}(p_1; p_2)\rangle_{\pm}$ satisfies

$$H^{(2)} |\text{sc}(p_1; p_2)\rangle_{\pm} = 4 \cos \frac{p_1}{2} \cos p_2 |\text{sc}(p_1; p_2)\rangle_{\pm}$$

and is delta-function normalized as

$${}_{\pm} \langle \text{sc}(p'_1; p'_2) | \text{sc}(p_1; p_2) \rangle_{\pm} = 4\pi^2 \delta(p_1 - p'_1) \delta(p_2 - p'_2). \quad (1.4)$$

Proof. Expand $|\psi(0)\rangle$ in the basis of eigenstates of the Hamiltonian to get

$$|\psi(t)\rangle = e^{-iH^{(2)}t} |\psi(0)\rangle = |\psi_1(t)\rangle + |\psi_2(t)\rangle$$

where

$$|\psi_1(t)\rangle = \iint_{D_{\epsilon}} \frac{d\phi_1 d\phi_2}{4\pi^2} e^{-it4 \cos(\frac{p_1}{2} + \frac{\phi_1}{2}) \cos(p_2 + \phi_2)} |\text{sc}(p_1 + \phi_1; p_2 + \phi_2)\rangle_{\pm} ({}_{\pm} \langle \text{sc}(p_1 + \phi_1; p_2 + \phi_2) | \psi(0) \rangle)$$

with $D_\epsilon = [-\epsilon, \epsilon] \times [-\epsilon, \epsilon]$, $p_1 = \pi/2 - \pi/4 = \pi/4$, $p_2 = (\pi/2 + \pi/4)/2 = 3\pi/8$, and with $|\psi_2(t)\rangle$ orthogonal to $|\psi_1(t)\rangle$. We take $\epsilon = a/\sqrt{L}$ for some constant a . Using equation (1.3) we get

$$|\psi_1(t)\rangle = |\psi_A(t)\rangle \pm |\psi_B(t)\rangle$$

where

$$\begin{aligned} |\psi_A(t)\rangle &= \iint_{D_\epsilon} \frac{d\phi_1 d\phi_2}{4\pi^2} e^{-it4 \cos(\frac{\pi}{8} + \frac{\phi_1}{2}) \cos(\frac{3\pi}{8} + \phi_2)} A(\phi_1, \phi_2) |\text{sc}(\frac{\pi}{4} + \phi_1; \frac{3\pi}{8} + \phi_2)\rangle_\pm \\ |\psi_B(t)\rangle &= \iint_{D_\epsilon} \frac{d\phi_1 d\phi_2}{4\pi^2} e^{-it4 \cos(\frac{\pi}{8} + \frac{\phi_1}{2}) \cos(\frac{3\pi}{8} + \phi_2)} e^{-i\theta_\pm(\frac{\pi}{4} + \phi_1, \frac{3\pi}{8} + \phi_2)} B(\phi_1, \phi_2, \frac{3\pi}{8}) |\text{sc}(\frac{\pi}{4} + \phi_1; \frac{3\pi}{8} + \phi_2)\rangle_\pm \end{aligned} \quad (1.5)$$

with

$$\begin{aligned} A(\phi_1, \phi_2) &= \frac{1}{L} \sum_{x=-(M+L)}^{-(M+1)} \sum_{y=M+1}^{M+L} e^{i\phi_1 \frac{x+y}{2}} e^{i\phi_2(x-y)} \\ B(\phi_1, \phi_2, k) &= \frac{1}{L} \sum_{x=-(M+L)}^{-(M+1)} \sum_{y=M+1}^{M+L} e^{i\phi_1 \frac{x+y}{2}} e^{i(\phi_2 + 2k)(y-x)}. \end{aligned} \quad (1.6)$$

Using the delta-function normalization of the scattering states (equation (1.4)) we get

$$\begin{aligned} \langle \psi_B(t) | \psi_B(t) \rangle &= \iint_{D_\epsilon} \frac{d\phi_1 d\phi_2}{4\pi^2} |B(\phi_1, \phi_2, \frac{3\pi}{8})|^2 \\ &\leq \frac{16\pi^2}{L^2 \epsilon^2} \end{aligned}$$

by Lemma 3 (as long as $\epsilon < 3\pi/8$, which holds for L sufficiently large). Similarly,

$$\begin{aligned} 1 &\geq \langle \psi_A(t) | \psi_A(t) \rangle \\ &= \iint_{D_\epsilon} \frac{d\phi_1 d\phi_2}{4\pi^2} |A(\phi_1, \phi_2)|^2 \\ &\geq 1 - \frac{4\pi}{L\epsilon} \end{aligned}$$

(from the first two facts in Lemma 3) and therefore

$$\begin{aligned} \langle \psi_1(t) | \psi_1(t) \rangle &= \langle \psi_A(t) | \psi_A(t) \rangle + \langle \psi_B(t) | \psi_B(t) \rangle + \langle \psi_A(t) | \psi_B(t) \rangle + \langle \psi_B(t) | \psi_A(t) \rangle \\ &\geq 1 - \frac{4\pi}{L\epsilon} - 2 |\langle \psi_A(t) | \psi_B(t) \rangle| \\ &\geq 1 - \frac{4\pi}{L\epsilon} - 2 |\langle \psi_A(t) | \psi_A(t) \rangle|^{\frac{1}{2}} |\langle \psi_B(t) | \psi_B(t) \rangle|^{\frac{1}{2}} \\ &\geq 1 - \frac{12\pi}{L\epsilon}. \end{aligned}$$

Hence

$$\langle \psi_2(t) | \psi_2(t) \rangle \leq \frac{12\pi}{L\epsilon}$$

since

$$\langle \psi(t) | \psi(t) \rangle = \langle \psi_1(t) | \psi_1(t) \rangle + \langle \psi_2(t) | \psi_2(t) \rangle = 1.$$

Thus

$$\begin{aligned} \| |\psi(t)\rangle - |\psi_A(t)\rangle \| &= \| |\psi_B(t)\rangle + |\psi_2(t)\rangle \| \\ &\leq \| |\psi_B(t)\rangle \| + \| |\psi_2(t)\rangle \| \\ &\leq \frac{4\pi}{L\epsilon} + \sqrt{\frac{12\pi}{L\epsilon}}. \end{aligned}$$

Now

$$\begin{aligned} \| |\psi(t)\rangle - |\alpha(t)\rangle \| &\leq \| |\psi(t)\rangle - |\psi_A(t)\rangle \| + \| |\psi_A(t)\rangle - |\alpha(t)\rangle \| \\ &\leq \frac{4\pi}{L\epsilon} + \sqrt{\frac{12\pi}{L\epsilon}} + \| |\psi_A(t)\rangle - |\alpha(t)\rangle \| \\ &= \mathcal{O}(L^{-1/4}) + \| |\psi_A(t)\rangle - |\alpha(t)\rangle \| \end{aligned}$$

using our choice $\epsilon = a/\sqrt{L}$. To complete the proof, we now show that the second term in this expression is bounded by $\mathcal{O}(L^{-1/8})$.

Lemma 1. *With $|\psi_A(t)\rangle$ and $|\alpha(t)\rangle$ defined through equations (1.5) and (1.1), with $t \leq c_0 L$ (for some constant c_0),*

$$\| |\psi_A(t)\rangle - |\alpha(t)\rangle \| = \mathcal{O}(L^{-1/8}).$$

Proof. To simplify matters, note that both $|\psi_A(t)\rangle$ and $|\alpha(t)\rangle$ are either symmetric or anti-symmetric (i.e., $\langle x, y | \alpha(t) \rangle = \pm \langle y, x | \alpha(t) \rangle$ and $\langle x, y | \psi_A(t) \rangle = \pm \langle y, x | \psi_A(t) \rangle$). Taking C to be the maximum range of the interaction in our Hamiltonian, we have

$$\| |\psi_A(t)\rangle - |\alpha(t)\rangle \| \leq 2 \| P_1 |\psi_A(t)\rangle - P_1 |\alpha(t)\rangle \| + \| P_2 |\psi_A(t)\rangle \| + \| P_2 |\alpha(t)\rangle \|,$$

where

$$P_1 = \sum_{y-x \geq C} |x, y\rangle \langle x, y| \quad P_2 = \sum_{|x-y| < C} |x, y\rangle \langle x, y|.$$

Now, for $y - x \geq C$,

$$\begin{aligned} \langle x, y | \psi_A(t) \rangle &= \iint_{D_\epsilon} \frac{d\phi_1 d\phi_2}{4\pi^2} e^{-it4 \cos(\frac{\pi}{8} + \frac{\phi_1}{2}) \cos(\frac{3\pi}{8} + \phi_2)} A(\phi_1, \phi_2) \frac{e^{-i(\frac{\pi}{4} + \phi_1)(\frac{x+y}{2})}}{\sqrt{2}} \\ &\quad \left(e^{i(\frac{3\pi}{8} + \phi_2)(y-x)} \pm e^{-i(\frac{3\pi}{8} + \phi_2)(y-x) + i\theta_\pm(\frac{\pi}{4} + \phi_1, \frac{3\pi}{8} + \phi_2)} \right) \\ &= \iint_{D_\epsilon} \frac{d\phi_1 d\phi_2}{4\pi^2} \left[\frac{1}{\sqrt{2}} e^{-it4 \cos(\frac{\pi}{8} + \frac{\phi_1}{2}) \cos(\frac{3\pi}{8} + \phi_2)} A(\phi_1, \phi_2) \right. \\ &\quad \left(e^{-i\pi x/2} e^{i\pi y/4} e^{-i\phi_1(\frac{x+y}{2})} e^{i\phi_2(y-x)} \right. \\ &\quad \left. \left. \pm e^{i\pi x/4} e^{-i\pi y/2} e^{-i\phi_1(\frac{x+y}{2})} e^{-i\phi_2(y-x)} e^{i\theta_\pm(\frac{\pi}{4} + \phi_1, \frac{3\pi}{8} + \phi_2)} \right) \right]. \end{aligned}$$

From Lemma 4, for $x \leq y$, the state $|\alpha(t)\rangle$ takes the form

$$\begin{aligned} \langle x, y | \alpha(t) \rangle = & \frac{1}{\sqrt{2}} e^{-it\sqrt{2}} \left[e^{-i\pi x/2} e^{i\pi y/4} \left(\iint_{D_\pi} \frac{d\phi_1 d\phi_2}{4\pi^2} \right. \right. \\ & A(\phi_1, \phi_2) e^{-i\phi_1(-[t] + \lfloor \frac{t}{\sqrt{2}} \rfloor + \frac{x+y}{2})} e^{-2i\phi_2(-[t] - \lfloor \frac{t}{\sqrt{2}} \rfloor + \frac{x-y}{2})} \\ & \pm e^{i\theta} e^{i\pi x/4} e^{-i\pi y/2} \left(\iint_{D_\pi} \frac{d\phi_1 d\phi_2}{4\pi^2} \right. \\ & \left. \left. A(\phi_1, \phi_2) e^{-i\phi_1(-[t] + \lfloor \frac{t}{\sqrt{2}} \rfloor + \frac{x+y}{2})} e^{-2i\phi_2(-[t] - \lfloor \frac{t}{\sqrt{2}} \rfloor + \frac{x-y}{2})} \right) \right], \end{aligned}$$

where $D_\pi = [-\pi, \pi] \times [-\pi, \pi]$. Using these expressions for $|\psi_A(t)\rangle$ and $|\alpha(t)\rangle$, we now write

$$P_1 |\psi_A(t)\rangle - P_1 |\alpha(t)\rangle = \pm |e_1(t)\rangle + |e_2(t)\rangle \pm |f_1(t)\rangle + |f_2(t)\rangle \pm |g_1(t)\rangle + |g_2(t)\rangle \pm |h(t)\rangle$$

where each term in the above equation is supported only on states $|x, y\rangle$ such that $y - x \geq C$, and (for $y - x \geq C$)

$$\begin{aligned} \langle x, y | e_1(t) \rangle &= \frac{e^{i\theta}}{\sqrt{2}} e^{-it\sqrt{2}} e^{i\pi x/4} e^{-i\pi y/2} \iint_{D_\pi} \frac{d\phi_1 d\phi_2}{4\pi^2} A(\phi_1, \phi_2) \left[e^{-i\phi_1(-t + \frac{t}{\sqrt{2}} + \frac{x+y}{2})} \right. \\ & \quad \left. e^{-2i\phi_2(-t - \frac{t}{\sqrt{2}} + \frac{y-x}{2})} - e^{-i\phi_1(-[t] + \lfloor \frac{t}{\sqrt{2}} \rfloor + \frac{x+y}{2})} e^{-2i\phi_2(-[t] - \lfloor \frac{t}{\sqrt{2}} \rfloor + \frac{y-x}{2})} \right] \\ \langle x, y | e_2(t) \rangle &= \frac{1}{\sqrt{2}} e^{-it\sqrt{2}} e^{-i\pi x/2} e^{i\pi y/4} \iint_{D_\pi} \frac{d\phi_1 d\phi_2}{4\pi^2} A(\phi_1, \phi_2) \left[e^{-i\phi_1(-t + \frac{t}{\sqrt{2}} + \frac{x+y}{2})} \right. \\ & \quad \left. e^{-2i\phi_2(-t - \frac{t}{\sqrt{2}} + \frac{x-y}{2})} - e^{-i\phi_1(-[t] + \lfloor \frac{t}{\sqrt{2}} \rfloor + \frac{x+y}{2})} e^{-2i\phi_2(-[t] - \lfloor \frac{t}{\sqrt{2}} \rfloor + \frac{x-y}{2})} \right] \\ \langle x, y | f_1(t) \rangle &= -\frac{e^{i\theta}}{\sqrt{2}} e^{-it\sqrt{2}} e^{i\pi x/4} e^{-i\pi y/2} \iint_{D_\pi \setminus D_\epsilon} \frac{d\phi_1 d\phi_2}{4\pi^2} A(\phi_1, \phi_2) \\ & \quad e^{-i\phi_1(-t + \frac{t}{\sqrt{2}} + \frac{x+y}{2})} e^{-2i\phi_2(-t - \frac{t}{\sqrt{2}} + \frac{y-x}{2})} \\ \langle x, y | f_2(t) \rangle &= -\frac{1}{\sqrt{2}} e^{-it\sqrt{2}} e^{-i\pi x/2} e^{i\pi y/4} \iint_{D_\pi \setminus D_\epsilon} \frac{d\phi_1 d\phi_2}{4\pi^2} A(\phi_1, \phi_2) \\ & \quad e^{-i\phi_1(-t + \frac{t}{\sqrt{2}} + \frac{x+y}{2})} e^{-2i\phi_2(-t - \frac{t}{\sqrt{2}} + \frac{x-y}{2})} \\ \langle x, y | g_1(t) \rangle &= \frac{e^{i\theta}}{\sqrt{2}} e^{i\pi x/4} e^{-i\pi y/2} \iint_{D_\epsilon} \frac{d\phi_1 d\phi_2}{4\pi^2} A(\phi_1, \phi_2) e^{-i\phi_1(\frac{x+y}{2})} e^{-2i\phi_2(\frac{y-x}{2})} \\ & \quad \left[e^{-it4 \cos(\frac{\pi}{8} + \frac{\phi_1}{2}) \cos(\frac{3\pi}{8} + \phi_2)} - e^{-it(\sqrt{2} + \sqrt{2}(\frac{\phi_1}{2} - \phi_2) - 2(\frac{\phi_1}{2} + \phi_2))} \right] \\ \langle x, y | g_2(t) \rangle &= \frac{1}{\sqrt{2}} e^{-i\pi x/2} e^{i\pi y/4} \iint_{D_\epsilon} \frac{d\phi_1 d\phi_2}{4\pi^2} A(\phi_1, \phi_2) e^{-i\phi_1(\frac{x+y}{2})} e^{-2i\phi_2(\frac{x-y}{2})} \\ & \quad \left[e^{-it4 \cos(\frac{\pi}{8} + \frac{\phi_1}{2}) \cos(\frac{3\pi}{8} + \phi_2)} - e^{-it(\sqrt{2} + \sqrt{2}(\frac{\phi_1}{2} - \phi_2) - 2(\frac{\phi_1}{2} + \phi_2))} \right] \\ \langle x, y | h(t) \rangle &= \frac{1}{\sqrt{2}} e^{i\pi x/4} e^{-i\pi y/2} \iint_{D_\epsilon} \frac{d\phi_1 d\phi_2}{4\pi^2} A(\phi_1, \phi_2) e^{-i\phi_1(\frac{x+y}{2})} e^{-2i\phi_2(\frac{y-x}{2})} \\ & \quad e^{-it4 \cos(\frac{\pi}{8} + \frac{\phi_1}{2}) \cos(\frac{3\pi}{8} + \phi_2)} \left(e^{i\theta \pm (\frac{\pi}{4} + \phi_1, \frac{3\pi}{8} + \phi_2)} - e^{i\theta} \right). \end{aligned}$$

We now proceed to bound the norm of each of these states. We repeatedly use the fact that, for $(\phi_1, \phi_2) \in D_\pi$,

$$\sum_{x,y=-\infty}^{\infty} e^{ix(\frac{1}{2}(\phi_1-\tilde{\phi}_1)-(\phi_2-\tilde{\phi}_2))} e^{iy(\frac{1}{2}(\phi_1-\tilde{\phi}_1)+(\phi_2-\tilde{\phi}_2))} = 4\pi^2 \delta(\phi_1 - \tilde{\phi}_1) \delta(\phi_2 - \tilde{\phi}_2).$$

Using this formula we get

$$\begin{aligned} \langle e_1(t) | e_1(t) \rangle &= \sum_{y-x \geq C} \langle e_1(t) | x, y \rangle \langle x, y | e_1(t) \rangle \\ &\leq \sum_{x=-\infty}^{\infty} \sum_{y=-\infty}^{\infty} \left| \frac{1}{\sqrt{2}} \iint_{D_\pi} \frac{d\phi_1 d\phi_2}{4\pi^2} A(\phi_1, \phi_2) \left[e^{-i\phi_1(-t+\frac{t}{\sqrt{2}}+\frac{x+y}{2})} \right. \right. \\ &\quad \left. \left. e^{-2i\phi_2(-t-\frac{t}{\sqrt{2}}+\frac{y-x}{2})} - e^{-i\phi_1(-[t]+\lfloor \frac{t}{\sqrt{2}} \rfloor + \frac{x+y}{2})} e^{-2i\phi_2(-[t]-\lfloor \frac{t}{\sqrt{2}} \rfloor + \frac{y-x}{2})} \right] \right|^2 \\ &= \frac{1}{2} \iint_{D_\pi} \frac{d\phi_1 d\phi_2}{4\pi^2} |A(\phi_1, \phi_2)|^2 \left| e^{-i\phi_1(-t+\frac{t}{\sqrt{2}})} e^{-2i\phi_2(-t-\frac{t}{\sqrt{2}})} \right. \\ &\quad \left. - e^{-i\phi_1(-[t]+\lfloor \frac{t}{\sqrt{2}} \rfloor)} e^{-2i\phi_2(-[t]-\lfloor \frac{t}{\sqrt{2}} \rfloor)} \right|^2. \end{aligned}$$

Now use the fact that $|e^{-ic} - 1|^2 \leq c^2$ for $c \in \mathbb{R}$ to get

$$\begin{aligned} \langle e_1(t) | e_1(t) \rangle &\leq \frac{1}{2} \iint_{D_\pi} \left(\frac{d\phi_1 d\phi_2}{4\pi^2} \right) |A(\phi_1, \phi_2)|^2 \left(-\phi_1 \left(-t + \frac{t}{\sqrt{2}} + [t] - \left\lfloor \frac{t}{\sqrt{2}} \right\rfloor \right) \right. \\ &\quad \left. - 2\phi_2 \left(-t - \frac{t}{\sqrt{2}} + [t] + \left\lfloor \frac{t}{\sqrt{2}} \right\rfloor \right) \right)^2 \\ &\leq 4 \iint_{D_\pi} \frac{d\phi_1 d\phi_2}{4\pi^2} |A(\phi_1, \phi_2)|^2 (\phi_1^2 + 4\phi_2^2) \end{aligned}$$

using the Cauchy-Schwarz inequality and the fact that $|t - t/\sqrt{2} - [t] - \lfloor t/\sqrt{2} \rfloor| \leq 2$. So

$$\begin{aligned} \langle e_1(t) | e_1(t) \rangle &\leq 4 \left(\iint_{D_\pi \setminus D_\epsilon} \frac{d\phi_1 d\phi_2}{4\pi^2} + \iint_{D_\epsilon} \frac{d\phi_1 d\phi_2}{4\pi^2} \right) |A(\phi_1, \phi_2)|^2 (\phi_1^2 + 4\phi_2^2) \\ &\leq 4 (5\pi^2) \left(\frac{4\pi}{L\epsilon} \right) + 20\epsilon^2 \\ &= \frac{80\pi^3}{L\epsilon} + 20\epsilon^2 \end{aligned}$$

where we have used [Lemma 3](#) and the fact that $\phi_1^2 + 4\phi_2^2 \leq 5\epsilon^2$ on D_ϵ . Similarly,

$$\langle e_2(t) | e_2(t) \rangle \leq \frac{80\pi^3}{L\epsilon} + 20\epsilon^2.$$

Now

$$\begin{aligned}\langle f_1(t)|f_1(t)\rangle &\leq \frac{1}{2} \iint_{D_\pi \setminus D_\epsilon} \frac{d\phi_1 d\phi_2}{4\pi^2} |A(\phi_1, \phi_2)|^2 \\ &\leq \frac{2\pi}{L\epsilon}\end{aligned}$$

by [Lemma 3](#), and similarly

$$\langle f_2(t)|f_2(t)\rangle \leq \frac{2\pi}{L\epsilon}.$$

Moving on to the next term,

$$\begin{aligned}\langle g_1(t)|g_1(t)\rangle &\leq \frac{1}{2} \iint_{D_\epsilon} \frac{d\phi_1 d\phi_2}{4\pi^2} |A(\phi_1, \phi_2)|^2 \left| e^{-it4 \cos(\frac{\pi}{8} + \frac{\phi_1}{2}) \cos(\frac{3\pi}{8} + \phi_2)} \right. \\ &\quad \left. - e^{-it(\sqrt{2} + \sqrt{2}(\frac{\phi_1}{2} - \phi_2) - 2(\frac{\phi_1}{2} + \phi_2))} \right|^2 \\ &\leq \frac{1}{2} \iint_{D_\epsilon} \frac{d\phi_1 d\phi_2}{4\pi^2} \left[|A(\phi_1, \phi_2)|^2 t^2 \left(4 \cos\left(\frac{\pi}{8} + \frac{\phi_1}{2}\right) \cos\left(\frac{3\pi}{8} + \phi_2\right) \right. \right. \\ &\quad \left. \left. - \sqrt{2} - \sqrt{2}\left(\frac{\phi_1}{2} - \phi_2\right) + 2\left(\frac{\phi_1}{2} + \phi_2\right) \right)^2 \right] \quad (1.7)\end{aligned}$$

using $|e^{-ic} - 1|^2 \leq c^2$ for $c \in \mathbb{R}$. Now

$$\begin{aligned}4 \cos\left(\frac{\pi}{8} + \frac{\phi_1}{2}\right) \cos\left(\frac{3\pi}{8} + \phi_2\right) &= 2 \cos\left(\frac{\pi}{2} + \frac{\phi_1}{2} + \phi_2\right) + 2 \cos\left(-\frac{\pi}{4} + \frac{\phi_1}{2} - \phi_2\right) \\ &= -2 \sin\left(\frac{\phi_1}{2} + \phi_2\right) + \sqrt{2} \cos\left(\frac{\phi_1}{2} - \phi_2\right) + \sqrt{2} \sin\left(\frac{\phi_1}{2} - \phi_2\right)\end{aligned}$$

so

$$\begin{aligned}&\left| 4 \cos\left(\frac{\pi}{8} + \frac{\phi_1}{2}\right) \cos\left(\frac{3\pi}{8} + \phi_2\right) - \sqrt{2} - \sqrt{2}\left(\frac{\phi_1}{2} - \phi_2\right) + 2\left(\frac{\phi_1}{2} + \phi_2\right) \right| \\ &\leq \left| \sqrt{2} \left(\cos\left(\frac{\phi_1}{2} - \phi_2\right) - 1 \right) \right| + \left| \sqrt{2} \left(\sin\left(\frac{\phi_1}{2} - \phi_2\right) - \left(\frac{\phi_1}{2} - \phi_2\right) \right) \right| \\ &\quad + \left| 2 \left(\sin\left(\frac{\phi_1}{2} + \phi_2\right) - \left(\frac{\phi_1}{2} + \phi_2\right) \right) \right| \\ &\leq \sqrt{2} \left(\frac{\phi_1}{2} - \phi_2 \right)^2 + \sqrt{2} \left(\frac{\phi_1}{2} - \phi_2 \right)^2 + 2 \left(\frac{\phi_1}{2} + \phi_2 \right)^2 \\ &\leq 4 \left(\left(\frac{\phi_1}{2} + \phi_2 \right)^2 + \left(\frac{\phi_1}{2} - \phi_2 \right)^2 \right),\end{aligned}$$

using $|\cos x - 1| \leq x^2$ and $|\sin x - x| \leq x^2$ for $x \in \mathbb{R}$. Plugging this into equation (1.7) we get

$$\begin{aligned}
\langle g_1(t) | g_1(t) \rangle &\leq \frac{1}{2} \iint_{D_\epsilon} \frac{d\phi_1 d\phi_2}{4\pi^2} 16 |A(\phi_1, \phi_2)|^2 t^2 \left(\left(\frac{\phi_1}{2} + \phi_2 \right)^2 + \left(\frac{\phi_1}{2} - \phi_2 \right)^2 \right)^2 \\
&\leq 16t^2 \iint_{D_\epsilon} \frac{d\phi_1 d\phi_2}{4\pi^2} |A(\phi_1, \phi_2)|^2 \left(\left(\frac{\phi_1}{2} + \phi_2 \right)^4 + \left(\frac{\phi_1}{2} - \phi_2 \right)^4 \right) \\
&\leq \frac{16t^2}{L^2} \iint_{D_\epsilon} \frac{d\phi_1 d\phi_2}{4\pi^2} \frac{\sin^2(\frac{L}{2}[\frac{\phi_1}{2} + \phi_2])}{\sin^2(\frac{1}{2}[\frac{\phi_1}{2} + \phi_2])} \frac{\sin^2(\frac{L}{2}[-\frac{\phi_1}{2} + \phi_2])}{\sin^2(\frac{1}{2}[-\frac{\phi_1}{2} + \phi_2])} \\
&\quad \left(\left(\frac{\phi_1}{2} + \phi_2 \right)^4 + \left(\frac{\phi_1}{2} - \phi_2 \right)^4 \right)
\end{aligned}$$

where we used the Cauchy-Schwarz inequality in the second line and equation (1.11) in the last line. Changing coordinates to

$$\alpha_1 = \phi_1 + \frac{\phi_2}{2} \quad \alpha_2 = \frac{\phi_1}{2} - \phi_2$$

and realizing that $|\alpha_1|, |\alpha_2| < 3\epsilon/2$ for $(\phi_1, \phi_2) \in D_\epsilon$, we see that

$$\begin{aligned}
\langle g_1(t) | g_1(t) \rangle &\leq \frac{16t^2}{L^2} \int_{-3\epsilon/2}^{3\epsilon/2} \frac{d\alpha_1}{2\pi} \int_{-3\epsilon/2}^{3\epsilon/2} \frac{d\alpha_2}{2\pi} \frac{\sin^2(\frac{1}{2}L\alpha_1)}{\sin^2(\frac{1}{2}\alpha_1)} \frac{\sin^2(\frac{1}{2}L\alpha_2)}{\sin^2(\frac{1}{2}\alpha_2)} (\alpha_1^4 + \alpha_2^4) \\
&= \frac{32t^2}{L^2} \int_{-3\epsilon/2}^{3\epsilon/2} \frac{d\alpha_1}{2\pi} \int_{-3\epsilon/2}^{3\epsilon/2} \frac{d\alpha_2}{2\pi} \frac{\sin^2(\frac{1}{2}L\alpha_1)}{\sin^2(\frac{1}{2}\alpha_1)} \frac{\sin^2(\frac{1}{2}L\alpha_2)}{\sin^2(\frac{1}{2}\alpha_2)} \alpha_1^4 \\
&\leq \frac{32t^2}{L} \int_{-3\epsilon/2}^{3\epsilon/2} \frac{d\alpha_1}{2\pi} \frac{\sin^2(\frac{1}{2}L\alpha_1)}{\sin^2(\frac{1}{2}\alpha_1)} \alpha_1^4 \\
&\leq \frac{32t^2}{L} \int_{-3\epsilon/2}^{3\epsilon/2} \frac{d\alpha_1}{2\pi} \frac{\pi^2}{\alpha_1^2} \alpha_1^4 \\
&= \frac{36\pi t^2 \epsilon^3}{L},
\end{aligned}$$

with a similar bound on $\langle g_2(t) | g_2(t) \rangle$.

Finally,

$$\langle h(t) | h(t) \rangle \leq \frac{1}{2} \iint_{D_\epsilon} \frac{d\phi_1 d\phi_2}{4\pi^2} |A(\phi_1, \phi_2)|^2 \left| e^{i\theta \pm (\frac{\pi}{4} + \phi_1, \frac{3\pi}{8} + \phi_2)} - e^{i\theta} \right|^2.$$

Recall that $e^{i\theta \pm (p_1, p_2)} = T(p_1, p_2) \pm R(p_1, p_2)$ is obtained by solving for the effective single-particle S-matrix for the Hamiltonian (??). For p_1 near $\pi/4$ we divide this Hamiltonian by $2 \cos(p_1/2)$ to put it in the form considered in [?], where the potential term is now $\mathcal{V}(|r|)/(2 \cos(p_1/2))$. The entries $T(p_1, p_2)$ and $R(p_1, p_2)$ of this S-matrix are bounded rational functions of $z = e^{ip_2}$ and $(2 \cos(p_1/2))^{-1}$ [?], so they are differentiable as a function of p_1 and

p_2 on some neighborhood U of $(\pi/4, 3\pi/8)$ (and have bounded partial derivatives on this neighborhood).

For ϵ small enough that $D_\epsilon \subset U$ we get, using the mean value theorem and the fact that $\theta = \theta_\pm(\pi/4, 3\pi/8)$,

$$\begin{aligned} \left| e^{i\theta_\pm(\frac{\pi}{4}+\phi_1, \frac{3\pi}{8}+\phi_2)} - e^{i\theta} \right| &\leq \sqrt{\phi_1^2 + \phi_2^2} \max_U |\vec{\nabla} e^{i\theta_\pm}| \quad \text{for } (\phi_1, \phi_2) \in D_\epsilon \\ &\leq \epsilon \Gamma \end{aligned}$$

for some constant Γ (independent of L). Therefore

$$\begin{aligned} \langle h(t) | h(t) \rangle &\leq \frac{1}{2} \iint_{D_\epsilon} \frac{d\phi_1 d\phi_2}{4\pi^2} |A(\phi_1, \phi_2)|^2 \epsilon^2 \Gamma^2 \\ &\leq \frac{1}{2} \Gamma^2 \epsilon^2. \end{aligned}$$

Putting these bounds together, we get

$$\begin{aligned} \|P_1|\psi_A(t)\rangle - P_1|\alpha(t)\rangle\| &\leq \| |e_1(t)\rangle \| + \| |e_2(t)\rangle \| + \| |f_1(t)\rangle \| + \| |f_2(t)\rangle \| \\ &\quad + \| |g_1(t)\rangle \| + \| |g_2(t)\rangle \| + \| |h(t)\rangle \| \\ &\leq 2 \left(\frac{80\pi^3}{L\epsilon} + 20\epsilon^2 \right)^{\frac{1}{2}} + 2 \left(\frac{2\pi}{L\epsilon} \right)^{\frac{1}{2}} + 2 \left(\frac{36\pi t^2 \epsilon^3}{L} \right)^{\frac{1}{2}} + \frac{1}{\sqrt{2}} \Gamma \epsilon. \end{aligned}$$

Letting $\epsilon = a/\sqrt{L}$ and $t \leq c_0 L$ we get

$$\|P_1|\psi_A(t)\rangle - P_1|\alpha(t)\rangle\| = \mathcal{O}(L^{-1/4}). \quad (1.8)$$

Since $P_2|\alpha(t)\rangle$ has support on at most $4CL$ basis states $|x, y\rangle$, and since $|\langle x, y | P_2|\alpha(t)\rangle|^2 = \mathcal{O}(L^{-2})$, we get

$$\|P_2|\alpha(t)\rangle\| = \mathcal{O}(L^{-1/2}). \quad (1.9)$$

We now use the bounds (1.8) and (1.9) and Lemma 2 to show that

$$\| |\psi_A(t)\rangle - |\alpha(t)\rangle \| = \mathcal{O}(L^{-1/8}). \quad (1.10)$$

First consider the case where the interaction range is $C = 0$ (as in the Bose-Hubbard model). In this case equation (1.10) follows directly from equation (1.8) and the facts that $\langle x, y | \alpha(t) \rangle = \pm \langle y, x | \alpha(t) \rangle$ and $\langle x, y | \psi_A(t) \rangle = \pm \langle y, x | \psi_A(t) \rangle$.

Now suppose $C \neq 0$. In this case

$$\begin{aligned} \|(1 - P_2) |\psi_A(t)\rangle\|^2 &= 2 \|P_1|\psi_A(t)\rangle\|^2 \\ &= 2 (\|P_1|\alpha(t)\rangle\| + \mathcal{O}(L^{-1/4}))^2 \\ &= 2 \left(\frac{1}{2} \|(1 - P_2)|\alpha(t)\rangle\|^2 + \mathcal{O}(L^{-1/4}) \right) \\ &= 1 + \mathcal{O}(L^{-1}) - \langle \alpha(t) | P_2 | \alpha(t) \rangle + \mathcal{O}(L^{-1/4}) \\ &= 1 + \mathcal{O}(L^{-1/4}) \end{aligned}$$

where in the next-to-last line we have used [Lemma 2](#). So

$$\begin{aligned}
\| |\psi_A(t)\rangle - |\alpha(t)\rangle \| &\leq 2 \| P_1 |\psi_A(t)\rangle - P_1 |\alpha(t)\rangle \| + \| P_2 |\alpha(t)\rangle \| + \| P_2 |\psi_A(t)\rangle \| \\
&= \mathcal{O}(L^{-1/4}) + \mathcal{O}(L^{-1/2}) + (1 - \|(1 - P_2) |\psi_A(t)\rangle\|)^{\frac{1}{2}} \\
&= \mathcal{O}(L^{-1/4}) + \mathcal{O}(L^{-1/2}) + \mathcal{O}(L^{-1/8}) \\
&= \mathcal{O}(L^{-1/8})
\end{aligned}$$

which completes the proof. □

□

1.2.1.1 Technical lemmas

In this section we prove three lemmas that are used in the proof of [Theorem 1](#).

Lemma 2. *Let $|\alpha(t)\rangle$ be defined as in [Theorem 1](#). Then*

$$\langle \alpha(t) | \alpha(t) \rangle = 1 + \mathcal{O}(L^{-1}).$$

Proof. Define

$$\Pi = \sum_{x \leq y} |x, y\rangle \langle x, y|.$$

Note that, since $\langle x, y | \alpha(t) \rangle = \pm \langle y, x | \alpha(t) \rangle$,

$$\begin{aligned}
\langle \alpha(t) | \alpha(t) \rangle &= 2 \langle \alpha(t) | \Pi | \alpha(t) \rangle - \sum_{x=-\infty}^{\infty} \langle \alpha(t) | x, x \rangle \langle x, x | \alpha(t) \rangle \\
&= 2 \langle \alpha(t) | \Pi | \alpha(t) \rangle + \mathcal{O}(L^{-1})
\end{aligned}$$

where the last line follows since $|\langle x, x | \alpha(t) \rangle|^2$ is nonzero for at most L values of x and $|\langle x, x | \alpha(t) \rangle|^2 = \mathcal{O}(L^{-2})$. We now show that

$$\langle \alpha(t) | \Pi | \alpha(t) \rangle = \frac{1}{2} + \mathcal{O}(L^{-1}).$$

Note that

$$\begin{aligned}
\langle \alpha(t) | \Pi | \alpha(t) \rangle &= \frac{1}{2L^2} \sum_{x \leq y} \left(F(x, y, t) + F(y, x, t) \right. \\
&\quad \left. \pm e^{i\theta} e^{\frac{3i\pi}{4}x} e^{-\frac{3i\pi}{4}y} F(x, y, t) F(y, x, t) \right. \\
&\quad \left. \pm e^{-i\theta} e^{-\frac{3i\pi}{4}x} e^{\frac{3i\pi}{4}y} F(x, y, t) F(y, x, t) \right).
\end{aligned}$$

Now $F(x, y, t) = 1$ if and only if $x \in \{-M - L + 2[t], \dots, -M - 1 + 2[t]\}$ and $y \in \{M + 1 - 2[t/\sqrt{2}], \dots, M + L - 2[t/\sqrt{2}]\}$. Similarly $F(y, x, t) = 1$ if and only if $x \in \{M + 1 - 2[t/\sqrt{2}], \dots, M + L - 2[t/\sqrt{2}]\}$ and $y \in \{-M - L + 2[t], \dots, -M - 1 + 2[t]\}$. So

$$\sum_{x \leq y} F(y, x, t) = \sum_{y \leq x} F(x, y, t)$$

and

$$\begin{aligned} \frac{1}{2L^2} \sum_{x \leq y} [F(x, y, t) + F(y, x, t)] &= \frac{1}{2L^2} \left(\sum_{x=-\infty}^{\infty} \sum_{y=-\infty}^{\infty} F(x, y, t) - \sum_{x=-\infty}^{\infty} F(x, x, t) \right) \\ &= \frac{1}{2} + \mathcal{O}(L^{-1}). \end{aligned}$$

We now establish the bound

$$\left| \frac{1}{2L^2} \sum_{x \leq y} e^{\frac{3i\pi}{4}x} e^{-\frac{3i\pi}{4}y} F(x, y, t) F(y, x, t) \right| = \mathcal{O}(L^{-1})$$

to complete the proof. To get this bound, note that both $F(x, y, t) = 1$ and $F(y, x, t) = 1$ if and only if

$$\begin{aligned} &x, y \in \{-M - L + 2 \lfloor t \rfloor, \dots, -M - 1 + 2 \lfloor t \rfloor\} \\ \text{and } &x, y \in \left\{ M + 1 - 2 \left\lfloor \frac{t}{\sqrt{2}} \right\rfloor, \dots, M + L - 2 \left\lfloor \frac{t}{\sqrt{2}} \right\rfloor \right\}. \end{aligned}$$

Letting

$$B = \{-M - L + 2 \lfloor t \rfloor, \dots, -M - 1 + 2 \lfloor t \rfloor\} \cap \left\{ M + 1 - 2 \left\lfloor \frac{t}{\sqrt{2}} \right\rfloor, \dots, M + L - 2 \left\lfloor \frac{t}{\sqrt{2}} \right\rfloor \right\},$$

we have

$$B = \{j, j+1, \dots, j+l\}$$

for some $j, l \in \mathbb{Z}$ with $l < L$. So

$$\begin{aligned} \frac{1}{2L^2} \left| \sum_{x \leq y} e^{\frac{3i\pi}{4}x} e^{-\frac{3i\pi}{4}y} F(x, y, t) F(y, x, t) \right| &= \frac{1}{2L^2} \left| \sum_{x, y \in B, x \leq y} e^{\frac{3i\pi}{4}x} e^{-\frac{3i\pi}{4}y} \right| \\ &= \frac{1}{2L^2} \left| \sum_{y=j}^{j+l} \sum_{x=j}^y e^{\frac{3i\pi}{4}x} e^{-\frac{3i\pi}{4}y} \right| \\ &= \frac{1}{2L^2} \left| \sum_{y=j}^{j+l} e^{-\frac{3i\pi}{4}y} e^{3i\frac{\pi}{4}j} \frac{e^{3i\frac{\pi}{4}(y+1-j)} - 1}{e^{3i\frac{\pi}{4}} - 1} \right| \\ &\leq \frac{(l+1)}{2L^2} \frac{2}{|e^{3i\frac{\pi}{4}} - 1|} \\ &= \mathcal{O}(L^{-1}) \end{aligned}$$

since $l < L$. □

Lemma 3. Let $k \in (-\pi, 0) \cup (0, \pi)$ and $0 < \epsilon < \min \{\pi - |k|, |k|\}$. Let

$$\begin{aligned} D_\epsilon &= [-\epsilon, \epsilon] \times [-\epsilon, \epsilon] \\ D_\pi &= [-\pi, \pi] \times [-\pi, \pi]. \end{aligned}$$

Then

$$\begin{aligned} \iint_{D_\pi} \frac{d\phi_1 d\phi_2}{4\pi^2} |A(\phi_1, \phi_2)|^2 &= 1 \\ \iint_{D_\pi \setminus D_\epsilon} \frac{d\phi_1 d\phi_2}{4\pi^2} |A(\phi_1, \phi_2)|^2 &\leq \frac{4\pi}{L\epsilon} \\ \iint_{D_\epsilon} \frac{d\phi_1 d\phi_2}{4\pi^2} |B(\phi_1, \phi_2, k)|^2 &\leq \frac{4\pi^2}{L^2\epsilon^2}. \end{aligned}$$

where $A(\phi_1, \phi_2)$ and $B(\phi_1, \phi_2, k)$ are given by equation (1.6).

Proof. Using equation (1.6) we get

$$|A(\phi_1, \phi_2)|^2 = \frac{1}{L^2} \sum_{x, \tilde{x} = -(M+L)}^{-(M+1)} \sum_{y, \tilde{y} = M+1}^{M+L} e^{i\frac{\phi_1}{2}(x+y-(\tilde{x}+\tilde{y}))} e^{i\phi_2(x-y-(\tilde{x}-\tilde{y}))}.$$

Now

$$\int_{-\pi}^{\pi} \frac{d\phi_2}{2\pi} e^{i\phi_2(x-y-(\tilde{x}-\tilde{y}))} = \delta_{x-y, \tilde{x}-\tilde{y}},$$

so (suppressing the limits of summation for readability)

$$\begin{aligned} \iint_{D_\pi} \frac{d\phi_1 d\phi_2}{4\pi^2} |A(\phi_1, \phi_2)|^2 &= \frac{1}{L^2} \int_{-\pi}^{\pi} \frac{d\phi_1}{2\pi} \sum_{x, \tilde{x}} \sum_{y, \tilde{y}} e^{i\phi_1(y-\tilde{y})} \delta_{x-y, \tilde{x}-\tilde{y}} \\ &= \frac{1}{L^2} \sum_{x, \tilde{x}} \sum_{y, \tilde{y}} \delta_{y, \tilde{y}} \delta_{x-y, \tilde{x}-\tilde{y}} \\ &= 1 \end{aligned}$$

which proves the first part.

By performing the sums in equation (1.6) we get

$$|A(\phi_1, \phi_2)|^2 = \frac{1}{L^2} \frac{\sin^2(\frac{1}{2}L[\frac{\phi_1}{2} + \phi_2])}{\sin^2(\frac{1}{2}[\frac{\phi_1}{2} + \phi_2])} \frac{\sin^2(\frac{1}{2}L[\frac{\phi_1}{2} - \phi_2])}{\sin^2(\frac{1}{2}[\frac{\phi_1}{2} - \phi_2])}. \quad (1.11)$$

Letting $\alpha_1 = \phi_1/2 + \phi_2$ and $\alpha_2 = \phi_1/2 - \phi_2$, we see that $|\alpha_1| \leq 3\pi/2$, $|\alpha_2| \leq 3\pi/2$, and $\alpha_1^2 + \alpha_2^2 \geq 5\epsilon^2/2$ whenever $(\phi_1, \phi_2) \in D_\pi \setminus D_\epsilon$. Defining $D_{3\pi/2} = [-3\pi/2, 3\pi/2]^2$ we get

$(\alpha_1, \alpha_2) \in D_{3\pi/2} \setminus D_\epsilon$ whenever $(\phi_1, \phi_2) \in D_\pi \setminus D_\epsilon$. Hence

$$\begin{aligned}
\iint_{D_\pi \setminus D_\epsilon} \frac{d\phi_1 d\phi_2}{4\pi^2} |A(\phi_1, \phi_2)|^2 &\leq \frac{1}{L^2} \iint_{D_{3\pi/2} \setminus D_\epsilon} \frac{d\alpha_1 d\alpha_2}{4\pi^2} \frac{\sin^2(\frac{1}{2}L\alpha_1)}{\sin^2(\frac{1}{2}\alpha_1)} \frac{\sin^2(\frac{1}{2}L\alpha_2)}{\sin^2(\frac{1}{2}\alpha_2)} \\
&\leq \frac{4}{L} \left(\frac{1}{L} \int_{-\frac{3\pi}{2}}^{\frac{3\pi}{2}} \frac{d\alpha_1}{2\pi} \frac{\sin^2(\frac{1}{2}L\alpha_1)}{\sin^2(\frac{1}{2}\alpha_1)} \right) \left(\int_{\epsilon}^{3\pi/2} \frac{d\alpha_2}{2\pi} \frac{\sin^2(\frac{1}{2}L\alpha_2)}{\sin^2(\frac{1}{2}\alpha_2)} \right) \\
&\leq \frac{4}{L} \left(\int_{-2\pi}^{2\pi} \frac{d\alpha_1}{2\pi} \frac{1}{L} \frac{\sin^2(\frac{1}{2}L\alpha_1)}{\sin^2(\frac{1}{2}\alpha_1)} \right) \left(\int_{\epsilon}^{\frac{3\pi}{2}} \frac{d\alpha_2}{2\pi} \frac{1}{\sin^2(\frac{1}{2}\alpha_2)} \right) \\
&= \frac{8}{L} \left(\int_{\epsilon}^{\pi} \frac{d\alpha_2}{2\pi} \frac{1}{\sin^2(\frac{1}{2}\alpha_2)} + \int_{\pi}^{\frac{3\pi}{2}} \frac{d\alpha_2}{2\pi} \frac{1}{\sin^2(\frac{1}{2}\alpha_2)} \right) \\
&\leq \frac{8}{L} \left(\int_{\epsilon}^{\pi} \frac{d\alpha_2}{2\pi} \frac{\pi^2}{\alpha_2^2} + 2 \int_{\pi}^{\frac{3\pi}{2}} \frac{d\alpha_2}{2\pi} \right) \\
&= \frac{4\pi}{L\epsilon}
\end{aligned}$$

which proves the second inequality (in the next-to-last line we have used the fact that $\sin(x/2) > x/\pi$ for $x \in (0, \pi)$ and $\sin^2(x/2) > 1/2$ for $x \in (\pi, 3\pi/2)$).

Now

$$\begin{aligned}
|B(\phi_1, \phi_2, k)|^2 &= |A(\phi_1, -\phi_2 - 2k)|^2 \\
&\leq \frac{1}{L^2} \frac{1}{\sin^2\left(\frac{1}{2}\left[\frac{\phi_1}{2} + \phi_2 + 2k\right]\right)} \frac{1}{\sin^2\left(\frac{1}{2}\left[-\frac{\phi_1}{2} + \phi_2 + 2k\right]\right)}.
\end{aligned}$$

If $(\phi_1, \phi_2) \in D_\epsilon$ then $|k| - 3\epsilon/4 \leq |\pm\phi_1/4 + \phi_2/2 + k| \leq |k| + 3\epsilon/4$. Noting that ϵ is chosen such that $0 < \epsilon < \min\{\pi - |k|, |k|\}$, we get

$$\frac{\epsilon}{4} \leq \left| \pm \frac{\phi_1}{4} + \frac{\phi_2}{2} + k \right| \leq \pi - \frac{\epsilon}{4}$$

so

$$\begin{aligned}
|B(\phi_1, \phi_2, k)|^2 &\leq \frac{1}{L^2} \frac{1}{\sin^4(\frac{\epsilon}{4})} \\
&\leq \frac{16\pi^4}{L^2\epsilon^4}
\end{aligned}$$

and

$$\begin{aligned}
\iint_{D_\epsilon} \frac{d\phi_1 d\phi_2}{4\pi^2} |B(\phi_1, \phi_2, k)|^2 &\leq \frac{1}{4\pi^2} (2\epsilon)^2 \left(\frac{16\pi^4}{L^2\epsilon^4} \right) \\
&= \frac{16\pi^2}{L^2\epsilon^2}.
\end{aligned}$$

□

Lemma 4. Let $a_{xy}(t)$ be as in Theorem 1. For $x \leq y$,

$$\begin{aligned} a_{xy}(t) = & \frac{1}{\sqrt{2}} e^{-it\sqrt{2}} \left[e^{-i\pi x/2} e^{i\pi y/4} \left(\iint_{D_\pi} \frac{d\phi_1 d\phi_2}{4\pi^2} A(\phi_1, \phi_2) \right. \right. \\ & \left. \left. e^{-i\phi_1 \left(-\lfloor t \rfloor + \lfloor \frac{t}{\sqrt{2}} \rfloor + \frac{x+y}{2} \right)} e^{-2i\phi_2 \left(-\lfloor t \rfloor - \lfloor \frac{t}{\sqrt{2}} \rfloor + \frac{y-x}{2} \right)} \right) \right. \\ & \left. \pm e^{i\theta} e^{i\pi x/4} e^{-i\pi y/2} \left(\iint_{D_\pi} \frac{d\phi_1 d\phi_2}{4\pi^2} A(\phi_1, \phi_2) \right. \right. \\ & \left. \left. e^{-i\phi_1 \left(-\lfloor t \rfloor + \lfloor \frac{t}{\sqrt{2}} \rfloor + \frac{x+y}{2} \right)} e^{-2i\phi_2 \left(-\lfloor t \rfloor - \lfloor \frac{t}{\sqrt{2}} \rfloor + \frac{y-x}{2} \right)} \right) \right]. \end{aligned}$$

Proof. The lemma follows from (1.2) and the fact that, for any two numbers γ_1, γ_2 such that $\gamma_1 + \gamma_2, \gamma_1 - \gamma_2 \in \mathbb{Z}$,

$$\iint_{D_\pi} \frac{d\phi_1 d\phi_2}{4\pi^2} A(\phi_1, \phi_2) e^{i\gamma_1 \phi_1 + 2i\gamma_2 \phi_2} = \begin{cases} \frac{1}{L} & \text{if } (-\gamma_1 - \gamma_2, -\gamma_1 + \gamma_2) \in S \\ 0 & \text{otherwise} \end{cases}$$

where $S = \{-M-L, \dots, -M-1\} \times \{M+1, \dots, M+L\}$. To establish this formula, observe that

$$\begin{aligned} \iint_{D_\pi} \frac{d\phi_1 d\phi_2}{4\pi^2} A(\phi_1, \phi_2) e^{i\gamma_1 \phi_1 + 2i\gamma_2 \phi_2} &= \frac{1}{L} \sum_{x=-M-L}^{-M-1} \sum_{y=M+1}^{M+L} \iint_{D_\pi} \frac{d\phi_1 d\phi_2}{4\pi^2} e^{i\phi_1 \left(\gamma_1 + \frac{x+y}{2} \right)} e^{i\phi_2 (x-y+2\gamma_2)} \\ &= \frac{1}{L} \sum_{x=-M-L}^{-M-1} \sum_{y=M+1}^{M+L} \int_{-\pi}^{\pi} \frac{d\phi_1}{2\pi} e^{i\phi_1 \left(\gamma_1 + \frac{x-y}{2} \right)} \delta_{y, -x-2\gamma_2}. \end{aligned}$$

Here we have performed the integral over ϕ_2 using the fact that $2\gamma_2$ is an integer. We then have

$$\begin{aligned} \iint_{D_\pi} \frac{d\phi_1 d\phi_2}{4\pi^2} A(\phi_1, \phi_2) e^{i\gamma_1 \phi_1 + 2i\gamma_2 \phi_2} &= \frac{1}{L} \sum_{x=-M-L}^{-M-1} \sum_{y=M+1}^{M+L} \int_{-\pi}^{\pi} \frac{d\phi_1}{2\pi} e^{i\phi_1 (\gamma_1 + x + \gamma_2)} \delta_{y, -x-2\gamma_1} \\ &= \frac{1}{L} \sum_{x=-M-L}^{-M-1} \sum_{y=M+1}^{M+L} \delta_{x, -\gamma_1 - \gamma_2} \delta_{y, \gamma_2 - \gamma_1} \end{aligned}$$

as claimed. □

1.2.2 Construction of $C\theta$ -gate

1.3 Impossibility of some momentum switches

1.4 Universal Computation

1.4.1 Two-qubit blocks

1.4.2 Combining blocks

1.5 Improvements and Modifications

What about long-range interactions, but where the interactions die off? Additionally, what about error correction?