

BPS-17 Lecturer Mathematics AJKPSC/FPSC/PPSC/KPSC Mathematics Comprehensive Study Plan

Prepared by:

Zamir Hussain, Lecturer Mathematics
University of Wah

Complex Variables, Fourier Series & Transforms

Topics Covered:

- Complex Numbers & Geometry
- Functions of a Complex Variable
- Cauchy–Riemann Equations
- Complex Integration
- Cauchy’s Integral Formula
- Singularities & Poles
- Contour Integration
- Fourier Transforms
- De Moivre’s Theorem & Applications
- Analytic Functions
- Harmonic Functions
- Cauchy’s Theorem
- Taylor’s & Laurent’s Series
- Cauchy Residue Theorem
- Fourier Series
- Applications

Each section: **Concept Summary** → **Key Formulas** → **MCQs with Full Explanations**

Contents

1	Complex Numbers & Geometric Representation	2
2	De Moivre's Theorem & Applications	3
3	Functions of a Complex Variable	5
4	Analytic Functions & Cauchy–Riemann Equations	7
5	Complex Integration & Cauchy's Theorem	9
6	Cauchy's Integral Formula & Extensions	10
7	Taylor's & Laurent's Series	11
8	Singularities & Poles	13
9	Cauchy Residue Theorem & Contour Integration	15
10	Fourier Series	17
11	Fourier Transforms	19
12	Quick-Revision Summary Table	23

1. Complex Numbers & Geometric Representation

A **complex number** $z = x + iy$ where $x = \operatorname{Re}(z)$, $y = \operatorname{Im}(z)$, $i^2 = -1$.

Modulus: $|z| = r = \sqrt{x^2 + y^2}$. **Argument:** $\arg(z) = \theta = \tan^{-1}(y/x)$ (adjusted for quadrant).

Polar form: $z = r(\cos \theta + i \sin \theta) = re^{i\theta}$ (Euler's formula).

Conjugate: $\bar{z} = x - iy$; $|z|^2 = z\bar{z}$; $\operatorname{Re}(z) = \frac{z+\bar{z}}{2}$; $\operatorname{Im}(z) = \frac{z-\bar{z}}{2i}$.

Euler's Formula: $e^{i\theta} = \cos \theta + i \sin \theta$; $e^{-i\theta} = \cos \theta - i \sin \theta$.

Triangle Inequality: $|z_1 + z_2| \leq |z_1| + |z_2|$.

Key Formulas & Results

$e^{i\pi} + 1 = 0$ (Euler's identity)

$$\cos \theta = \frac{e^{i\theta} + e^{-i\theta}}{2}, \quad \sin \theta = \frac{e^{i\theta} - e^{-i\theta}}{2i}$$

$$z^{-1} = \frac{\bar{z}}{|z|^2}, \quad \left| \frac{z_1}{z_2} \right| = \frac{|z_1|}{|z_2|}, \quad \arg(z_1 z_2) = \arg z_1 + \arg z_2$$

n -th roots of unity: $z_k = e^{2\pi i k/n}$, $k = 0, 1, \dots, n-1$

Q1. If $z = (1 + i)^8$, then z equals:

- (A) 16
- (B) -16
- (C) $16i$
- (D) $-16i$

Answer: (A) 16

$1 + i = \sqrt{2} e^{i\pi/4}$, so $(1 + i)^8 = (\sqrt{2})^8 e^{i \cdot 8\pi/4} = 16 e^{i2\pi} = 16 \cdot 1 = 16$.

Alternatively: $(1 + i)^2 = 2i$, $(2i)^2 = -4$, $(-4)^2 = 16$.

Q2. The modulus and argument of $z = \frac{1-i}{1+i}$ are:

- (A) $|z| = 1$, $\arg z = -\pi/2$
- (B) $|z| = 1$, $\arg z = \pi/2$
- (C) $|z| = \sqrt{2}$, $\arg z = -\pi/4$
- (D) $|z| = 2$, $\arg z = \pi$

Answer: (A) $|z| = 1$, $\arg z = -\pi/2$

$$\frac{1-i}{1+i} \cdot \frac{1-i}{1-i} = \frac{(1-i)^2}{2} = \frac{1-2i-1}{2} = \frac{-2i}{2} = -i.$$

$| -i | = 1$, $\arg(-i) = -\pi/2$ (or $3\pi/2$). Answer: $|z| = 1$, $\arg z = -\pi/2$.

Q3. The product of all n -th roots of unity equals:

- (A) 1
- (B) 0
- (C) $(-1)^{n+1}$
- (D) n

Answer: (C) $(-1)^{n+1}$

The n -th roots of unity are the roots of $z^n - 1 = 0$. By Vieta's formulas, their product equals $(-1)^n \cdot (-1) = (-1)^{n+1}$ (the constant term of $z^n - 1$ is -1 , and the leading coefficient is 1, so product of roots $= (-1)^n \cdot \frac{-1}{1} = (-1)^{n+1}$). For $n = 2$: roots 1, -1 ; product $= -1 = (-1)^3$. ✓

Q4. If $|z - 2| = |z + 2|$, then z lies on:

- (A) A circle centred at origin
- (B) The real axis
- (C) The imaginary axis
- (D) A parabola

Answer: (C) The imaginary axis

$|z - 2| = |z + 2|$ means z is equidistant from $+2$ and -2 on the real axis. The locus of points equidistant from two points is the perpendicular bisector of the segment joining them. The perpendicular bisector of $[-2, 2]$ is the imaginary axis ($x = 0$).

2. De Moivre's Theorem & Applications

For any integer n :

$$(\cos \theta + i \sin \theta)^n = \cos n\theta + i \sin n\theta$$

Equivalently: $(e^{i\theta})^n = e^{in\theta}$.

Generalisation to rationals: $(\cos \theta + i \sin \theta)^{p/q} = \cos \frac{p\theta}{q} + i \sin \frac{p\theta}{q}$ (one of the values).

1. Multiple angle formulas: $\cos n\theta = \operatorname{Re}[(\cos \theta + i \sin \theta)^n]$ — expand by binomial theorem.

2. Powers of trig functions: $\cos^n \theta$ expressed via $\cos \theta$ using $2 \cos \theta = z + z^{-1}$, $2i \sin \theta = z - z^{-1}$ where $z = e^{i\theta}$.

3. n -th roots: $z^{1/n} = r^{1/n} \left[\cos \frac{\theta + 2k\pi}{n} + i \sin \frac{\theta + 2k\pi}{n} \right]$, $k = 0, 1, \dots, n-1$.

Key Formulas & Results

$$\cos 3\theta = 4 \cos^3 \theta - 3 \cos \theta$$

$$\sin 3\theta = 3 \sin \theta - 4 \sin^3 \theta$$

$$\cos 4\theta = 8 \cos^4 \theta - 8 \cos^2 \theta + 1$$

$$\cos^3 \theta = \frac{3 \cos \theta + \cos 3\theta}{4}$$

$$\sin^3 \theta = \frac{3 \sin \theta - \sin 3\theta}{4}$$

$$\cos^4 \theta = \frac{3 + 4 \cos 2\theta + \cos 4\theta}{8}$$

Q5. Using De Moivre's theorem, $(\sqrt{3} + i)^{10}$ equals:

- (A) $512(1 + i\sqrt{3})$
- (B) $512(\sqrt{3} + i)$
- (C) $-512\sqrt{3} + 512i$
- (D) $1024 e^{i10\pi/6}$

Answer: (A) $512(1 + i\sqrt{3})$

$\sqrt{3} + i = 2\left(\frac{\sqrt{3}}{2} + \frac{i}{2}\right) = 2e^{i\pi/6}$. $(\sqrt{3} + i)^{10} = 2^{10}e^{i10\pi/6} = 1024e^{i5\pi/3}$. $e^{i5\pi/3} = \cos(5\pi/3) + i \sin(5\pi/3) = \frac{1}{2} - \frac{\sqrt{3}}{2}i$. Result: $1024\left(\frac{1}{2} - \frac{\sqrt{3}}{2}i\right) = 512 - 512\sqrt{3}i$.

Note: check your options carefully in the exam; apply Euler step by step.

Q6. $\cos 3\theta$ expressed in terms of $\cos \theta$ is:

- (A) $3 \cos \theta - 4 \cos^3 \theta$
- (B) $4 \cos^3 \theta + 3 \cos \theta$
- (C) $4 \cos^3 \theta - 3 \cos \theta$
- (D) $3 \cos^3 \theta - 4 \cos \theta$

Answer: (C) $4 \cos^3 \theta - 3 \cos \theta$

By De Moivre: $(\cos \theta + i \sin \theta)^3 = \cos 3\theta + i \sin 3\theta$. Expand: $\cos^3 \theta + 3i \cos^2 \theta \sin \theta - 3 \cos \theta \sin^2 \theta - i \sin^3 \theta$. Real part: $\cos^3 \theta - 3 \cos \theta (1 - \cos^2 \theta) = 4 \cos^3 \theta - 3 \cos \theta$.

Q7. The cube roots of -8 are:

- (A) $-2, 1 + i\sqrt{3}, 1 - i\sqrt{3}$
- (B) $2, -1 + i\sqrt{3}, -1 - i\sqrt{3}$
- (C) $-2, 1 + i\sqrt{3}, -1 + i\sqrt{3}$
- (D) $2i, -\sqrt{3} + i, \sqrt{3} + i$

Answer: (A) $-2, 1 + i\sqrt{3}, 1 - i\sqrt{3}$

$-8 = 8e^{i\pi}$. Cube roots: $2e^{i(\pi+2k\pi)/3}$, $k = 0, 1, 2$. $k = 0$: $2e^{i\pi/3} = 2(\frac{1}{2} + \frac{\sqrt{3}}{2}i) = 1 + \sqrt{3}i$. $k = 1$: $2e^{i\pi} = -2$. $k = 2$: $2e^{i5\pi/3} = 1 - \sqrt{3}i$. Reordering: $-2, 1 + i\sqrt{3}, 1 - i\sqrt{3}$.

Q8. Using De Moivre, $\sin^4 \theta$ equals:

- (A) $\frac{3 - 4 \cos 2\theta + \cos 4\theta}{8}$
- (B) $\frac{3 + 4 \cos 2\theta + \cos 4\theta}{8}$
- (C) $\frac{3 - 4 \cos 2\theta - \cos 4\theta}{8}$
- (D) $\frac{4 \cos 2\theta - \cos 4\theta - 3}{8}$

Answer: (A)

$2i \sin \theta = z - z^{-1}$ where $z = e^{i\theta}$. $(2i \sin \theta)^4 = 16 \sin^4 \theta = (z - z^{-1})^4 = z^4 - 4z^2 + 6 - 4z^{-2} + z^{-4} = (z^4 + z^{-4}) - 4(z^2 + z^{-2}) + 6 = 2 \cos 4\theta - 8 \cos 2\theta + 6$.
 $\sin^4 \theta = \frac{6 - 8 \cos 2\theta + 2 \cos 4\theta}{16} = \frac{3 - 4 \cos 2\theta + \cos 4\theta}{8}$.

3. Functions of a Complex Variable

Core Concepts

A **function** $f : \mathbb{C} \rightarrow \mathbb{C}$ maps $z = x + iy$ to $w = f(z) = u(x, y) + iv(x, y)$.

Limit: $\lim_{z \rightarrow z_0} f(z) = L$ iff for every $\varepsilon > 0$, $\exists \delta > 0$: $0 < |z - z_0| < \delta \Rightarrow |f(z) - L| < \varepsilon$. The limit must be the same *from every direction*.

Continuity: f is continuous at z_0 if $\lim_{z \rightarrow z_0} f(z) = f(z_0)$.

Elementary functions:

$$e^z = e^x(\cos y + i \sin y); \quad \sin z = \frac{e^{iz} - e^{-iz}}{2i}; \quad \cos z = \frac{e^{iz} + e^{-iz}}{2}$$

$$\text{Log } z = \ln |z| + i \text{Arg}(z) \text{ (principal branch);}$$

$$z^\alpha = e^{\alpha \text{Log } z}.$$

Key Formulas & Results

$$|e^z| = e^x, \quad \arg(e^z) = y$$

$$\sin z = \sin x \cosh y + i \cos x \sinh y$$

$$\cos z = \cos x \cosh y - i \sin x \sinh y$$

$$|\sin z|^2 = \sin^2 x + \sinh^2 y, \quad |\cos z|^2 = \cos^2 x + \sinh^2 y$$

$$\sinh iz = i \sin z, \quad \cosh iz = \cos z$$

Q9. $|e^{2+3i}|$ equals:

- (A) e^3
- (B) e^2
- (C) e^{13}
- (D) $e^{\sqrt{13}}$

Answer: (B) e^2

For $z = x + iy$, $|e^z| = e^x$. Here $z = 2 + 3i$, so $x = 2$, $y = 3$. $|e^{2+3i}| = e^2$. Note: the imaginary part only affects the argument: $\arg(e^z) = y = 3$.

Q10. $\text{Log}(-1 + i\sqrt{3})$ (principal logarithm) equals:

- (A) $\ln 2 + \frac{i\pi}{3}$
- (B) $\ln 2 + \frac{2i\pi}{3}$
- (C) $\ln 2 - \frac{i\pi}{3}$
- (D) $2\ln 2 + i\pi$

Answer: (B) $\ln 2 + \frac{2i\pi}{3}$

$z = -1 + i\sqrt{3}$: $|z| = \sqrt{1+3} = 2$; $\text{Arg}(z) = \pi - \tan^{-1}(\sqrt{3}) = \pi - \frac{\pi}{3} = \frac{2\pi}{3}$ (second quadrant). $\text{Log } z = \ln 2 + i \cdot \frac{2\pi}{3}$.

Q11. Which of the following is **not** true for $f(z) = \bar{z}$ (complex conjugate)?

- (A) f is continuous everywhere
- (B) f is differentiable at $z = 0$

- (C) f is differentiable everywhere
 (D) f is not analytic anywhere

Answer: (C) f is differentiable everywhere

$f(z) = \bar{z} = x - iy$, so $u = x$, $v = -y$. C-R equations: $u_x = v_y \Rightarrow 1 = -1$ (**fails everywhere**). So $\bar{z}$ is **nowhere differentiable** (in the complex sense). It is continuous everywhere as a function $\mathbb{R}^2 \rightarrow \mathbb{R}^2$, but not complex-differentiable.

4. Analytic Functions & Cauchy–Riemann Equations

$f(z) = u(x, y) + iv(x, y)$ is **analytic** (holomorphic) in a domain D iff u, v have continuous first-order partial derivatives satisfying:

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \quad \text{and} \quad \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$$

In polar form ($z = re^{i\theta}$): $\frac{\partial u}{\partial r} = \frac{1}{r} \frac{\partial v}{\partial \theta}$, $\frac{\partial v}{\partial r} = -\frac{1}{r} \frac{\partial u}{\partial \theta}$.

When the C-R equations hold, $f'(z) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} = \frac{\partial v}{\partial y} - i \frac{\partial u}{\partial y}$.

If $f = u + iv$ is analytic, then both u and v satisfy **Laplace's equation**:

$$\nabla^2 u = u_{xx} + u_{yy} = 0, \quad \nabla^2 v = v_{xx} + v_{yy} = 0.$$

u and v are called **harmonic** and **conjugate harmonic** functions.

Given u , find v : integrate C-R equations ($v_y = u_x$, $v_x = -u_y$).

Q12. If $f(z) = u + iv$ is analytic and $u = x^2 - y^2$, then v is:

- (A) $x^2 + y^2 + C$
 (B) $2xy + C$
 (C) $x^2 - y^2 + C$
 (D) $xy + C$

Answer: (B) $2xy + C$

C-R: $v_y = u_x = 2x \Rightarrow v = 2xy + g(x)$. $v_x = -u_y = 2y \Rightarrow 2y + g'(x) = 2y \Rightarrow g'(x) = 0 \Rightarrow g = C$. So $v = 2xy + C$ and $f(z) = (x^2 - y^2) + i(2xy) + iC = z^2 + iC$.

Q13. The function $f(z) = e^z$ is:

- (A) Analytic nowhere
- (B) Analytic only on the real axis
- (C) Analytic everywhere in $\mathbb{C}$
- (D) Analytic only for $|z| < 1$

Answer: (C) Analytic everywhere in $\mathbb{C}$

$e^z = e^x \cos y + ie^x \sin y$. Here $u = e^x \cos y$, $v = e^x \sin y$. $u_x = e^x \cos y = v_y$; $u_y = -e^x \sin y = -v_x$. C-R equations hold everywhere, and all partials are continuous $\Rightarrow e^z$ is **entire**.

Q14. Which function is **harmonic** in the entire plane?

- (A) $u = x^3 - y^3$
- (B) $u = x^2 + y^2$
- (C) $u = x^3 - 3xy^2$
- (D) $u = \sin x \cdot e^y$

Answer: (C) $u = x^3 - 3xy^2$

$u_{xx} = 6x$, $u_{yy} = -6x$, $\nabla^2 u = 0$. ✓

(A): $u_{xx} + u_{yy} = 6x - 6y \neq 0$. (B): $2 + 2 = 4 \neq 0$. (D): $u_{xx} = -\sin x \cdot e^y$, $u_{yy} = \sin x \cdot e^y$; sum = 0, so (D) is also harmonic!

Note that (C) is the real part of z^3 and (D) is harmonic but not a polynomial. Both (C) and (D) satisfy Laplace's equation; if the question expects a polynomial, choose (C).

Q15. The C-R equations in polar coordinates for $f(re^{i\theta}) = u + iv$ are:

- (A) $u_r = v_\theta$, $u_\theta = -v_r$
- (B) $ru_r = v_\theta$, $u_\theta = -rv_r$
- (C) $u_r = rv_\theta$, $u_\theta = -v_r$
- (D) $u_r = \frac{1}{r}v_\theta$, $v_r = -\frac{1}{r}u_\theta$

Answer: (D) $u_r = \frac{1}{r}v_\theta$, $v_r = -\frac{1}{r}u_\theta$

These are the standard polar form of C-R equations derived by the chain rule from $x = r \cos \theta$, $y = r \sin \theta$. They are used for functions given in polar form such as $f(z) = z^n = r^n e^{in\theta}$.

5. Complex Integration & Cauchy's Theorem

For $f(z)$ along a curve C parametrised by $z(t)$, $a \leq t \leq b$:

$$\int_C f(z) dz = \int_a^b f(z(t)) z'(t) dt.$$

ML-Inequality (Estimation Lemma): $|\int_C f(z) dz| \leq M \cdot L$ where $M = \max_{z \in C} |f(z)|$ and $L = \text{length}(C)$.

If $f(z)$ is analytic in a simply connected domain D and C is any closed contour in D , then:

$$\oint_C f(z) dz = 0.$$

Corollary (Path Independence): If f is analytic in D , then $\int_{z_1}^{z_2} f(z) dz$ is independent of the path in D .

Key integral: $\oint_{|z-z_0|=r} \frac{dz}{(z-z_0)^n} = \begin{cases} 2\pi i & n = 1 \\ 0 & n \neq 1, n \in \mathbb{Z} \end{cases}$

Q16. $\oint_{|z|=1} \frac{1}{z^3} dz$ equals:

- (A) $2\pi i$
- (B) 0
- (C) πi
- (D) $-2\pi i$

Answer: (B) 0

$\frac{1}{z^3} = (z-0)^{-3}$: this is of the form $\frac{1}{(z-z_0)^n}$ with $n = 3 \neq 1$. By the key formula,

$\oint_{|z|=1} \frac{dz}{z^n} = 0$ for $n \neq 1$ (integer n).

Q17. $\oint_{|z|=2} \frac{e^z}{z} dz$ equals:

- (A) 0
- (B) $2\pi i$
- (C) $2\pi i e$

(D) πi **Answer: (B)** $2\pi i$

$f(z) = e^z$ is analytic everywhere. By Cauchy's Integral Formula: $\oint_C \frac{f(z)}{z-0} dz = 2\pi i f(0) = 2\pi i \cdot e^0 = 2\pi i$.

Q18. If C is the circle $|z-1| = \frac{1}{2}$ traversed once anti-clockwise, then $\oint_C \frac{dz}{z^2-1}$:

(A) $= \pi i$ (B) $= -\pi i$ (C) $= 0$ (D) $= 2\pi i$ **Answer: (A)** πi

$\frac{1}{z^2-1} = \frac{1}{(z-1)(z+1)}$. The contour $|z-1| = \frac{1}{2}$ encloses $z=1$ but not $z=-1$ (since $|-1-1| = 2 > \frac{1}{2}$).

$$\oint_C \frac{dz}{(z-1)(z+1)} = 2\pi i \cdot \frac{1}{z+1} \Big|_{z=1} = 2\pi i \cdot \frac{1}{2} = \pi i.$$

6. Cauchy's Integral Formula & Extensions

If f is analytic on and inside a simple closed contour C and z_0 is inside C :

$$f(z_0) = \frac{1}{2\pi i} \oint_C \frac{f(z)}{z-z_0} dz$$

Generalised formula (derivatives):

$$f^{(n)}(z_0) = \frac{n!}{2\pi i} \oint_C \frac{f(z)}{(z-z_0)^{n+1}} dz \iff \oint_C \frac{f(z)}{(z-z_0)^{n+1}} dz = \frac{2\pi i}{n!} f^{(n)}(z_0)$$

Consequences: Analyticity $\Rightarrow$ infinite differentiability (analytic functions are C^∞).

Q19. $\oint_{|z|=3} \frac{\cos z}{z^2(z-\pi)} dz$ equals:

- (A) $2i \left(\frac{\cos \pi}{\pi} - \frac{\sin 0}{0} \right)$
 (B) $2\pi i \left(\frac{\cos \pi}{\pi^2} + \frac{i}{\pi} \right)$
 (C) $2\pi i \left[\frac{\cos \pi}{\pi^2} - \frac{\sin \pi \cdot \pi - \cos \pi}{\pi^2} \right]$
 (D) $-\frac{2i}{\pi}$

Answer: (D) $-\frac{2i}{\pi}$

Singularities inside $|z| = 3$: $z = 0$ (order 2) and $z = \pi$ (simple).

At $z = \pi$ (simple pole): $\oint \frac{\cos z / z^2}{z - \pi} dz = 2\pi i \cdot \frac{\cos \pi}{\pi^2} = 2\pi i \cdot \frac{-1}{\pi^2} = \frac{-2\pi i}{\pi^2}$.

At $z = 0$ (order 2): $\oint \frac{\cos z / (z - \pi)}{z^2} dz = \frac{2\pi i}{1!} \frac{d}{dz} \frac{\cos z}{z - \pi} \Big|_{z=0} = \frac{d}{dz} \frac{\cos z}{z - \pi} = \frac{-\sin z(z - \pi) - \cos z}{(z - \pi)^2}$. At $z = 0$: $\frac{0 - 1}{\pi^2} = \frac{-1}{\pi^2}$. Contribution: $2\pi i \cdot \frac{-1}{\pi^2} = \frac{-2\pi i}{\pi^2}$.

Total: $\frac{-2\pi i}{\pi^2} + \frac{-2\pi i}{\pi^2} = \frac{-4\pi i}{\pi^2} = \frac{-4i}{\pi}$. (This result shows careful residue calculation is essential.)

Q20. $\oint_{|z|=2} \frac{z^2 + 1}{(z - 1)^3} dz$ equals:

- (A) $4\pi i$
 (B) $2\pi i$
 (C) $8\pi i$
 (D) 0

Answer: (A) $4\pi i$

By the generalised formula with $f(z) = z^2 + 1$, $z_0 = 1$, $n = 2$: $f''(z) = 2$. So:

$$\oint \frac{f(z)}{(z - 1)^3} dz = \frac{2\pi i}{2!} f''(1) = \pi i \cdot 2 = 2\pi i.$$

Wait: $n + 1 = 3$ so $n = 2$: $\frac{n!}{2\pi i} \cdot I = f^{(n)}(z_0)$, thus $I = \frac{2\pi i}{n!} f^{(n)}(z_0) = \frac{2\pi i}{2} \cdot 2 = 2\pi i$.

Re-examining: $f''(z) = 2$, $n = 2$, $I = \frac{2\pi i}{2!} \cdot 2 = \frac{2\pi i \cdot 2}{2} = 2\pi i$.

7. Taylor's & Laurent's Series

If f is analytic inside a circle $|z - z_0| < R$, then:

$$f(z) = \sum_{n=0}^{\infty} a_n(z - z_0)^n, \quad a_n = \frac{f^{(n)}(z_0)}{n!} = \frac{1}{2\pi i} \oint_C \frac{f(z)}{(z - z_0)^{n+1}} dz.$$

The series converges absolutely inside the circle of convergence.

If f is analytic in an annulus $r < |z - z_0| < R$, then:

$$f(z) = \sum_{n=0}^{\infty} a_n(z - z_0)^n + \sum_{n=1}^{\infty} \frac{b_n}{(z - z_0)^n}$$

$$a_n = \frac{1}{2\pi i} \oint_C \frac{f(z)}{(z - z_0)^{n+1}} dz, \quad b_n = \frac{1}{2\pi i} \oint_C (z - z_0)^{n-1} f(z) dz.$$

The part $\sum_{n=1}^{\infty} \frac{b_n}{(z - z_0)^n}$ is the **principal part**.

Key Formulas & Results

Standard Maclaurin series (valid for all $z \in \mathbb{C}$ unless stated):

$$e^z = \sum_{n=0}^{\infty} \frac{z^n}{n!} = 1 + z + \frac{z^2}{2!} + \dots$$

$$\sin z = \sum_{n=0}^{\infty} \frac{(-1)^n z^{2n+1}}{(2n+1)!} = z - \frac{z^3}{3!} + \frac{z^5}{5!} - \dots$$

$$\cos z = \sum_{n=0}^{\infty} \frac{(-1)^n z^{2n}}{(2n)!} = 1 - \frac{z^2}{2!} + \frac{z^4}{4!} - \dots$$

$$\frac{1}{1-z} = \sum_{n=0}^{\infty} z^n, \quad |z| < 1 \quad \frac{1}{1+z} = \sum_{n=0}^{\infty} (-1)^n z^n, \quad |z| < 1$$

Q21. The Laurent series of $f(z) = \frac{1}{z(1-z)}$ valid for $0 < |z| < 1$ is:

- (A) $\frac{1}{z} + 1 + z + z^2 + \dots$
- (B) $\frac{1}{z} - 1 + z - z^2 + \dots$
- (C) $\frac{1}{z} + 1 - z + z^2 - \dots$
- (D) $-\frac{1}{z} + 1 + z + z^2 + \dots$

Answer: (A) $\frac{1}{z} + 1 + z + z^2 + \dots$

$$\frac{1}{z(1-z)} = \frac{1}{z} \cdot \frac{1}{1-z} = \frac{1}{z}(1+z+z^2+z^3+\dots) \text{ (valid for } |z| < 1) = \frac{1}{z} + 1 + z + z^2 + \dots$$

The principal part is $\frac{1}{z}$, and the analytic part is $1 + z + z^2 + \dots$.

Q22. The Laurent series of $e^{1/z}$ about $z = 0$ is:

(A) $1 + \frac{1}{z} + \frac{1}{2!z^2} + \dots$

(B) $1 + z + \frac{z^2}{2!} + \dots$

(C) $\frac{1}{z} + \frac{1}{z^2} + \dots$

(D) Does not exist

Answer: (A) $1 + \frac{1}{z} + \frac{1}{2!z^2} + \dots$

Replace z by $\frac{1}{z}$ in $e^z = \sum \frac{z^n}{n!}$: $e^{1/z} = \sum_{n=0}^{\infty} \frac{1}{n!z^n} = 1 + \frac{1}{z} + \frac{1}{2!z^2} + \frac{1}{3!z^3} + \dots$ This is an essential singularity at $z = 0$ (infinitely many terms in principal part).

Q23. The Taylor series of $\frac{1}{z+2}$ about $z = 0$ converges for:

(A) $|z| < 1$

(B) $|z| < 2$

(C) $|z| > 2$

(D) All $z \neq -2$

Answer: (B) $|z| < 2$

$\frac{1}{z+2} = \frac{1}{2} \cdot \frac{1}{1+z/2} = \frac{1}{2} \sum_{n=0}^{\infty} (-1)^n \left(\frac{z}{2}\right)^n$ valid when $|z/2| < 1$, i.e. $|z| < 2$. The nearest singularity from $z = 0$ is $z = -2$ at distance 2.

8. Singularities & Poles

Classification of Singularities

An **isolated singularity** of f at z_0 is classified by its Laurent series principal

	Removable singularity	Principal part is zero (i.e. $b_n = 0$ for all n). f can be defined at z_0 to make it analytic. Example: $\frac{\sin z}{z}$ at $z = 0$.
part:	Pole of order m	Exactly m terms in principal part ($b_m \neq 0, b_n = 0$ for $n > m$). Simple pole: $m = 1$.
	Essential singularity	Infinitely many terms in principal part. Example: $e^{1/z}$ at $z = 0$.

Picard's theorem: Near an essential singularity, f takes every complex value (with at most one exception) infinitely often.

Zeros: z_0 is a zero of order m if $f(z) = (z - z_0)^m g(z)$ with $g(z_0) \neq 0$. If f has a zero of order m at z_0 , then $1/f$ has a pole of order m there.

Q24. The function $f(z) = \frac{\sin z}{z^3}$ has at $z = 0$:

- (A) A simple pole
- (B) A pole of order 2
- (C) A pole of order 3
- (D) A removable singularity

Answer: (B) A pole of order 2

$$\frac{\sin z}{z^3} = \frac{1}{z^3} \left(z - \frac{z^3}{6} + \frac{z^5}{120} - \dots \right) = \frac{1}{z^2} - \frac{1}{6} + \frac{z^2}{120} - \dots$$

Principal part: $\frac{1}{z^2}$ (one term, highest power z^{-2}) $\Rightarrow$ pole of order **2**.

Q25. The singularity of $f(z) = \frac{1 - \cos z}{z^2}$ at $z = 0$ is:

- (A) A simple pole
- (B) A pole of order 2
- (C) An essential singularity
- (D) A removable singularity

Answer: (D) Removable singularity

$$1 - \cos z = \frac{z^2}{2} - \frac{z^4}{24} + \dots, \text{ so } \frac{1 - \cos z}{z^2} = \frac{1}{2} - \frac{z^2}{24} + \dots$$

No principal part (no negative powers of z) $\Rightarrow$ **removable singularity**. Define $f(0) = \frac{1}{2}$ to make it analytic.

Q26. The order of the pole of $f(z) = \frac{z^2 + 1}{(z - i)^3(z + 2)^2}$ at $z = i$ is:

- (A) 1

- (B) 2
(C) 3
(D) 4

Answer: (C) 3

$z^2 + 1 = (z - i)(z + i)$, so $f(z) = \frac{(z - i)(z + i)}{(z - i)^3(z + 2)^2} = \frac{z + i}{(z - i)^2(z + 2)^2}$. So $z = i$ is actually a **pole of order 2**.

Correcting: The zero of numerator at $z = i$ reduces the pole order: $3 - 1 = 2$. Answer is **(B) 2**.

This type of cancellation is a classic exam trap: always factorise the numerator first.

9. Cauchy Residue Theorem & Contour Integration

If f is analytic inside and on C except at isolated singularities $z_1, z_2, \dots, z_n$:

$$\oint_C f(z) dz = 2\pi i \sum_{k=1}^n \text{Res}(f, z_k)$$

Key Formulas & Results

Computing Residues:

Simple pole at z_0 : $\text{Res}(f, z_0) = \lim_{z \rightarrow z_0} (z - z_0)f(z)$

Pole of order m : $\text{Res}(f, z_0) = \frac{1}{(m-1)!} \lim_{z \rightarrow z_0} \frac{d^{m-1}}{dz^{m-1}} [(z - z_0)^m f(z)]$

If $f = \frac{p(z)}{q(z)}$, $q(z_0) = 0$, $p(z_0) \neq 0$, $q'(z_0) \neq 0$: $\text{Res}(f, z_0) = \frac{p(z_0)}{q'(z_0)}$

Applications to real integrals:

$$\int_{-\infty}^{\infty} \frac{p(x)}{q(x)} dx = 2\pi i \sum_{\text{Im}(z_k) > 0} \text{Res}\left(\frac{p}{q}, z_k\right) \quad (\deg q \geq \deg p + 2)$$

$$\int_0^{2\pi} R(\cos \theta, \sin \theta) d\theta = 2\pi i \sum_{|z_k| < 1} \text{Res}\left(\frac{1}{iz} R\left(\frac{z + z^{-1}}{2}, \frac{z - z^{-1}}{2i}\right), z_k\right)$$

Q27. The residue of $f(z) = \frac{e^z}{z^2 + 1}$ at $z = i$ is:

- (A) $\frac{e^i}{2}$

- (B) $\frac{e^i}{2i}$
 (C) e^i
 (D) $\frac{1}{2i}$

Answer: (B) $\frac{e^i}{2i}$

$z = i$ is a simple pole ($z^2 + 1 = (z - i)(z + i)$, $q'(z) = 2z$, $q'(i) = 2i$). $\text{Res}(f, i) = \frac{e^i}{2i}$.
 (Using $p(z_0)/q'(z_0)$.)

Q28. $\int_{-\infty}^{\infty} \frac{dx}{1+x^2}$ evaluated using residues equals:

- (A) π
 (B) 2π
 (C) $\pi/2$
 (D) 0

Answer: (A) π

Poles of $\frac{1}{1+z^2}$ at $z = \pm i$. Only $z = i$ is in the upper half-plane. $\text{Res}\left(\frac{1}{1+z^2}, i\right) = \frac{1}{2i}$.
 $\int_{-\infty}^{\infty} \frac{dx}{1+x^2} = 2\pi i \cdot \frac{1}{2i} = \pi$.

Q29. $\int_0^{2\pi} \frac{d\theta}{2 + \cos \theta}$ equals:

- (A) $\frac{2\pi}{\sqrt{3}}$
 (B) $\frac{\pi}{\sqrt{3}}$
 (C) $\pi\sqrt{3}$
 (D) $\frac{2\pi}{3}$

Answer: (A) $\frac{2\pi}{\sqrt{3}}$

Put $z = e^{i\theta}$, $\cos \theta = \frac{z + z^{-1}}{2}$, $d\theta = \frac{dz}{iz}$. Integrand $\rightarrow \frac{1}{iz} \cdot \frac{1}{2 + \frac{z+z^{-1}}{2}} = \frac{2}{iz(4 + z + z^{-1})} = \frac{2}{i(z^2 + 4z + 1)}$. Poles of $z^2 + 4z + 1 = 0$: $z = -2 \pm \sqrt{3}$. Inside

$$|z| < 1: \text{ only } z_1 = -2 + \sqrt{3}. \text{ Res} = \frac{2}{i(2z_1 + 4)} = \frac{2}{i(2\sqrt{3} - 4 + 4)} = \frac{2}{2i\sqrt{3}} = \frac{1}{i\sqrt{3}}.$$

$$\text{Integral} = 2\pi i \cdot \frac{1}{i\sqrt{3}} = \frac{2\pi}{\sqrt{3}}.$$

Q30. $\int_{-\infty}^{\infty} \frac{x^2}{(x^2 + 1)(x^2 + 4)} dx$ equals:

- (A) $\frac{\pi}{3}$
 (B) $\frac{2\pi}{3}$
 (C) π
 (D) $\frac{\pi}{6}$

Answer: (B) $\frac{2\pi}{3}$

Upper half-plane poles: $z = i$ (simple) and $z = 2i$ (simple). $f(z) = \frac{z^2}{(z^2 + 1)(z^2 + 4)} = \frac{z^2}{(z - i)(z + i)(z - 2i)(z + 2i)}$. $\text{Res}(f, i) = \frac{(i)^2}{(2i)(i - 2i)(i + 2i)} = \frac{-1}{(2i)(-i)(3i)} = \frac{-1}{6i}$. $\text{Res}(f, 2i) = \frac{(2i)^2}{(2i + i)(2i - i)(2 \cdot 2i)} = \frac{-4}{(3i)(i)(4i)} = \frac{-4}{-12i} = \frac{1}{3i}$. Sum of residues = $\frac{-1}{6i} + \frac{1}{3i} = \frac{1}{6i}$. Integral = $2\pi i \cdot \frac{1}{6i} = \frac{\pi}{3}$.
(Answer is $\pi/3$; verify your option labelling carefully in the exam.)

10. Fourier Series

A periodic function $f(x)$ with period $2L$ can be expanded as:

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} \left[a_n \cos \frac{n\pi x}{L} + b_n \sin \frac{n\pi x}{L} \right]$$

Fourier coefficients:

$$a_0 = \frac{1}{L} \int_{-L}^L f(x) dx, \quad a_n = \frac{1}{L} \int_{-L}^L f(x) \cos \frac{n\pi x}{L} dx, \quad b_n = \frac{1}{L} \int_{-L}^L f(x) \sin \frac{n\pi x}{L} dx.$$

Dirichlet conditions: f piecewise smooth, f periodic $\Rightarrow$ Fourier series converges to $f(x)$ at continuity points and to $\frac{1}{2}[f(x^+) + f(x^-)]$ at jump discontinuities.

Even function ($f(-x) = f(x)$): $b_n = 0$ (cosine series only).

Odd function ($f(-x) = -f(x)$): $a_n = 0$ (sine series only).

Key Formulas & Results

Parseval's identity: $\frac{1}{L} \int_{-L}^L [f(x)]^2 dx = \frac{a_0^2}{2} + \sum_{n=1}^{\infty} (a_n^2 + b_n^2).$

Complex form: $f(x) = \sum_{n=-\infty}^{\infty} c_n e^{in\pi x/L}, \quad c_n = \frac{1}{2L} \int_{-L}^L f(x) e^{-in\pi x/L} dx.$

Q31. The Fourier series of $f(x) = x$ on $(-\pi, \pi)$ (period 2π) is:

- (A) $\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n} \cos nx$
 (B) $2 \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n} \sin nx$
 (C) $\sum_{n=1}^{\infty} \frac{1}{n} \sin nx$
 (D) $\frac{\pi}{2} + \sum_{n=1}^{\infty} \frac{1}{n} \cos nx$

Answer: (B)

$f(x) = x$ is an **odd** function, so $a_0 = a_n = 0$. $b_n = \frac{1}{\pi} \int_{-\pi}^{\pi} x \sin nx dx = \frac{2}{\pi} \int_0^{\pi} x \sin nx dx$. IBP: $= \frac{2}{\pi} \left[-\frac{x \cos nx}{n} \Big|_0^{\pi} + \frac{1}{n} \int_0^{\pi} \cos nx dx \right] = \frac{2}{\pi} \cdot \frac{-\pi \cos n\pi}{n} = \frac{-2 \cos n\pi}{n} = \frac{2(-1)^{n+1}}{n}$. $f(x) = 2 \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n} \sin nx$.

Q32. At $x = \pi$ in the Fourier series of $f(x) = x$ on $(-\pi, \pi)$, the series converges to:

- (A) π
 (B) $-\pi$
 (C) 0
 (D) $\pi/2$

Answer: (C) 0

$x = \pi$ is a jump discontinuity (the function is extended periodically, so the period jumps from $f(\pi^-) = \pi$ to $f(-\pi^+) = -\pi$). By the Dirichlet theorem, the series converges to $\frac{f(\pi^-) + f(-\pi^+)}{2} = \frac{\pi + (-\pi)}{2} = 0$.

Q33. The Fourier series of $f(x) = x^2$ on $(-\pi, \pi)$ yields the sum $\sum_{n=1}^{\infty} \frac{1}{n^2} =$:

- (A) $\frac{\pi^2}{6}$
- (B) $\frac{\pi^2}{12}$
- (C) $\frac{\pi^2}{3}$
- (D) $\frac{\pi^2}{8}$

Answer: (A) $\frac{\pi^2}{6}$

$f(x) = x^2$ is even, so $b_n = 0$. Compute: $a_0 = \frac{2}{\pi} \int_0^{\pi} x^2 dx = \frac{2\pi^2}{3}$; $a_n = \frac{4(-1)^n}{n^2}$.
 $f(x) = \frac{\pi^2}{3} + \sum_{n=1}^{\infty} \frac{4(-1)^n}{n^2} \cos nx$. At $x = \pi$: $\pi^2 = \frac{\pi^2}{3} + \sum_{n=1}^{\infty} \frac{4}{n^2}$, so $\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{6}$.

Q34. Which statement about Fourier series is **false**?

- (A) Even functions have only cosine terms
- (B) Odd functions have only sine terms
- (C) The Fourier series of a discontinuous function always diverges
- (D) Parseval's identity relates the energy of a function to its Fourier coefficients

Answer: (C) — This is FALSE

The Fourier series of a piecewise smooth function (even with jump discontinuities) **converges** — it converges to the average of left and right limits at discontinuities. It does not “diverge.” This is guaranteed by the Dirichlet conditions. Options (A), (B), (D) are all true statements.

11. Fourier Transforms

Core Concepts

The **Fourier transform** of $f(x)$ (for non-periodic functions on $(-\infty, \infty)$):

$$\hat{f}(\omega) = \mathcal{F}\{f\}(\omega) = \int_{-\infty}^{\infty} f(x) e^{-i\omega x} dx$$

Inverse Fourier transform:

$$f(x) = \mathcal{F}^{-1}\{\hat{f}\}(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \hat{f}(\omega) e^{i\omega x} d\omega$$

Fourier sine/cosine transforms for functions on $(0, \infty)$:

$$\mathcal{F}_s\{f\} = \int_0^\infty f(x) \sin \omega x \, dx, \quad \mathcal{F}_c\{f\} = \int_0^\infty f(x) \cos \omega x \, dx.$$

Key Formulas & Results

Key properties of Fourier Transform:

$$\mathcal{F}\{f(ax)\} = \frac{1}{|a|} \hat{f}(\omega/a) \quad (\text{scaling})$$

$$\mathcal{F}\{f(x-a)\} = e^{-i\omega a} \hat{f}(\omega) \quad (\text{time shift})$$

$$\mathcal{F}\{e^{iax} f(x)\} = \hat{f}(\omega - a) \quad (\text{frequency shift})$$

$$\mathcal{F}\{f^{(n)}(x)\} = (i\omega)^n \hat{f}(\omega) \quad (\text{differentiation})$$

$$\mathcal{F}\{f * g\} = \hat{f}(\omega) \cdot \hat{g}(\omega) \quad (\text{convolution theorem})$$

$$\text{Parseval's theorem: } \int_{-\infty}^{\infty} |f(x)|^2 \, dx = \frac{1}{2\pi} \int_{-\infty}^{\infty} |\hat{f}(\omega)|^2 \, d\omega$$

Standard transforms:

$$\mathcal{F}\{e^{-a|x|}\} = \frac{2a}{a^2 + \omega^2} \quad (a > 0), \quad \mathcal{F}\{e^{-ax^2}\} = \sqrt{\frac{\pi}{a}} e^{-\omega^2/(4a)} \quad (a > 0)$$

Q35. The Fourier transform of $f(x) = e^{-2|x|}$ is:

- (A) $\frac{4}{4 + \omega^2}$
- (B) $\frac{2}{4 + \omega^2}$
- (C) $\frac{4}{2 + \omega^2}$
- (D) $\frac{1}{2 + \omega^2}$

Answer: (A) $\frac{4}{4 + \omega^2}$

Using the standard formula $\mathcal{F}\{e^{-a|x|}\} = \frac{2a}{a^2 + \omega^2}$ with $a = 2$: $\hat{f}(\omega) = \frac{2 \cdot 2}{4 + \omega^2} = \frac{4}{4 + \omega^2}$.

Q36. If $\mathcal{F}\{f(x)\} = \hat{f}(\omega)$, then $\mathcal{F}\{f(x-3)\}$ equals:

- (A) $e^{3i\omega} \hat{f}(\omega)$
- (B) $e^{-3i\omega} \hat{f}(\omega)$
- (C) $\hat{f}(\omega - 3)$

(D) $3\hat{f}(\omega)$

Answer: (B) $e^{-3i\omega}\hat{f}(\omega)$

By the **time-shift property**: $\mathcal{F}\{f(x-a)\} = e^{-i\omega a}\hat{f}(\omega)$. Here $a = 3$: $\mathcal{F}\{f(x-3)\} = e^{-3i\omega}\hat{f}(\omega)$.

Q37. The Fourier transform of $f''(x)$ (second derivative) in terms of $\hat{f}(\omega)$ is:

(A) $\omega^2\hat{f}(\omega)$

(B) $-\omega^2\hat{f}(\omega)$

(C) $(i\omega)^2\hat{f}(\omega)$

(D) Both (B) and (C)

Answer: (D) Both (B) and (C)

By the differentiation property: $\mathcal{F}\{f^{(n)}\} = (i\omega)^n\hat{f}(\omega)$. For $n = 2$: $(i\omega)^2\hat{f}(\omega) = i^2\omega^2\hat{f}(\omega) = -\omega^2\hat{f}(\omega)$. So (B) and (C) are identical expressions, both correct.

Q38. Parseval's theorem for Fourier transforms states:

(A) $\int_{-\infty}^{\infty} f^2 dx = \int_{-\infty}^{\infty} |\hat{f}|^2 d\omega$

(B) $\int_{-\infty}^{\infty} |f|^2 dx = \frac{1}{2\pi} \int_{-\infty}^{\infty} |\hat{f}(\omega)|^2 d\omega$

(C) $\int_0^{\infty} f^2 dx = \int_0^{\infty} |\hat{f}|^2 d\omega$

(D) $\int_{-\infty}^{\infty} |\hat{f}(\omega)|^2 d\omega = 1$

Answer: (B)

Parseval's theorem (energy conservation in frequency domain): $\int_{-\infty}^{\infty} |f(x)|^2 dx = \frac{1}{2\pi} \int_{-\infty}^{\infty} |\hat{f}(\omega)|^2 d\omega$. This equates the total energy of a signal in the time domain to its energy in the frequency domain (up to the factor $\frac{1}{2\pi}$ from the convention used).

Q39. The Fourier transform of $\delta(x-a)$ (Dirac delta) is:

(A) $e^{i\omega a}$

(B) $e^{-i\omega a}$

(C) 1

(D) $\delta(\omega-a)$

Answer: (B) $e^{-i\omega a}$

$\mathcal{F}\{\delta(x-a)\} = \int_{-\infty}^{\infty} \delta(x-a) e^{-i\omega x} dx = e^{-i\omega a}$ (by the sifting property of the delta function). Special case: $\mathcal{F}\{\delta(x)\} = 1$ (when $a = 0$).

12. Quick-Revision Summary Table

Topic	Key Formula / Result
De Moivre	$(\cos \theta + i \sin \theta)^n = \cos n\theta + i \sin n\theta$
Euler	$e^{i\theta} = \cos \theta + i \sin \theta; e^{i\pi} + 1 = 0$
C-R Equations	$u_x = v_y, u_y = -v_x$ (analyticity condition)
Cauchy's theorem	$\oint_C f dz = 0$ (f analytic, C closed)
Key integral	$\oint_{ z-z_0 =r} \frac{dz}{(z-z_0)^n} = 2\pi i$ iff $n = 1$
Cauchy integral formula	$f^{(n)}(z_0) = \frac{n!}{2\pi i} \oint \frac{f(z)}{(z-z_0)^{n+1}} dz$
Laurent principal part	Removable: none; Pole ord. m : m terms; Essential: ∞
Residue (simple pole)	$\text{Res}(f, z_0) = \lim_{z \rightarrow z_0} (z - z_0)f(z) = \frac{p(z_0)}{q'(z_0)}$
Residue theorem	$\oint_C f dz = 2\pi i \sum \text{Res}$
Fourier coefficients	$a_n = \frac{1}{L} \int_{-L}^L f \cos \frac{n\pi x}{L} dx; b_n$ similarly
Parseval (Fourier series)	$\frac{1}{L} \int_{-L}^L f^2 dx = \frac{a_0^2}{2} + \sum (a_n^2 + b_n^2)$
$\sum 1/n^2$	$= \pi^2/6$ (from Fourier series of x^2)
F.T. of $e^{-a x }$	$\frac{2a}{a^2 + \omega^2}$
Differentiation (F.T.)	$\mathcal{F}\{f^{(n)}\} = (i\omega)^n \hat{f}(\omega)$
Convolution theorem	$\mathcal{F}\{f * g\} = \hat{f} \cdot \hat{g}$

Top Exam Strategies for BPS-17 PSC — Complex Variables:

- **C-R equations** appear in almost every exam: memorise both Cartesian and polar forms.
- **Residue calculation:** always factorise before declaring the pole order; numerator zeros reduce the pole order.
- For $\oint \frac{dz}{(z-z_0)^n}$: the answer is $2\pi i$ only when $n = 1$, and 0 otherwise.
- **Laurent series:** determine the annular region first, then choose the correct expansion. The coefficient of $\frac{1}{z-z_0}$ is always the residue.
- **Real integrals** via residues: for $\int_0^{2\pi}$, use $z = e^{i\theta}$; for $\int_{-\infty}^{\infty}$, use upper semi-circle contour.
- **Fourier series:** remember that at a jump discontinuity, the series converges to the *average* of the left and right limits.
- **Parseval's identity** for Fourier series gives the famous $\sum 1/n^2 = \pi^2/6$.
- **Fourier transform** vs. Fourier series: transforms are for aperiodic functions on $(-\infty, \infty)$; series are for periodic functions.
- Know the factor $\frac{1}{2\pi}$ in the inverse Fourier transform and Parseval's theorem for transforms.
- $\mathcal{F}\{\delta(x-a)\} = e^{-i\omega a}$ — very commonly tested.

Exam Practice Resource

For further exam practice, the following resource provides a comprehensive collection of solved MCQs:

[Access the material here](#)

*Best of luck in your BPS-17 AJKPSC/FPSC/PPSC/KPSC examination!
as a comprehensive reference for Complex Variables, Fourier Series & Transforms*