

Care Guidance Summary for PT-97BCFE

Overview

This report summarizes the latest diabetes risk review, practical lifestyle advice, medication considerations, and follow-up priorities in patient-friendly language.

Risk Summary

Higher diabetes risk (100%)

Clinical Snapshot

Age: 18 | Sex: Male | HbA1c: 6.8% | Fasting glucose: 95 mg/dL | Random glucose: 140 mg/dL | Cholesterol: 190 mg/dL | BMI: 27.5 | Blood pressure: 120/80 mmHg | Smoking: Never | Drinking: Never | Exercise: Light

Key Risk Drivers

Glucose (fasting) (impact +1.00) | BMI (impact +0.71) | Total cholesterol (impact +0.65) | LDL (impact +0.64) | Age (impact +0.64)

Lifestyle Recommendations

Follow a balanced diet with portion control and lower refined sugar intake. | Aim for consistent physical activity on most days of the week. | Keep regular sleep timing and reduce sedentary time during the day.

Medication Considerations

No medication recommendation could be confidently summarized from the current data.

Monitoring Plan

Repeat HbA1c follow-up according to the clinician plan and response to treatment. | Continue glucose monitoring and compare trends against the current baseline. | Track blood pressure during follow-up visits and home monitoring when available. | Prioritize earlier follow-up because the latest prediction suggests elevated diabetes risk.

Next Steps

Follow a balanced diet with portion control and lower refined sugar intake. | No medication recommendation could be confidently summarized from the current data. | Repeat HbA1c follow-up according to the clinician plan and response to treatment.