

Objectives

Develop an tensor autoregressive model for time series.

- Introducing estimators based on projection, least squares, and maximum likelihood.
- Establishing asymptotics in fix and high dimensions.
- Establishing the model selection consistency.

Introduction




Figure 1: Matrix and Tensor-valued time series.

Our tensor autoregressive model:

- Generalizes the vector and matrix model.
- Preserves tensor structures and interpretations.
- Admits a more comprehensive modeling framework with multiple terms and multiple lagged terms.

Tensor Autoregressive Models

Consider tensor time series $\{\mathcal{X}_t\}$, where at each time t , an order- K tensor $\mathcal{X}_t \in \mathbb{R}^{d_1 \times d_2 \times \dots \times d_K}$ is observed. We introduce the tensor autoregressive model [TenAR\(1\)](#),

$$\mathcal{X}_t = \sum_{r=1}^R \mathcal{X}_{t-1} \times_1 \mathbf{A}_1^{(r)} \times_2 \dots \times_K \mathbf{A}_K^{(r)} + \mathcal{E}_t,$$

where $\mathbf{A}_k^{(r)} \in \mathbb{R}^{d_k \times d_k}$ are coefficient matrices, and $\mathcal{E}_t \in \mathbb{R}^{d_1 \times d_2 \times \dots \times d_K}$ is a tensor white noise. The model can be extended directly to [TenAR\(p\)](#),

$$\mathcal{X}_t = \sum_{i=1}^p \sum_{r=1}^{R_i} \mathcal{X}_{t-i} \times_1 \mathbf{A}_1^{(ir)} \times_2 \dots \times_K \mathbf{A}_K^{(ir)} + \mathcal{E}_t.$$

Estimation and Asymptotics

- Alternating least squares (LSE):

$$\min_{\mathbf{A}_1, \dots, \mathbf{A}_K} \sum_{t=2}^T \|\mathcal{X}_t - \mathcal{X}_{t-1} \times_1 \mathbf{A}_1 \dots \times_K \mathbf{A}_K\|_F^2.$$

- Gaussian MLE when $\text{Cov}(\text{vec}(\mathcal{E}_t))$ is separable:

$$\text{Cov}(\text{vec}(\mathcal{E}_t)) = \Sigma_K \otimes \Sigma_{K-1} \otimes \dots \otimes \Sigma_1.$$

Theorem: Asymptotics

- Central Limit theorems are established.
 - Convergence rates under high dimensionality: let $\Phi^{(i)} = \sum_{r=1}^{R_i} \mathbf{A}_K^{(ir)} \otimes \dots \otimes \mathbf{A}_1^{(ir)}$, and $\hat{\Phi}^{(i)} = \sum_{r=1}^{R_i} \hat{\mathbf{A}}_K^{(ir)} \otimes \dots \otimes \hat{\mathbf{A}}_1^{(ir)}$ be its corresponding estimator, constructed using the LSE.
 - Assume the error tensors \mathcal{E}_t are IID sub-Gaussian.
 - Assume $d \log d/T \rightarrow 0$, where $d = d_1 d_2 \dots d_K$.
- $$\|\hat{\Phi}^{(i)} - \Phi^{(i)}\| = O_p(\sqrt{d/T}), \quad 1 \leq i \leq p.$$

Initialization of the Algorithm

- Initial values of the alternating algorithms (for both LSE and MLE) is critical to find the global minimum.
- Use projection method to initialize the coefficient matrices and Σ_k .
- Projection Method:

$$\min_{\mathbf{A}_1, \dots, \mathbf{A}_K} \|\hat{\Phi} - \mathbf{A}_K \otimes \dots \otimes \mathbf{A}_1\|_F^2$$

where $\hat{\Phi}$ is the estimated coefficient matrix of the VAR model.

$$\min_{\Sigma_1, \dots, \Sigma_K} \|\hat{\Sigma} - \Sigma_K \otimes \dots \otimes \Sigma_1\|_F^2$$

where $\hat{\Sigma}$ is the residual (of VAR) sample covariance matrix.

Model Selection

For given order \tilde{p} and K-ranks $\tilde{\mathbf{R}}_{\tilde{p}} = (\tilde{R}_1, \dots, \tilde{R}_{\tilde{p}})'$, define the information criterion and the estimated K-ranks as

$$\text{IC}(\tilde{\mathbf{R}}_{\tilde{p}}) := \frac{1}{2} \log \left(\frac{1}{dT} \sum_t \left\| \mathcal{X}_t - \sum_{i=1}^{\tilde{p}} \sum_{r=1}^{\tilde{R}_i} \mathcal{X}_{t-i} \times_1 \hat{\mathbf{A}}_1^{(ir)} \times_2 \dots \times_K \hat{\mathbf{A}}_K^{(ir)} \right\|_F^2 \right) + g(d, T) \sum_{i=1}^{\tilde{p}} \tilde{R}_i,$$

$$\hat{\mathbf{R}}_{\hat{p}} := \arg \min_{\tilde{p} \leq P_{\max}, \tilde{R}_1 \leq R_{\max}, \dots, \tilde{R}_{\tilde{p}} \leq R_{\max}} \text{IC}(\tilde{\mathbf{R}}).$$

where $\hat{\mathbf{A}}_k^{(ir)}$ are the estimates obtained under given order and K-ranks.

Theorem: Model Selection

- $g(d, T) \rightarrow 0$ and $\frac{T}{d}g(d, T) \rightarrow \infty$ as $T \rightarrow \infty$.
- Assume $d \log d/T \rightarrow 0$, where $d = d_1 d_2 \dots d_K$.

Then we can select the right K-ranks consistently,

$$\lim_{T \rightarrow \infty} P(\hat{\mathbf{R}}_{\hat{p}} = \mathbf{R}_p) = 1.$$

Numerical Results

- Fama-French portfolio monthly returns: $2 \times 4 \times 4$ portfolio returns formed on Size, Book-to-Market and Operating Profitability.
- Manhattan taxi traffic data: number of rides moving among five zones within 7am to 7pm in a day, formed a $5 \times 5 \times 12$ tensor.

LSE	MLE	iAR	VAR	MEAN	TOTAL
36.63	36.49	36.91	38.76	36.96	37.48

Table 1: Mean square errors of rolling forecast on Fama-French data. Starting from 2001 Jan to 2018 Dec.

LSE	MLE	iAR	VAR	MEAN	TOTAL
33.86	34.80	53.94	41.06	76.68	69.26

Table 2: Mean square errors of rolling forecast on holiday taxi traffic data. Starting from 2018 Jan to 2019 Dec.

- LSE and MLE for taxi data are multi-term $R = 5$ TenAR(1) model; for Fama-French data $R = 1$.
- "iAR" denotes univariate AR(2) model.
- MEAN denotes using mean value as prediction.
- TOTAL denotes mean squares for whole time series,

Conclusion

We have proposed the autoregressive model for tensor-valued time series. Different estimation methods are studied. Asymptotic normality and the convergence rates are established. Information criteria is proposed, and the model selection consistency is established.