

02180: INTRODUCTION TO ARTIFICIAL INTELLIGENCE

LECTURE 11: BELIEF REVISION

Nina Gierasimczuk



RECALL: THE PROBLEM OF BELIEF REVISION

Belief revision is a topic of much interest in theoretical computer science and logic, and it forms a central problem in research into artificial intelligence. In simple terms: how do you update a database of knowledge in the light of new information? What if the new information is in conflict with something that was previously held to be true?

Gärdenfors, Belief Revision

BELIEF SETS

Belief set is a set of formulas that is **deductively closed**.

DEFINITION

For any set B of sentences, $Cn(B)$ is the set of **logical consequences** of B .

If φ can be derived from B by classical propositional logic, then $\varphi \in Cn(B)$.

THREE PARTS OF TAKING IN NEW INFORMATION

What can I do to my belief set?

1. **Revision:** $B * \varphi$; φ is added and other things are removed, so that the resulting new belief set B' is consistent.
2. **Contraction:** $B \div \varphi$; φ is removed from B giving a new belief set B' .
3. **Expansion:** $B + \varphi$; φ is added to B giving a new belief set B' .

AGM \div RATIONALITY POSTULATES OF CONTRACTION

1. **Closure:** $B \div \varphi = Cn(B \div \varphi)$
the outcome is logically closed
2. **Success:** If $\varphi \notin Cn(\emptyset)$, then $\varphi \notin Cn(B \div \varphi)$
the outcome does not contain φ
3. **Inclusion:** $B \div \varphi \subseteq B$
the outcome is a subset of the original set
4. **Vacuity:** If $\varphi \notin Cn(B)$, then $B \div \varphi = B$
if the incoming sentence is not in the original set then there is no effect
5. **Extensionality:** If $\varphi \leftrightarrow \psi \in Cn(\emptyset)$, then $B \div \varphi = B \div \psi$.
the outcomes of contracting with equivalent sentences are the same
6. **Recovery:** $B \subseteq (B \div \varphi) + \varphi$.
contraction leads to the loss of as few previous beliefs as possible
7. **Conjunctive inclusion:** If $\varphi \notin B \div (\varphi \wedge \psi)$, then $B \div (\varphi \wedge \psi) \subseteq B \div \varphi$.
8. **Conjunctive overlap:** $(B \div \varphi) \cap (B \div \psi) \subseteq B \div (\varphi \wedge \psi)$.

AGM* RATIONALITY POSTULATES OF REVISION

1. **Closure:** $B * \varphi = Cn(B * \varphi)$
2. **Success:** $\varphi \in B * \varphi$
3. **Inclusion:** $B * \varphi \subseteq B + \varphi$
4. **Vacuity:** If $\neg\varphi \notin B$, then $B * \varphi = B + \varphi$
5. **Consistency:** $B * \varphi$ is consistent if φ is consistent.
6. **Extensionality:** If $(\varphi \leftrightarrow \psi) \in Cn(\emptyset)$, then $B * \varphi = B * \psi$.
7. **Superexpansion:** $B * (\varphi \wedge \psi) \subseteq (B * \varphi) + \psi$
8. **Subexpansion:** If $\neg\psi \notin B * \varphi$, then $(B * \varphi) + \psi \subseteq B * (\varphi \wedge \psi)$.

LEVI IDENTITY

One formal way to combine those operations is to use:

LEVI IDENTITY

$$B * \varphi := (B \div \neg\varphi) + \varphi.$$

Belief revision can be defined as first removing any inconsistency with the incoming information and then adding the information itself.

WORKING WITH SETS OF FORMULAS

Obviously, we want $B \div \varphi$ to be a subset of B that does not imply φ .

WORKING WITH SETS OF FORMULAS

Obviously, we want $B \div \varphi$ to be a subset of B that does not imply φ .

But carefully: **limit the removals** from B !

WORKING WITH SETS OF FORMULAS

Obviously, we want $B \div \varphi$ to be a subset of B that does not imply φ .

But carefully: **limit the removals** from B !

DEFINITION

For any set A and sentence φ the **remainder for A and φ** is any set C s.t.:

WORKING WITH SETS OF FORMULAS

Obviously, we want $B \div \varphi$ to be a subset of B that does not imply φ .

But carefully: **limit the removals** from B !

DEFINITION

For any set A and sentence φ the **remainder for A and φ** is any set C s.t.:

- C is a subset of A ,

WORKING WITH SETS OF FORMULAS

Obviously, we want $B \div \varphi$ to be a subset of B that does not imply φ .

But carefully: **limit the removals** from B !

DEFINITION

For any set A and sentence φ the **remainder for A and φ** is any set C s.t.:

- ▶ C is a subset of A ,
- ▶ C does not imply φ , and

WORKING WITH SETS OF FORMULAS

Obviously, we want $B \div \varphi$ to be a subset of B that does not imply φ .

But carefully: **limit the removals** from B !

DEFINITION

For any set A and sentence φ the **remainder for A and φ** is any set C s.t.:

- ▶ C is a subset of A ,
- ▶ C does not imply φ , and
- ▶ there is no set C' such that C' does not imply φ and $C \subset C' \subseteq A$.

CONTRACTION REMAINDERS

DEFINITION

For any set A and sentence φ the **remainder set** $A \perp \varphi$ is the set of inclusion-maximal subsets of A that do not imply φ .

Formally: a set C is an element of $A \perp \varphi$ just in case $C \subset A$, $C \not\models \varphi$, and there is no set C' such that $C' \not\models \varphi$ and $C \subset C' \subseteq A$.

CONTRACTION REMAINDERS

DEFINITION

For any set A and sentence φ the **remainder set** $A \perp \varphi$ is the set of inclusion-maximal subsets of A that do not imply φ .

Formally: a set C is an element of $A \perp \varphi$ just in case $C \subset A$, $C \not\models \varphi$, and there is no set C' such that $C' \not\models \varphi$ and $C \subset C' \subseteq A$.

So, in principle, the result of contraction **could be** chosen to be just one of the remainders, but...

OBJECTION

So, in principle, the result of contraction **could be** chosen to be just one of the remainders...

BUT:

For belief set A and any φ inconsistent with A , A revised with φ will be complete, i.e., for any formula ψ , $\psi \in A * \varphi$ or $\neg\psi \in A * \varphi$.

This means that any agent after revising their belief becomes omniscient.

OBJECTION

So, in principle, the result of contraction **could be** chosen to be just one of the remainders...

BUT:

For belief set A and any φ inconsistent with A , A revised with φ will be complete, i.e., for any formula ψ , $\psi \in A * \varphi$ or $\neg\psi \in A * \varphi$.

This means that any agent after revising their belief becomes omniscient.

Argument:

Assume our agent has a belief set A .

Incoming information is φ , such that $\neg\varphi \in A$.

OBJECTION

So, in principle, the result of contraction **could be** chosen to be just one of the remainders...

BUT:

For belief set A and any φ inconsistent with A , A revised with φ will be complete, i.e., for any formula ψ , $\psi \in A * \varphi$ or $\neg\psi \in A * \varphi$.

This means that any agent after revising their belief becomes omniscient.

Argument:

Assume our agent has a belief set A .

Incoming information is φ , such that $\neg\varphi \in A$.

Then, for every formula ψ both $\neg\varphi \vee \psi$ and $\neg\varphi \vee \neg\psi$ are in A .

Upon contracting $\neg\varphi$ according to the above-described way, for any formula ψ , either ψ or $\neg\psi$ are in $A \div \neg\varphi$.

PARTIAL MEET CONTRACTION

Solution: Let $B \div \varphi$ be the **intersection** of some of the remainders!

DEFINITION

A **selection function** for B is a function γ such that if $B \perp \varphi$ is non-empty, then $\gamma(B \perp \varphi)$ is a non-empty subset of $B \perp \varphi$.

PARTIAL MEET CONTRACTION

Solution: Let $B \div \varphi$ be the **intersection** of some of the remainders!

DEFINITION

A **selection function** for B is a function γ such that if $B \perp \varphi$ is non-empty, then $\gamma(B \perp \varphi)$ is a non-empty subset of $B \perp \varphi$.

The outcome of the **partial meet contraction** is equal to the intersection of the selected elements of $B \perp \varphi$, i.e., $B \div \varphi = \cap \gamma(B \perp \varphi)$.

BELIEF SETS VS BELIEF BASES

In AGM every sentence that follows from the belief set...

BELIEF SETS VS BELIEF BASES

In AGM every sentence that follows from the belief set...is already in it.

DEFINITION

B is a belief set just in case $B = Cn(B)$.

BELIEF SETS VS BELIEF BASES

In AGM every sentence that follows from the belief set...is already in it.

DEFINITION

B is a belief set just in case $B = Cn(B)$.

OBJECTION

Suppose that the belief set contains the sentence p : 'Shakespeare wrote Hamlet'. Due to logical closure it then also contains the sentence $p \vee q$, 'Shakespeare wrote Hamlet or Dickens wrote Hamlet'. The latter sentence is a mere logical consequence that should perhaps have no standing of its own.

BELIEF SETS VS BELIEF BASES

In AGM every sentence that follows from the belief set...is already in it.

DEFINITION

B is a belief set just in case $B = Cn(B)$.

OBJECTION

Suppose that the belief set contains the sentence p : 'Shakespeare wrote Hamlet'. Due to logical closure it then also contains the sentence $p \vee q$, 'Shakespeare wrote Hamlet or Dickens wrote Hamlet'. The latter sentence is a mere logical consequence that should perhaps have no standing of its own.

DISCUSSION:

- ▶ Logical omniscience; belief sets are theories.
- ▶ 'Belief set is not what is believed, but what one is committed to believe' (Levi 1991).
- ▶ Computationally feasible: **belief base** instead of belief set.

BELIEF BASE

DEFINITION

A **belief base** is a set of sentences that is not necessarily deductively closed. Its elements represent beliefs that are held independently of any other belief or set of beliefs.

BELIEF BASE

DEFINITION

A **belief base** is a set of sentences that is not necessarily deductively closed. Its elements represent beliefs that are held independently of any other belief or set of beliefs.

Those elements of the belief set that are not in the belief base are 'merely derived', i.e., they have no independent standing.

BELIEF BASE

DEFINITION

A **belief base** is a set of sentences that is not necessarily deductively closed. Its elements represent beliefs that are held independently of any other belief or set of beliefs.

Those elements of the belief set that are not in the belief base are 'merely derived', i.e., they have no independent standing.

Changes are performed on the belief base. The underlying intuition is that the merely derived beliefs are not worth retaining for their own sake. If one of them loses the support that it had in basic beliefs, then it will be automatically and implicitly discarded.

BELIEF BASES AND BELIEF SETS

For every belief base A , there is a belief set $Cn(A)$ that represents the beliefs according to A .

BELIEF BASES AND BELIEF SETS

For every belief base A , there is a belief set $Cn(A)$ that represents the beliefs according to A .

One belief set can be represented by different belief bases.

BELIEF BASES AND BELIEF SETS

For every belief base A , there is a belief set $Cn(A)$ that represents the beliefs according to A .

One belief set can be represented by different belief bases.

So, belief bases have more expressive power than belief sets.

BELIEF BASES AND BELIEF SETS

For every belief base A , there is a belief set $Cn(A)$ that represents the beliefs according to A .

One belief set can be represented by different belief bases.

So, belief bases have more expressive power than belief sets.

EXAMPLE

Alice's basic beliefs: p and q .

Bob's basic beliefs: p and $p \leftrightarrow q$.

Are Alice's and Bob's beliefs the same?

Go to: www.menti.com and enter code: 4560 3653

EXAMPLE CONTINUED

Alice's basic beliefs: p and q .

Bob's basic beliefs: p and $p \leftrightarrow q$.

EXAMPLE CONTINUED

Alice's basic beliefs: p and q .

Bob's basic beliefs: p and $p \leftrightarrow q$.

Their **belief sets** with respect to p and q are the same.

EXAMPLE CONTINUED

Alice's basic beliefs: p and q .

Bob's basic beliefs: p and $p \leftrightarrow q$.

Their **belief sets** with respect to p and q are the same.

So, Alice's and Bob's beliefs are **statically equivalent**.

EXAMPLE CONTINUED

Alice's basic beliefs: p and q .

Bob's basic beliefs: p and $p \leftrightarrow q$.

Their **belief sets** with respect to p and q are the same.

So, Alice's and Bob's beliefs are **statically equivalent**.

On the other hand, they are not **dynamically equivalent**:

Say Alice and Bob receive and accept the information that p is false.

So, they both revise their belief bases to include the new belief $\neg p$.

EXAMPLE CONTINUED

Alice's basic beliefs: p and q .

Bob's basic beliefs: p and $p \leftrightarrow q$.

Their **belief sets** with respect to p and q are the same.

So, Alice's and Bob's beliefs are **statically equivalent**.

On the other hand, they are not **dynamically equivalent**:

Say Alice and Bob receive and accept the information that p is false.

So, they both revise their belief bases to include the new belief $\neg p$.

After that, Alice has the basic beliefs $\neg p$ and q , whereas Bob has the basic beliefs $\neg p$ and $p \leftrightarrow q$.

EXAMPLE CONTINUED

Alice's basic beliefs: p and q .

Bob's basic beliefs: p and $p \leftrightarrow q$.

Their **belief sets** with respect to p and q are the same.

So, Alice's and Bob's beliefs are **statically equivalent**.

On the other hand, they are not **dynamically equivalent**:

Say Alice and Bob receive and accept the information that p is false.

So, they both revise their belief bases to include the new belief $\neg p$.

After that, Alice has the basic beliefs $\neg p$ and q , whereas Bob has the basic beliefs $\neg p$ and $p \leftrightarrow q$.

Now, their belief sets are no longer the same:

Alice believes that q whereas Bob believes that $\neg q$.

BELIEF BASE CONTRACTION

Partial meet contraction, as defined before, is applicable to belief bases.

BELIEF BASE CONTRACTION

Partial meet contraction, as defined before, is applicable to belief bases.

Note that $A \perp p$ is the set of maximal subsets of A that do not imply p ; it is still not sufficient that they do not contain p .

EXAMPLE

$$\{p, p \wedge q, p \vee q, p \leftrightarrow q\} \perp p =$$

BELIEF BASE CONTRACTION

Partial meet contraction, as defined before, is applicable to belief bases.

Note that $A \perp p$ is the set of maximal subsets of A that do not imply p ; it is still not sufficient that they do not contain p .

EXAMPLE

$$\{p, p \wedge q, p \vee q, p \leftrightarrow q\} \perp p = \{\{p \vee q\}, \{p \leftrightarrow q\}\}$$

BELIEF REVISION ASSIGNMENT: HIGHLIGHTS

- ▶ The assignment is now online.
- ▶ Work with belief bases.
- ▶ Implement your own method for generating a plausibility order and a selection function;
- ▶ Use AGM postulates to test your software.

ENTRENCHMENT

When forced to give up previous beliefs, the epistemic agent should give up beliefs that have as little explanatory power and informational value as possible.

ENTRENCHMENT

When forced to give up previous beliefs, the epistemic agent should give up beliefs that have as little explanatory power and informational value as possible.

E.g., the choice between giving up beliefs in natural laws and beliefs in single factual statements, beliefs in the natural laws, that have much higher explanatory power, should in general be retained.

ENTRENCHMENT

When forced to give up previous beliefs, the epistemic agent should give up beliefs that have as little explanatory power and informational value as possible.

E.g., the choice between giving up beliefs in natural laws and beliefs in single factual statements, beliefs in the natural laws, that have much higher explanatory power, should in general be retained.

Contraction can be ruled by a binary relation of *epistemic entrenchment*.

ENTRENCHMENT

When forced to give up previous beliefs, the epistemic agent should give up beliefs that have as little explanatory power and informational value as possible.

E.g., the choice between giving up beliefs in natural laws and beliefs in single factual statements, beliefs in the natural laws, that have much higher explanatory power, should in general be retained.

Contraction can be ruled by a binary relation of *epistemic entrenchment*.

Say p and q are in the belief set, q **is more entrenched than** p means that q is more useful in inquiry or deliberation, or has more “epistemic value” than p . So, the beliefs with the lowest entrenchment should be the ones that are most readily given up.

ENTRENCHMENT AS AN ORDER

- ▶ $p \leq q$: p is at most as entrenched as q
- ▶ $p < q$: p is less entrenched than q , i.e., $p \leq q$ and $\neg(q \leq p)$
- ▶ $p \leq\!\!=\!\! q$: p and q are equally entrenched, i.e., $p \leq q$ and $q \leq p$

ENTRENCHMENT AS AN ORDER

- ▶ $p \leq q$: p is at most as entrenched as q
- ▶ $p < q$: p is less entrenched than q , i.e., $p \leq q$ and $\neg(q \leq p)$
- ▶ $p \leq \geq q$: p and q are equally entrenched, i.e., $p \leq q$ and $q \leq p$

The rationality postulates for entrenchment:

1. If $p \leq q$ and $q \leq r$, then $p \leq r$ (transitivity);
2. If $p \models q$, then $p \leq q$ (dominance);
3. Either $p \leq (p \wedge q)$ or $q \leq (p \wedge q)$ (conjunctiveness);
4. If the belief set K is consistent, then $p \notin K$ iff $p \leq q$, for all q (minimality);
5. if $q \leq p$ for all q , then $p \in \text{Cn}(\emptyset)$ (maximality).

ENTRENCHMENT-BASED CONTRACTION

An entrenchment relation \leq gives rise to an operation \div of **entrenchment-based contraction** according to the following definition:

$$q \in K \div p \text{ iff } q \in K \text{ and either } p < (p \vee q) \text{ or } p \in \text{Cn}(\emptyset).$$

End of Lecture 11