

# Simple Linear Regression



Dr. Kiah Wah Ong

# Inferences Concerning $\beta_0$ and $\beta_1$ using R

```
← → | 📄 | 💾 Source on Save | 🔍 ✨ | 📄 | ➡ Run | ↺ ➡ | ➡ Source | ⌵
```

```
1 #check working directory
2 getwd()
3 #naming the data as data1
4 data1<-read.csv("Experiment1.CSV", header=TRUE, sep=",")
5 x<-data1$x
6 y<-data1$y
7 show(data1)
8 plot(x,y)
9 model2=lm(y~x)
10 abline(model2)
11 summary(model2)|
```

# Inferences Concerning $\beta_0$ and $\beta_1$ using R



## Inferences Concerning $\beta_0$ and $\beta_1$ using R

```
Call:
lm(formula = y ~ x)

Residuals:
    Min       1Q   Median       3Q      Max
-125.78  -32.71   -2.81   31.49  122.62

Coefficients:
              Estimate Std. Error t value Pr(>|t|)
(Intercept)  18.9446     9.4293    2.009  0.0473 *
x             2.2040     0.1621   13.596 <2e-16 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 46.79 on 98 degrees of freedom
Multiple R-squared:  0.6535,    Adjusted R-squared:  0.65
F-statistic: 184.9 on 1 and 98 DF,  p-value: < 2.2e-16
```

## Inferences Concerning $\beta_0$ and $\beta_1$ using R

```
Call:
lm(formula = y ~ x)

Residuals:
    Min       1Q   Median       3Q      Max
-125.78  -32.71   -2.81   31.49  122.62

Coefficients:
              Estimate Std. Error t value Pr(>|t|)
(Intercept)  18.9446     9.4293   2.009   0.0473 *
x             2.2040     0.1621  13.596  <2e-16 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 46.79 on 98 degrees of freedom
Multiple R-squared:  0.6535,    Adjusted R-squared:  0.65
F-statistic: 184.9 on 1 and 98 DF,  p-value: < 2.2e-16
```

# Inferences Concerning $\beta_0$ and $\beta_1$ using R

```
1 #check working directory
2 getwd()
3 #naming the data as data1
4 data1<-read.csv("Experiment1.CSV", header=TRUE, sep=",")
5 x<-data1$x
6 y<-data1$y
7 show(data1)
8 plot(x,y)
9 model2=lm(y~x)
10 abline(model2)
11 summary(model2)
12 #confidence interval, name of the model, 95% confidence interval)
13 confint(model2, level=0.95)|
```

# Inferences Concerning $\beta_0$ and $\beta_1$ using R

```
Call:
```

```
lm(formula = y ~ x)
```

```
Residuals:
```

Min	1Q	Median	3Q	Max
-125.78	-32.71	-2.81	31.49	122.62

```
Coefficients:
```

	Estimate	Std. Error	t value	Pr(> t )
(Intercept)	18.9446	9.4293	2.009	0.0473 *
x	2.2040	0.1621	13.596	<2e-16 ***

```
---
```

```
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
Residual standard error: 46.79 on 98 degrees of freedom
```

```
Multiple R-squared:  0.6535,    Adjusted R-squared:  0.65
```

```
F-statistic: 184.9 on 1 and 98 DF,  p-value: < 2.2e-16
```

```
> confint(model2, level=0.95)
```

	2.5 %	97.5 %
(Intercept)	0.2323693	37.65683
x	1.8823229	2.52571